

# Supplementary Material for "Leftward, Rightward and Complete Exit Time Distributions of Jump Processes"

J. Klinger and O. Bénichou

*Laboratoire de Physique Théorique de la Matière Condensée,  
CNRS, UPMC, 4 Place Jussieu, 75005 Paris, France*

R. Voituriez

*Laboratoire Jean Perrin, CNRS, UPMC, 4 Place Jussieu, 75005 Paris, France*

(Dated: December 13, 2022)

This Supplementary Material presents details of calculations:

- on the eigenvalues and eigenfunctions of the fractional Laplacian.
- on the small  $x_0$  behavior of the continuous quantities.
- on the rightward exit time probability for the case  $1 \leq \mu < 2$ .
- on the non zero initial condition  $x_0$ .
- on the scaling of the splitting probability  $\pi_{0,x}$ .
- on the small  $\tau$  behavior of  $g_2(\tau)$ .

## I. EIGENVALUES AND EIGENFUNCTIONS OF THE FRACTIONAL DIFFUSION EQUATION

In this section, we reproduce results from<sup>3</sup> regarding the eigenvalues and eigenfunctions of the fractional Laplacian. We emphasize that the Fractional Laplacian is an important technical tool to describe Levy-like processes, but that the eigenfunctions  $\tilde{\phi}_k$  and eigenvalues  $\lambda_k$  of the operator in a bounded interval are not known analytically. Here we present approximates of these quantities ; we closely follow the notations of<sup>3</sup>. Let  $D = (-1, 1)$  be the domain of interest; we aim at finding solutions to the following equation:

$$\left( \frac{d^2}{dx^2} \right)^{\frac{\mu}{2}} \phi(x) = \lambda \phi(x) \quad \text{for } x \in (-1, 1) \quad (\text{S1})$$

with  $\phi(-1) = \phi(1) = 0$ . Denoting  $\lambda_k$  the  $k$ th eigenvalue, sorted in increasing order, we have:

$$\lambda_k = \left( \frac{k\pi}{2} - \frac{(2-\mu)\pi}{8} \right)^\mu + O\left(\frac{1}{k}\right). \quad (\text{S2})$$

Approximate expressions  $\psi_k$  of the corresponding eigenfunctions can be obtained by combining infinite and semi-infinite eigenfunctions of the fractional laplacian, which are known explicitly. It is found that

$$\psi_k(x) = q(-x)F_{\mu_k}(1+x) - (-1)^k q(x)F_{\mu_k}(1-x) \quad (\text{S3})$$

with

$$\mu_k = \frac{k\pi}{2} - \frac{(2-\mu)\pi}{8} \quad (\text{S4})$$

and  $F_\lambda$  defined in the following way:

$$F_\lambda(x) = \sin\left(\lambda x + \frac{(2-\mu)\pi}{8}\right) - G(\lambda x). \quad (\text{S5})$$

Here  $G$  is the laplace transform of a positive function  $\gamma(s)$ :

$$\gamma(s) = \frac{\sqrt{2\mu} \sin\left(\frac{\mu\pi}{2}\right)}{2\pi} \frac{s^\mu}{1 + s^{2\mu} - 2s^\mu \cos\left(\frac{\mu\pi}{2}\right)} \exp\left[\frac{1}{\pi} \int_0^\infty \frac{dr}{1+r^2} \ln\left(\frac{1-r^\mu s^\mu}{1-r^{2\mu} s^{2\mu}}\right)\right]. \quad (\text{S6})$$

Finally  $q$  is an interpolating function:

$$q(x) = \begin{cases} 0 & \text{for } x \in (-\infty, -\frac{1}{3}) \\ \frac{9}{2} \left(x + \frac{1}{3}\right)^2 & \text{for } x \in (-\frac{1}{3}, 0) \\ 1 - \frac{9}{2} \left(x - \frac{1}{3}\right)^2 & \text{for } x \in (0, \frac{1}{3}) \\ 1 & \text{for } x \in (\frac{1}{3}, \infty) \end{cases} \quad (\text{S7})$$

From these combined expressions, one can numerically evaluate the approximates  $\psi_k$ ; this is used to determine numerically  $C$  in equation (8) of the main text.

## II. SMALL $x_0$ BEHAVIOR OF THE CONTINUOUS QUANTITIES

In this section, we show that all continuous quantities considered in the main text, namely  $F_0^{(c)}(n|x_0)$ ,  $F_{0,\underline{x}}^{(c)}(n|x_0)$ ,  $F_{0,x}^{(c)}(n|x_0)$ ,  $\pi_{0,x}^{(c)}(x_0)$ , vanish similarly as  $x_0 \rightarrow 0$ . Denoting here  $\psi_k$  the eigenfunctions of the fractional Laplacian operator on  $[0, x]$  and  $\phi_k$  the eigenfunctions of the fractional Laplacian operator on  $[0, +\infty]$ , Kwasnicki shows that for  $x_0 \rightarrow 0$ :

$$\psi_k(x_0) \propto \phi_k(x_0) \propto \left[ \sqrt{\frac{\mu}{2}} \Gamma\left(\frac{\mu}{2}\right) \right]^{-1} [\mu_k x_0]^{\frac{\mu}{2}}. \quad (\text{S8})$$

Since all of the  $F_0^{(c)}(n|x_0)$ ,  $F_{0,\underline{x}}^{(c)}(n|x_0)$ ,  $F_{0,x}^{(c)}(n|x_0)$ ,  $\pi_{0,x}^{(c)}(x_0)$  can be projected either on the  $\psi_k$  or  $\phi_k$  basis, we obtain that they all vanish as  $x_0^{\frac{\mu}{2}}$  as  $x_0 \rightarrow 0$ .

## III. DETAILS ON THE RIGHTWARD EXIT TIME PROBABILITY FOR THE CASE $1 \leq \mu < 2$

In this section, we focus on the asymptotic behavior of  $F_{0,\underline{x}}(n|0)$  in the case  $1 \leq \mu < 2$ , by performing the same asymptotic expansion as in the main text for the  $\mu < 1$  case.

### A. $\mu = 1$ case

Let us list the various small  $p$  expansions necessary for the analysis, which can be found in<sup>4</sup>, or easily derived:

$$\begin{aligned} \widehat{G}_0(p, \xi|0) &= \frac{1}{\sqrt{1-\xi}} \left[ 1 + \left( \frac{\xi}{1-\xi} \right)^{\frac{1}{\mu}} \frac{a_1}{\pi} p \log(p) + O(p) \right] \\ \widetilde{f}(p) &= \frac{1}{2} + \frac{a_1}{\pi} p \log(p) + O(p) \\ \widetilde{F}(p) &= -\frac{a_1}{\pi} \log(p) + O(1) \end{aligned} \quad (\text{S9})$$

To lowest order in  $p$  one then obtains:

$$\widehat{F}_{0,\underline{x}}(\xi) = -\frac{\xi}{\sqrt{1-\xi}} \frac{a_1}{\pi} \log(p) + o(\log(p)) \quad (\text{S10})$$

which yields, after inversion:

$$F_{0,\underline{x}}(n|0) \underset{x \rightarrow \infty}{\sim} \frac{1}{\pi} q(n-1|0) \left[ \frac{a_1}{x} \right] \quad (\text{S11})$$

in agreement with equation (18) from the main text.

### B. $1 < \mu < 2$ case

Let us repeat this operation:

$$\begin{aligned} \widehat{\widehat{G}}_0(p, \xi|0) &= \frac{1}{\sqrt{1-\xi}} \left[ 1 + \left( \frac{\xi}{1-\xi} \right)^{\frac{1}{\mu}} \frac{a_\mu p}{\sin(\pi/\mu)} - \frac{\xi}{1-\xi} c_\mu (a_\mu p)^\mu + o(p^\mu) \right] \\ \widetilde{f}(p) &= \frac{1}{2} - p \langle f \rangle - c_\mu (a_\mu p)^\mu + o(p^\mu) \\ \widetilde{F}(p) &= \langle f \rangle + c_\mu a_\mu^\mu p^{\mu-1} + o(p^{\mu-1}) \end{aligned} \quad (\text{S12})$$

To lowest order in  $p$  one then obtains:

$$\widehat{\widehat{F}}_{0,\underline{p}}(\xi|0) = \frac{\xi}{\sqrt{1-\xi}} [\langle f \rangle + c_\mu a_\mu^\mu p^{\mu-1}] \quad (\text{S13})$$

which yields, after inversion:

$$F_{0,\underline{x}}(n|0) \underset{x \rightarrow \infty}{\sim} q(n-1|0) \left[ \langle f \rangle \delta(x) + \frac{1}{\pi} \Gamma(\mu) \sin\left(\frac{\pi\mu}{2}\right) \left[ \frac{a_\mu}{x} \right]^\mu \right]. \quad (\text{S14})$$

In the large  $x$  limit, we finally obtain:

$$F_{0,\underline{x}}(n|0) \underset{x \rightarrow \infty}{\sim} q(n-1|0) \frac{1}{\pi} \Gamma(\mu) \sin\left(\frac{\pi\mu}{2}\right) \left[ \frac{a_\mu}{x} \right]^\mu \quad (\text{S15})$$

in agreement with equation (18) of the main text.

## IV. NON ZERO INITIAL CONDITIONS

In this section we focus on the derivation of  $F_{0,\underline{x}}(n|x_0)$  and  $F_{0,x}(n|x_0)$  for  $0 < x_0 \ll a_\mu$ . Note that the assumption  $x_0 \ll a_\mu$  is important since in the opposite limit  $x_0 \gg a_\mu$ , the continuous limit is recovered and the rightward and leftward exit time distributions are known. Our objective here is thus to highlight the peculiar behaviors arising from the discrete time nature of the jump process. Recalling equations (5) and (9) of the main text, we argue that the dependence on the initial position of exit time probabilities is contained either in the splitting probability (rightward FETP) or in the semi infinite first passage time probability (leftward FETP):

$$\begin{aligned} F_{0,\underline{x}}(n|x_0) &\underset{\substack{n \rightarrow \infty \\ x \rightarrow \infty \\ \tau \text{ fixed}}}{\sim} \pi_{0,\underline{x}}(x_0) h_\mu(\tau) n^{-1} \\ F_{0,x}(n|x_0) &\underset{\substack{n \rightarrow \infty \\ x \rightarrow \infty \\ \tau \text{ fixed}}}{\sim} F_0(n|x_0) g_\mu(\tau) \end{aligned} \quad (\text{S16})$$

It was shown in<sup>1,4</sup> that both splitting and semi infinite first passage time probabilities exhibit similar asymptotic forms in the large  $n$  and  $x$  limit. More precisely, for a jump process with jump distribution  $p(l)$  and Fourier Transform  $\tilde{p}(k) = \int_{-\infty}^{\infty} e^{ik\ell} p(\ell) d\ell$  one has:

$$F_0(n|x_0) \underset{n \rightarrow \infty}{\sim} \frac{1}{\sqrt{4n^3}} \left[ \frac{1}{\sqrt{\pi}} + V(x_0) \right] \quad (\text{S17})$$

$$\pi_{0,\underline{x}}(x_0) \underset{x \rightarrow \infty}{\sim} 2^{\mu-1} \Gamma\left(\frac{1+\mu}{2}\right) \left[\frac{a_\mu}{x}\right]^{\frac{\mu}{2}} \left[\frac{1}{\sqrt{\pi}} + V(x_0)\right]$$

where  $a_\mu$  is the scale of the jump process defined by the small  $k$  expansion of  $\tilde{p}(k)$ :

$$\tilde{p}(k) \underset{k \rightarrow 0}{=} 1 - (a_\mu |k|)^\mu + o(k^\mu) \quad (\text{S18})$$

and  $V(x_0)$  is defined by its laplace transform:

$$\begin{aligned} \mathcal{L}V(\lambda) &= \int_0^\infty V(x_0) e^{-\lambda x_0} dx_0 \\ &= \frac{1}{\lambda \sqrt{\pi}} \left( \exp \left[ -\frac{\lambda}{\pi} \int_0^\infty \frac{dk}{k^2 + \lambda^2} \ln(1 - \tilde{p}(k)) \right] - 1 \right) \end{aligned} \quad (\text{S19})$$

Combining equations (S16) and (S17), we thus obtain the small  $x_0 \ll a_\mu$  behavior of the rightward and leftward FETPs, along with the complete exit time distribution  $F_{0,\underline{x}}(n|x_0) = F_{0,\underline{x}}(n|x_0) + F_{0,x}(n|x_0)$ . Agreement with numerical simulations is displayed in figure 1.




Figure 1:  $x_0$  dependence of exit time probabilities. (a) Leftward FETP for a Laplace jump process. After rescaling, the rightward FETP collapses to the  $U(x_0) = \frac{1}{\sqrt{\pi}} + V(x_0)$  function, for various  $\tau$  values. (b) Rescaled rightward FETP for a F-distributed jump process defined by  $p(\ell) \propto \left[ \sqrt{|\ell|}(1 + |\ell|) \right]^{-1}$ . The  $x_0$  dependence is in this case sub linear.

These expressions constitute the extension of the results derived in the main text to the full regime  $0 < x_0 \ll a_\mu$ .

## V. SCALING OF THE SPLITTING PROBABILITY

In this section, we show how to recover the scaling of the splitting probability derived in<sup>2</sup> from equations (5) and (18). Recall first that the splitting probability starting from  $x_0 = 0$  reads:

$$\pi_{0,\underline{x}} = \sum_{k=1}^{\infty} F_{0,\underline{x}}(k). \quad (\text{S20})$$

Splitting this sum into two parts we obtain the following decomposition:

$$\begin{aligned}\pi_{0,\underline{x}} &= \sum_{k=1}^{x^\mu} q(k-1) \frac{\Gamma(\mu)}{\pi} \sin\left(\frac{\pi\mu}{2}\right) \left[\frac{a_\mu}{x}\right]^\mu + \sum_{k=x^\mu}^{\infty} \pi_{0,\underline{x}} \frac{h_\mu(\frac{k}{x^\mu})}{k} \\ &\equiv A_1(x) + A_2(x).\end{aligned}\tag{S21}$$

Since the survival probability satisfies  $q(k) \propto k^{-\frac{1}{2}}$  in the large  $k$  limit, one has  $A_1(x) \propto x^{-\frac{\mu}{2}}$ . In the large  $x$  limit, the second sum can be rewritten as an integral:

$$A_2(x) \sim \pi_{0,\underline{x}} \int_1^\infty h_\mu(\tau) \tau^{-1} d\tau.\tag{S22}$$

Recalling the definition of  $h_\mu(\tau)$  from equation (3) of the main text, we rewrite  $A_2$  as:

$$A_2(x) \sim \pi_{0,\underline{x}} \sum_{k=1}^{\infty} P(k) e^{-\lambda_k 2^\mu}\tag{S23}$$

with  $P(k)$  some sub exponential function of  $k$  that guarantees the convergence of the sum. Summing  $A_1$  and  $A_2$  together, we thus recover the expected scaling of the splitting probability derived in<sup>2</sup>:

$$A_1(x) + A_2(x) \propto x^{-\frac{\mu}{2}}\tag{S24}$$

## VI. SMALL $\tau$ BEHAVIOR OF $g_2(\tau)$

In this section we give an alternate expression of  $g_2(\tau)$  defined in equation (10) of the main text, which is convenient to analyze small  $\tau$  values. We obtain this expression by using the Poisson Sum Formula:

$$\begin{aligned}g_2(\tau) &= 4\pi^{\frac{5}{2}} \tau^{\frac{3}{2}} \sum_{k=1}^{\infty} e^{-k^2 \pi^2 \tau} k^2 \\ &= 1 + \sum_{k=1}^{\infty} e^{-\frac{k^2}{\tau}} \left(2 - \frac{4k^2}{\tau}\right)\end{aligned}\tag{S25}$$

In particular, this yields  $g_2(\tau) \rightarrow 1$  for  $\tau \rightarrow 0$ .

- 
- <sup>1</sup> J. Klinger, R. Voituriez, and O. Bénichou. Distribution of the span of one-dimensional confined random processes before hitting a target. *Physical Review E*, 103(3):032107–, 03 2021.
- <sup>2</sup> J. Klinger, R. Voituriez, and O. Bénichou. Splitting probabilities of symmetric jump processes. *Phys. Rev. Lett.*, 129:140603, Sep 2022.
- <sup>3</sup> Mateusz Kwaśnicki. Eigenvalues of the fractional Laplace operator in the interval. *Journal of Functional Analysis*, 262(5):2379–2402, 2012.
- <sup>4</sup> Satya N. Majumdar, Philippe Mounaix, and Grégory Schehr. Survival probability of random walks and Lévy flights on a semi-infinite line. *Journal of Physics A: Mathematical and Theoretical*, 50(46):465002, nov 2017.