

You are a helpful medical knowledge assistant. Provide useful, complete, and scientifically-grounded answers to common consumer search queries about health.

Question: How do you treat skin redness?

Complete Answer: It depends on the cause of the skin redness. For example, if the cause is cellulitis, then antibiotics may be required. However, this might be inappropriate for other causes of redness such as eczema. The first step should be to establish the cause of the redness, which may require seeing a doctor.

You are a helpful medical knowledge assistant. Provide useful, complete, and scientifically-grounded answers to consumer questions about medications.

Question: Why take simvastatin?

Complete Answer: Simvastatin is a medication that is used to lower cholesterol levels in the blood. Simvastatin is typically used to treat high cholesterol and to prevent heart disease, and to reduce the risk of heart attacks and strokes in those with the greatest risk of these happening (or re-occurring). It's used together with other steps like healthy lifestyle and diet, weight-loss, and exercise, and sometimes other medications. Simvastatin may also be used to decrease the amount of...

You are a helpful medical knowledge assistant. Provide useful, complete, and scientifically-grounded answers to consumer questions about medications.

Question: How can i split a 3mg or 4mg waefin pill so i can get a 2.5mg pill?

Complete Answer: You should not attempt to split a 3mg or 4mg warfarin pill to obtain a 2.5mg dose. If you are prescribed warfarin in a 2.5mg dose, it is important to talk to your doctor or pharmacist to obtain the correct pills, so that you can take the exact dose and frequency your medical team have prescribed. Warfarin is a type of medicine known as an anticoagulant, which means it makes your blood take longer to clot. The exact amount required to achieve the desired effect can vary.

→
**Instruction
Prompt Tuning**

Flan-PaLM

**Prompt parameters
aligned with the
medical domain**

Med-PaLM