

Table 1. The derived parameters for the targets EBLM J0057-19, EBLM J0113+31, EBLM J0123+38, EBLM J0239-20, EBLM J0540-17 and EBLM J0546-18; calculated by our PYCHEOPS fits of CHEOPS and TESS light curves. We also present the spectroscopic surface gravity ($\log g_{1,\text{sp.}}$) and microturbulence velocity v_{mic} calculated by the CHEOPS TS3 team.

	J0057-19	J0113+31	J0123+38	J0239-20	J0540-17	J0546-18
Spectr. params						
$\log g_{1,\text{sp.}}$ (cgs)	4.27 ± 0.10	4.10 ± 0.11	4.30 ± 0.14	4.08 ± 0.20	3.955 ± 0.120	4.0 ± 0.8
v_{mic} (kms $^{-1}$)	1 ± 0	1.07 ± 0.06	1.67 ± 0.10	1 ± 0	1.191 ± 0.060	--
CHEOPS						
Model params						
T_0 (BJD)	1767.3843 ± 0.0002	2335.4973 ± 0.0001	2164.3360 ± 0.0002	2163.7080 ± 0.0001	2209.1209 ± 0.0002	2203.7145 ± 0.0003
P (days)	$= 4.30055$	$= 14.27684$	$= 7.95294$	$= 2.77869$	$= 6.00494$	$= 3.19190$
D	0.0195 ± 0.0004	0.0228 ± 0.0001	0.0300 ± 0.0002	0.0168 ± 0.0002	0.0140 ± 0.0002	0.0239 ± 0.0019
W	0.0365 ± 0.0002	0.0190 ± 0.0001	0.0387 ± 0.0003	0.0527 ± 0.0004	0.0382 ± 0.0002	0.0415 ± 0.0016
b	0.368 ± 0.063	0.320 ± 0.009	0.186 ± 0.095	0.654 ± 0.014	0.167 ± 0.105	0.777 ± 0.041
f_c	$= 0.0$	$0.0871 \pm 0.0002^\ddagger$	$= 0.0$	$= 0.0$	$= 0.0$	$= 0.0$
f_s	$= 0.0$	$-0.5489 \pm 0.0004^\ddagger$	$= 0.0$	$= 0.0$	$= 0.0$	$= 0.0$
L	0.00042 ± 0.00012	0.00083 ± 0.00005	0.00102 ± 0.00008	0.00037 ± 0.00005	0.00037 ± 0.00005	0.00111 ± 0.00013
h_1	0.800 ± 0.020	0.769 ± 0.004	0.818 ± 0.009	0.767 ± 0.020	0.767 ± 0.015	0.440 ± 0.144
h_2	0.502 ± 0.230	0.667 ± 0.093	0.622 ± 0.169	0.469 ± 0.219	0.54 ± 0.18	0.308 ± 0.137
Derived params						
R_2/R_1	0.1395 ± 0.0014	0.1509 ± 0.0003	0.1733 ± 0.0007	0.1296 ± 0.0007	0.1185 ± 0.0009	0.1547 ± 0.0062
R_1/a	0.1064 ± 0.0024	0.0539 ± 0.0003	0.1049 ± 0.0020	0.1797 ± 0.0003	0.1084 ± 0.0018	0.1534 ± 0.0060
R_2/a	0.0143 ± 0.0004	0.0081 ± 0.0001	0.0178 ± 0.0004	0.0229 ± 0.0004	0.0127 ± 0.0003	0.0223 ± 0.0015
i ($^\circ$)	87.75 ± 0.43	89.01 ± 0.03	88.88 ± 0.59	83.25 ± 0.24	88.96 ± 0.67	83.16 ± 0.56
e	0.0	0.3089 ± 0.0004	0.0	0.0	0.0	0.0
ω ($^\circ$)	--	-80.98 ± 0.03	--	--	--	--
Absolute params						
M_2 (M_\odot)	0.1290 ± 0.0052	0.1974 ± 0.0068	0.338 ± 0.012	0.1598 ± 0.0059	0.1633 ± 0.0058	0.2129 ± 0.0075
R_2 (R_\odot)	0.1722 ± 0.0054	0.2161 ± 0.0041	0.3498 ± 0.0097	0.2056 ± 0.0053	0.1939 ± 0.0050	0.2330 ± 0.0130
$\log g_1$ (cgs)	4.257 ± 0.022	4.141 ± 0.010	3.894 ± 0.019	4.053 ± 0.016	4.058 ± 0.017	4.099 ± 0.035
$\log g_2$ (cgs)	5.076 ± 0.021	5.064 ± 0.005	4.883 ± 0.017	5.015 ± 0.014	5.075 ± 0.015	5.028 ± 0.048
$T_{\text{eff},2}$ (K)	2958 ± 124	3258 ± 37	3404 ± 71	3027 ± 58	3220 ± 70	3412 ± 80
TESS						
Model params						
T_0 (BJD)	1338.9354 ± 0.0002	1778.7003 ± 0.0002	1776.6884 ± 0.0005	1413.4614 ± 0.0001	1470.5128 ± 0.0003	2174.9866 ± 0.0003
P (days)	4.30055 ± 0.00001	14.28099 ± 0.00322	7.95267 ± 0.00034	2.77869 ± 0.00001	6.00494 ± 0.00001	3.19192 ± 0.00003
D	0.0185 ± 0.0002	0.0239 ± 0.0002	0.0300 ± 0.0005	0.0166 ± 0.0001	0.0138 ± 0.0002	0.0233 ± 0.0008
W	0.0365 ± 0.0002	0.0191 ± 0.0001	0.0394 ± 0.0002	0.0527 ± 0.0001	0.0383 ± 0.0002	0.0402 ± 0.0005
b	0.376 ± 0.039	0.338 ± 0.017	0.188 ± 0.101	0.644 ± 0.009	0.254 ± 0.089	0.824 ± 0.013
f_c	$= 0.0$	$0.0869 \pm 0.0002^\ddagger$	$= 0.0$	$= 0.0$	$= 0.0$	$= 0.0$
f_s	$= 0.0$	$-0.5489 \pm 0.0004^\ddagger$	$= 0.0$	$= 0.0$	$= 0.0$	$= 0.0$
L	0.00054 ± 0.00008	0.00160 ± 0.00008	0.00261 ± 0.00013	0.00077 ± 0.00004	0.00066 ± 0.00008	0.00176 ± 0.00012
h_1	0.829 ± 0.013	0.820 ± 0.012	0.782 ± 0.018	0.841 ± 0.011	0.811 ± 0.013	0.717 ± 0.101
h_2	0.383 ± 0.278	0.517 ± 0.226	0.521 ± 0.205	0.598 ± 0.193	0.466 ± 0.205	0.355 ± 0.244
Derived params						
R_2/R_1	0.1361 ± 0.0009	0.1546 ± 0.0008	0.1733 ± 0.0015	0.1288 ± 0.0003	0.1175 ± 0.0007	0.1526 ± 0.0028
R_1/a	0.1069 ± 0.0017	0.0545 ± 0.0006	0.1069 ± 0.0017	0.1787 ± 0.0014	0.1105 ± 0.0023	0.1568 ± 0.0027
R_2/a	0.0143 ± 0.0003	0.0083 ± 0.0001	0.0182 ± 0.0004	0.0228 ± 0.0002	0.0126 ± 0.0003	0.0236 ± 0.0003
i ($^\circ$)	87.69 ± 0.28	88.94 ± 0.06	88.85 ± 0.64	83.39 ± 0.14	88.39 ± 0.60	82.58 ± 0.23
e	0.0	0.3088 ± 0.0004	0.0	0.0	0.0	0.0
ω ($^\circ$)	--	-81.00 ± 0.02	--	--	--	--
Absolute params						
M_2 (M_\odot)	0.1289 ± 0.0052	0.1975 ± 0.0068	0.338 ± 0.012	0.1597 ± 0.0059	0.1633 ± 0.0057	0.2132 ± 0.0075
R_2 (R_\odot)	0.1689 ± 0.0045	0.2236 ± 0.0049	0.3566 ± 0.0094	0.2032 ± 0.0043	0.1959 ± 0.0056	0.2356 ± 0.0074
$\log g_1$ (cgs)	4.252 ± 0.017	4.132 ± 0.013	3.877 ± 0.017	4.057 ± 0.011	4.042 ± 0.020	4.080 ± 0.018
$\log g_2$ (cgs)	5.093 ± 0.015	5.035 ± 0.010	4.866 ± 0.016	5.025 ± 0.007	5.066 ± 0.019	5.021 ± 0.022
$T_{\text{eff},2}$ (K)	2750 ± 77	3227 ± 38	3544 ± 63	3014 ± 41	3142 ± 66	3331 ± 56

‡ : Derived parameters based on Gaussian priors from spectroscopic or radial velocity data.

Table 2. The derived parameters for the *CHEOPS* targets EBLM J0719+25, EBLM J0941-31, EBLM J0955-39, EBLM J1013+01, EBLM J1305-31 and EBLM J1522+42; calculated by our *PYCHEOPS* fits of *CHEOPS* and *TESS* light curves. We also present the spectroscopic surface gravity ($\log g_{1,\text{sp.}}$) and microturbulence velocity v_{mic} calculated by the *CHEOPS* TS3 team.

	J0719+25	J0941-31	J0955-39	J1013+01	J1305-31	J1522+42
Spectr. parameters						
$\log g_{1,\text{spectr.}}$ (cgs)	4.35 ± 0.11	3.948 ± 0.191	4.5 ± 0.8	3.56 ± 0.80	4.083 ± 0.109	4.23 ± 0.11
v_{mic} (kms $^{-1}$)	1.12 ± 0.04	1.701 ± 0.108	--	--	1.154 ± 0.031	0.885 ± 0.028
CHEOPS						
Model params						
T_0 (BJD)	2216.3900 ± 0.0002	2278.9545 ± 0.0002	2290.7348 ± 0.0006	2244.3407 ± 0.0002	2311.3193 ± 0.0002	$= 2317.3518$
P (days)	$= 7.45629$	$= 5.54563$	$= 5.31360$	$= 2.89227$	$= 10.61913$	$= 7.66134$
D	0.0213 ± 0.0003	0.0180 ± 0.0003	0.0450 ± 0.0008	0.0432 ± 0.0007	0.0405 ± 0.0006	$= 0.0197$
W	0.0250 ± 0.0003	0.0407 ± 0.0001	0.0270 ± 0.0003	0.0431 ± 0.0002	0.0206 ± 0.0001	$= 0.0285$
b	0.502 ± 0.031	0.343 ± 0.062	0.489 ± 0.034	0.082 ± 0.065	0.704 ± 0.012	$= 0.148$
f_c	$-0.2589 \pm 0.0068^\ddagger$	$0.4436 \pm 0.0012^\ddagger$	$= 0.0$	$= 0.0$	$-0.1725 \pm 0.0014^\ddagger$	$-0.2184 \pm 0.0053^\ddagger$
f_s	$-0.1163 \pm 0.0228^\ddagger$	$0.0390 \pm 0.0040^\ddagger$	$= 0.0$	$= 0.0$	$-0.0859 \pm 0.0024^\ddagger$	$-0.2767 \pm 0.0137^\ddagger$
L	0.00064 ± 0.00012	0.00063 ± 0.00007	0.00153 ± 0.00026	0.00162 ± 0.00016	0.00104 ± 0.00014	0.00053 ± 0.00011
h_1	$0.749 \pm 0.009^\ddagger$	0.829 ± 0.017	0.808 ± 0.034	0.710 ± 0.017	0.785 ± 0.056	--
h_2	$0.408 \pm 0.051^\ddagger$	0.598 ± 0.193	0.397 ± 0.273	0.384 ± 0.192	0.344 ± 0.281	--
Derived params						
R_2/R_1	0.1460 ± 0.0010	0.1341 ± 0.0010	0.2121 ± 0.0019	0.2079 ± 0.0017	0.2013 ± 0.0016	--
R_1/a	0.0763 ± 0.0018	0.1183 ± 0.0021	0.0764 ± 0.0015	0.1124 ± 0.0009	0.0665 ± 0.0008	--
R_2/a	0.0108 ± 0.0003	0.0155 ± 0.0004	0.0158 ± 0.0004	0.0232 ± 0.0002	0.0133 ± 0.0001	--
i ($^\circ$)	87.81 ± 0.19	87.67 ± 0.46	87.86 ± 0.18	89.47 ± 0.42	87.32 ± 0.07	--
e	0.0808 ± 0.0041	0.1984 ± 0.0010	0.0	0.0	0.0371 ± 0.0006	--
ω ($^\circ$)	-155.80 ± 4.57	5.03 ± 0.52	--	--	-153.53 ± 0.69	--
Absolute params						
M_2 (M_\odot)	0.1584 ± 0.0055	0.2173 ± 0.0078	0.2211 ± 0.0080	0.1706 ± 0.0062	0.2820 ± 0.0095	--
R_2 (R_\odot)	0.1919 ± 0.0057	0.2340 ± 0.0064	0.2325 ± 0.0060	0.2093 ± 0.0045	0.3007 ± 0.0072	--
$\log g_1$ (cgs)	4.233 ± 0.022	4.027 ± 0.018	4.432 ± 0.019	4.424 ± 0.011	4.116 ± 0.014	$= 4.168 \pm 0.014$
$\log g_2$ (cgs)	5.071 ± 0.021	5.037 ± 0.017	5.048 ± 0.019	5.029 ± 0.010	4.932 ± 0.013	$= 5.093 \pm 0.013$
$T_{\text{eff},2}$ (K)	3212 ± 89	3448 ± 72	3332 ± 90	3043 ± 49	3135 ± 64	3064 ± 86
TESS						
Model params						
T_0 (BJD)	2559.3826 ± 0.0002	1519.2002 ± 0.0001	1546.8317 ± 0.0002	1518.3782 ± 0.0001	1578.5980 ± 0.0003	1934.2851 ± 0.0002
P (days)	7.45629 ± 0.00005	5.54565 ± 0.00001	5.31359 ± 0.00001	2.89228 ± 0.00001	10.61914 ± 0.00001	7.66134 ± 0.00001
D	0.0209 ± 0.0002	0.0179 ± 0.0001	0.0459 ± 0.0004	0.0417 ± 0.0002	0.0420 ± 0.0004	0.0197 ± 0.0002
W	0.0250 ± 0.0002	0.0410 ± 0.0001	0.0269 ± 0.0001	0.0425 ± 0.0001	0.0205 ± 0.0001	0.0285 ± 0.0002
b	0.520 ± 0.016	0.382 ± 0.022	0.471 ± 0.016	0.106 ± 0.046	0.672 ± 0.012	0.148 ± 0.075
f_c	$-0.2588 \pm 0.0054^\ddagger$	$0.4440 \pm 0.0005^\ddagger$	$= 0.0$	$= 0.0$	$-0.1742 \pm 0.0011^\ddagger$	$-0.2205 \pm 0.0042^\ddagger$
f_s	$-0.1385 \pm 0.0225^\ddagger$	$0.0389 \pm 0.0039^\ddagger$	$= 0.0$	$= 0.0$	$-0.0868 \pm 0.0024^\ddagger$	$-0.2751 \pm 0.0115^\ddagger$
L	0.00093 ± 0.00006	0.00121 ± 0.00005	0.00284 ± 0.00015	0.00292 ± 0.00004	0.00104 ± 0.00014	0.00112 ± 0.00006
h_1	0.812 ± 0.013	0.826 ± 0.007	0.828 ± 0.016	0.795 ± 0.004	0.836 ± 0.038	0.768 ± 0.009
h_2	0.553 ± 0.195	0.558 ± 0.184	0.492 ± 0.206	$= 0.420$	0.505 ± 0.203	0.584 ± 0.150
Derived params						
R_2/R_1	0.1446 ± 0.0006	0.1338 ± 0.0004	0.2141 ± 0.0010	0.2041 ± 0.0004	0.2050 ± 0.0009	0.1404 ± 0.0007
R_1/a	0.0769 ± 0.0010	0.1206 ± 0.0010	0.0756 ± 0.0006	0.1114 ± 0.0004	0.0644 ± 0.0006	0.0792 ± 0.0010
R_2/a	0.0110 ± 0.0002	0.0160 ± 0.0002	0.0160 ± 0.0002	0.0226 ± 0.0001	0.0130 ± 0.0002	0.0110 ± 0.0002
i ($^\circ$)	87.71 ± 0.10	87.36 ± 0.18	87.96 ± 0.09	89.32 ± 0.29	87.52 ± 0.06	89.33 ± 0.346
e	0.0862 ± 0.0036	0.1987 ± 0.0004	0.0	0.0	0.0379 ± 0.0005	0.1243 ± 0.0045
ω ($^\circ$)	-151.86 ± 4.35	5.00 ± 0.51	--	--	-153.53 ± 0.67	-128.70 ± 1.70
Absolute params						
M_2 (M_\odot)	0.1583 ± 0.0055	0.2173 ± 0.0078	0.2220 ± 0.0079	0.1712 ± 0.0062	0.2820 ± 0.0094	0.1657 ± 0.0063
R_2 (R_\odot)	0.1917 ± 0.0044	0.2380 ± 0.0050	0.2327 ± 0.0048	0.2042 ± 0.0039	0.2963 ± 0.0063	0.1915 ± 0.0043
$\log g_1$ (cgs)	4.225 ± 0.014	4.010 ± 0.012	4.442 ± 0.012	4.433 ± 0.010	4.144 ± 0.012	4.168 ± 0.014
$\log g_2$ (cgs)	5.072 ± 0.012	5.022 ± 0.008	5.049 ± 0.008	5.052 ± 0.004	4.945 ± 0.009	5.093 ± 0.013
$T_{\text{eff},2}$ (K)	3063 ± 40	3421 ± 47	3285 ± 41	3017 ± 30	3174 ± 44	3073 ± 36

†: Derived parameters based on Gaussian priors from spectroscopic or radial velocity data.

Table 3. The derived parameters for the *CHEOPS* targets EBLM J1559-05, EBLM J1741+31, EBLM J1928-38, EBLM J1934-42, EBLM J2040-41 and EBLM J2046-40; calculated by our PYCHEOPS fits of *CHEOPS* and *TESS* light curves. We also present the spectroscopic surface gravity ($\log g_{1,\text{sp.}}$) and microturbulence velocity v_{mic} calculated by the *CHEOPS* TS3 team.

	J1559-05	J1741+31	J1928-38	J1934-42	J2040-41	J2046-40
Spectr. params						
$\log g_{1,\text{spectr.}}$ (cgs)	4.37 ± 0.10	4.63 ± 0.11	4.015 ± 0.102	4.327 ± 0.119	4.084 ± 0.103	4.009 ± 0.117
v_{mic} (kms $^{-1}$)	--	1.25 ± 0.05	1.079 ± 0.021	1.099 ± 0.042	1.005 ± 0.035	0.983 ± 0.059
CHEOPS						
Model params						
T_0 (BJD)	2347.2225 ± 0.0001	2014.0489 ± 0.0001	2375.4273 ± 0.0002	2028.2295 ± 0.0002	2433.8913 ± 0.0005	$= 2604.942$
P (days)	$= 3.76008$	$= 7.71262$	$= 23.32286$	$= 6.35251$	$= 14.45626$	$= 37.0143$
D	0.0139 ± 0.0005	0.1446 ± 0.0445	0.0378 ± 0.0004	0.0540 ± 0.0122	0.0177 ± 0.0003	$= 0.0316$
W	0.0429 ± 0.0002	0.0104 ± 0.0027	0.0134 ± 0.0001	0.0191 ± 0.0003	0.0184 ± 0.0003	$= 0.0092$
b	0.701 ± 0.007	1.274 ± 0.116	0.395 ± 0.019	0.812 ± 0.053	0.176 ± 0.118	$= 0.152$
f_c	$= 0.0$	$0.3003 \pm 0.0016^\ddagger$	$-0.1989 \pm 0.0006^\ddagger$	$= 0.0$	$0.3810 \pm 0.0005^\ddagger$	$-0.6279 \pm 0.0002^\ddagger$
f_s	$= 0.0$	$0.4592 \pm 0.0012^\ddagger$	$-0.1840 \pm 0.0007^\ddagger$	$= 0.0$	$-0.2852 \pm 0.0007^\ddagger$	$0.2826 \pm 0.0006^\ddagger$
L	0.00029 ± 0.00002	--	0.00126 ± 0.00008	0.00133 ± 0.00043	0.00032 ± 0.00007	0.00087 ± 0.00007
h_1	0.660 ± 0.075	$0.765 \pm 0.011^\ddagger$	0.732 ± 0.009	$0.730 \pm 0.011^\ddagger$	0.733 ± 0.015	--
h_2	0.080 ± 0.085	$0.429 \pm 0.050^\ddagger$	0.553 ± 0.149	$0.396 \pm 0.050^\ddagger$	0.305 ± 0.197	--
Derived params						
R_2/R_1	0.1177 ± 0.0022	0.380 ± 0.057	0.1945 ± 0.0010	0.2323 ± 0.0247	0.1329 ± 0.0013	--
R_1/a	0.1547 ± 0.0015	0.0610 ± 0.0006	0.0373 ± 0.0003	0.0646 ± 0.0018	0.0517 ± 0.0016	--
R_2/a	0.0179 ± 0.0004	0.0203 ± 0.0035	0.0072 ± 0.0001	0.0141 ± 0.0021	0.0067 ± 0.0002	--
i ($^\circ$)	83.78 ± 0.12	85.53 ± 0.42	89.16 ± 0.05	86.99 ± 0.29	89.48 ± 0.37	--
e	0.0	0.3009 ± 0.0015	0.0734 ± 0.0004	0.0	0.2265 ± 0.0005	--
ω ($^\circ$)	--	56.81 ± 0.16	-137.24 ± 0.15	--	-36.82 ± 0.07	--
Absolute params						
M_2 (M_\odot)	0.1568 ± 0.0058	0.461 ± 0.015	0.2703 ± 0.0091	0.1960 ± 0.0076	0.1524 ± 0.0053	--
R_2 (R_\odot)	0.2011 ± 0.0058	0.450 ± 0.068	0.2692 ± 0.0057	0.239 ± 0.026	0.1797 ± 0.0065	--
$\log g_1$ (cgs)	4.024 ± 0.012	4.367 ± 0.013	4.153 ± 0.012	4.471 ± 0.027	4.174 ± 0.028	$= 4.273 \pm 0.011$
$\log g_2$ (cgs)	5.025 ± 0.019	4.790 ± 0.130	5.010 ± 0.009	4.997 ± 0.096	5.111 ± 0.028	$= 5.032 \pm 0.008$
$T_{\text{eff},2}$ (K)	3139 ± 50	--	3153 ± 36	3014 ± 98	2910 ± 89	3127 ± 43
TESS						
Model parameters						
T_0 (BJD)	--	1990.9112 ± 0.0001	--	1659.7838 ± 0.0002	2043.5742 ± 0.0006	1346.4552 ± 0.0003
P (days)	--	7.71264 ± 0.00001	--	6.35251 ± 0.00001	14.45534 ± 0.00083	37.01362 ± 0.00002
D	--	0.0986 ± 0.0020	--	0.0476 ± 0.0011	0.0176 ± 0.0002	0.0316 ± 0.0003
W	--	0.0127 ± 0.0002	--	0.0190 ± 0.0001	0.0187 ± 0.0001	0.0092 ± 0.0001
b	--	1.141 ± 0.011	--	0.787 ± 0.009	0.124 ± 0.090	0.152 ± 0.111
f_c	--	$0.3005 \pm 0.0020^\ddagger$	--	$= 0.0$	$0.3812 \pm 0.0005^\ddagger$	$-0.6276 \pm 0.0003^\ddagger$
f_s	--	$0.4589 \pm 0.0016^\ddagger$	--	$= 0.0$	$-0.2853 \pm 0.0007^\ddagger$	$0.2825 \pm 0.0006^\ddagger$
L	--	--	--	0.00212 ± 0.00028	0.00089 ± 0.00016	0.00194 ± 0.00015
h_1	--	$0.807 \pm 0.012^\ddagger$	--	$0.786 \pm 0.011^\ddagger$	0.780 ± 0.018	0.764 ± 0.013
h_2	--	$0.412 \pm 0.057^\ddagger$	--	$0.389 \pm 0.050^\ddagger$	0.428 ± 0.196	0.518 ± 0.179
Derived params						
R_2/R_1	--	0.3140 ± 0.0032	--	0.2182 ± 0.0024	0.1325 ± 0.0009	0.1778 ± 0.0010
R_1/a	--	0.0612 ± 0.0003	--	0.0641 ± 0.0007	0.0521 ± 0.0008	0.0247 ± 0.0002
R_2/a	--	0.0189 ± 0.0002	--	0.0137 ± 0.0003	0.0068 ± 0.0001	0.0044 ± 0.0001
i ($^\circ$)	--	86.00 ± 0.05	--	87.11 ± 0.06	89.63 ± 0.27	89.78 ± 0.16
e	--	0.3007 ± 0.0019	--	0.0	0.2267 ± 0.0005	0.4737 ± 0.0004
ω ($^\circ$)	--	56.79 ± 0.20	--	--	-36.81 ± 0.08	155.77 ± 0.05
Absolute params						
M_2 (M_\odot)	--	0.460 ± 0.015	--	0.1967 ± 0.0070	0.1524 ± 0.0053	0.1917 ± 0.0067
R_2 (R_\odot)	--	0.3729 ± 0.0083	--	0.2229 ± 0.0051	0.1804 ± 0.0044	0.2212 ± 0.0046
$\log g_1$ (cgs)	--	4.365 ± 0.011	--	4.478 ± 0.013	4.168 ± 0.016	4.273 ± 0.011
$\log g_2$ (cgs)	--	4.958 ± 0.010	--	5.038 ± 0.014	5.108 ± 0.014	5.032 ± 0.008
$T_{\text{eff},2}$ (K)	--	--	--	2959 ± 64	3014 ± 90	3163 ± 50

‡ : Derived parameters based on Gaussian priors from spectroscopic or radial velocity data.

Table 4. The derived parameters for the *CHEOPS* targets EBLM J2046+06, EBLM J2134+19, EBLM J2315+23, EBLM J2343+29 and EBLM J2359+44; calculated by our *PHYCHEOPS* fits of *CHEOPS* and *TESS* light curves. We also present the spectroscopic surface gravity ($\log g_{1,\text{sp.}}$) and microturbulence velocity v_{mic} calculated by the *CHEOPS* TS3 team.

	J2046+06	J2134+19	J2315+23	J2343+29	J2359+44
Spectr. parameters					
$\log g_{1,\text{spectr.}}$ (cgs)	3.984 ± 0.105	4.00 ± 0.11	4.26 ± 0.11	4.25 ± 0.18	4.68 ± 0.12
v_{mic} (kms^{-1})	1.612 ± 0.048	0.76 ± 0.04	1.18 ± 0.04	0.42 ± 0.20	2.10 ± 0.09
CHEOPS					
Model params					
T_0 (BJD)	2090.6246 ± 0.0001	2462.2840 ± 0.0011	2476.1268 ± 0.0002	2458.6659 ± 0.0001	1977.9726 ± 0.0001
P (days)	$= 10.10779$	16.58571 ± 0.00005	$= 9.13105$	$= 16.95353$	$= 11.35602$
D	0.0162 ± 0.0002	0.0410 ± 0.0003	0.0269 ± 0.0003	0.0256 ± 0.0003	0.0300 ± 0.0002
W	0.0261 ± 0.0002	0.0201 ± 0.0002	0.0279 ± 0.0002	0.01070 ± 0.00005	0.0260 ± 0.0001
b	0.254 ± 0.050	0.645 ± 0.013	0.190 ± 0.092	0.290 ± 0.038	0.0970 ± 0.0239
f_c	$-0.1904 \pm 0.0006^\dagger$	$0.4360 \pm 0.0061^\dagger$	$-0.3241 \pm 0.0009^\dagger$	$0.0806 \pm 0.0006^\dagger$	$-0.0532 \pm 0.0003^\dagger$
f_s	$0.5545 \pm 0.0004^\dagger$	$0.2706 \pm 0.0214^\dagger$	$-0.2088 \pm 0.0020^\dagger$	$0.3924 \pm 0.0004^\dagger$	$-0.6890 \pm 0.0007^\dagger$
L	0.00033 ± 0.00002	0.00388 ± 0.00013	0.00097 ± 0.00011	0.00031 ± 0.00005	0.00089 ± 0.00006
h_1	0.761 ± 0.011	0.746 ± 0.009	0.816 ± 0.0103	0.697 ± 0.009	0.776 ± 0.004
h_2	0.319 ± 0.161	0.598 ± 0.132	0.640 ± 0.157	0.377 ± 0.140	0.612 ± 0.131
Derived params					
R_2/R_1	0.1274 ± 0.0007	0.2024 ± 0.0007	0.1641 ± 0.0009	0.1601 ± 0.0010	0.1731 ± 0.0005
R_1/a	0.0747 ± 0.0005	0.0622 ± 0.0008	0.0765 ± 0.0015	0.0299 ± 0.0002	0.0698 ± 0.0003
R_2/a	0.0094 ± 0.0001	0.0124 ± 0.0002	0.0123 ± 0.0003	0.0047 ± 0.0001	0.0120 ± 0.0001
i ($^\circ$)	88.91 ± 0.22	87.70 ± 0.07	89.17 ± 0.42	89.50 ± 0.07	89.61 ± 0.10
e	0.3438 ± 0.0005	0.2633 ± 0.0064	0.1486 ± 0.0007	0.1604 ± 0.0003	0.4776 ± 0.0010
ω ($^\circ$)	108.95 ± 0.05	31.84 ± 2.40	-147.21 ± 0.29	78.39 ± 0.08	-94.41 ± 0.03
Absolute params					
M_2 (M_\odot)	0.1769 ± 0.0062	0.359 ± 0.019	0.2309 ± 0.0099	0.1202 ± 0.0046	0.293 ± 0.010
R_2 (R_\odot)	0.2043 ± 0.0042	0.3706 ± 0.0088	0.2517 ± 0.0068	0.1464 ± 0.0029	0.2965 ± 0.0058
$\log g_1$ (cgs)	4.079 ± 0.011	3.860 ± 0.016	4.095 ± 0.019	4.596 ± 0.011	4.067 ± 0.010
$\log g_2$ (cgs)	5.065 ± 0.008	4.854 ± 0.019	5.000 ± 0.020	5.191 ± 0.008	4.960 ± 0.005
$T_{\text{eff},2}$ (K)	3124 ± 34	3496 ± 35	3298 ± 62	2572 ± 82	3462 ± 45
TESS					
Model params					
T_0 (BJD)	2090.6165 ± 0.0100	$= 2462.2840$	2476.1443 ± 0.0156	$= 2458.6659$	1773.4225 ± 0.0026
P (days)	10.10771 ± 0.00014	$= 16.58558$	9.13069 ± 0.00040	$= 16.95353$	11.35338 ± 0.00264
D	0.0160 ± 0.0001	$= 0.0410$	0.0281 ± 0.0002	$= 0.0256$	0.0302 ± 0.0002
W	0.0259 ± 0.0002	0.0209 ± 0.0003	0.0278 ± 0.0001	$= 0.01070$	0.0260 ± 0.0002
b	0.353 ± 0.048	$= 0.645$	0.058 ± 0.048	$= 0.290$	0.120 ± 0.040
f_c	$-0.1908 \pm 0.0006^\dagger$	$0.4326 \pm 0.0080^\dagger$	$-0.3235 \pm 0.0009^\dagger$	$0.0818 \pm 0.0005^\dagger$	$-0.0524 \pm 0.0005^\dagger$
f_s	$0.5546 \pm 0.0004^\dagger$	$0.2805 \pm 0.0277^\dagger$	$-0.2084 \pm 0.0019^\dagger$	$0.3925 \pm 0.0004^\dagger$	$-0.6890 \pm 0.0007^\dagger$
L	0.00073 ± 0.00005	0.00700 ± 0.00011	0.00165 ± 0.00013	0.00071 ± 0.00005	0.00202 ± 0.00010
h_1	0.811 ± 0.010	--	0.824 ± 0.011	--	0.824 ± 0.009
h_2	0.595 ± 0.170	--	0.557 ± 0.186	--	0.565 ± 0.197
Derived params					
R_2/R_1	0.1267 ± 0.0005	--	0.1676 ± 0.0007	--	0.1739 ± 0.0007
R_1/a	0.0760 ± 0.0006	--	0.0751 ± 0.0004	--	0.0700 ± 0.0006
R_2/a	0.0095 ± 0.0001	--	0.0125 ± 0.0001	--	0.0120 ± 0.0001
i ($^\circ$)	88.46 ± 0.22	--	89.75 ± 0.21	--	89.52 ± 0.16
e	0.3439 ± 0.0005	--	0.1480 ± 0.0007	--	0.4775 ± 0.0010
ω ($^\circ$)	108.98 ± 0.06	--	-147.20 ± 0.28	--	-94.35 ± 0.04
Absolute params					
M_2 (M_\odot)	0.1770 ± 0.0061	--	0.2310 ± 0.0099	--	0.293 ± 0.010
R_2 (R_\odot)	0.2067 ± 0.0042	--	0.2523 ± 0.0059	--	0.2989 ± 0.0064
$\log g_1$ (cgs)	4.064 ± 0.011	$= 3.860 \pm 0.016$	4.111 ± 0.010	$= 4.596 \pm 0.011$	4.064 ± 0.012
$\log g_2$ (cgs)	5.055 ± 0.008	$= 4.854 \pm 0.019$	4.998 ± 0.011	$= 5.191 \pm 0.008$	4.952 ± 0.000
$T_{\text{eff},2}$ (K)	3124 ± 42	3574 ± 42	3188 ± 47	2745 ± 36	3508 ± 42

†: Derived parameters based on Gaussian priors from spectroscopic or radial velocity data.