Cyclopropenylidene $c\text{-}C_3H_2$ Mass: 38

C.A. Nixon
ACS

Photolysis and Photoionization

PRODUCTION:

$C_3H_3 + h\nu \rightarrow c\text{–}C_3H_2 + H$ **R094**

LOSS [a,b,c]	Yield	Lower λ
$c\text{-}C_3H_2 + h\nu \rightarrow l\text{-}C_3H + H$	0.3	266 nm
$\rightarrow c\text{-}C_3H + H$	0.3	274 nm
$\rightarrow C_3 + H_2$	0.4	289 nm

$c\text{-}C_3H_2$ may also convert to different isomers of C_3H_2 by absorption of a photon:

$c\text{-}C_3H_2 + h\nu \rightarrow$ HCCCH (propynylidene)
$\rightarrow$ HCCHC (propenediylidene)
$\rightarrow$ CCHCH (cyclopropyne)
$\rightarrow H_2CCC$ (propadienylidene)

Or become ionized: $c\text{-}C_3H_2 + h\nu \rightarrow c\text{-}C_3H_2^+$

References

[a] Seburg et al. (1997)
[b] Wu et al. (2010)
[C] Hébrard et al. (2013)

Example Ion Reactions

PRODUCTION

Dissociative electron recombination:

$c\text{-}C_3H_3^+ + e^- \rightarrow c\text{-}C_3H_2 + H$ **R101**

LOSS

Dissociative electron recombination (as $c\text{–}C_3H_3^+$):

$e^- + c\text{-}C_3H_3^+ \rightarrow C_2H + {}^3CH_2$ **R102**

$e^- + c\text{-}C_3H_3^+ \rightarrow C_3H + 2\,H$ **R104**

Ion-molecule reactions (as $c\text{–}C_3H_2^+$):

$c\text{-}C_3H_2^+ + CH_4 \rightarrow c\text{-}C_3H_3^+ + CH_3$ **R099**

$c\text{-}C_3H_2^+ + C_2H_6 \rightarrow c\text{-}C_3H_3^+ + C_2H_5$ **R100**

Example Neutral Reactions

PRODUCTION

Collisional isomerization:

$t\text{-}C_3H_2 + H \rightarrow c\text{-}C_3H_2 + H$ **R097**

$l\text{-}C_3H_2 + H \rightarrow c\text{-}C_3H_2 + H$ **R098**

Radical-molecule reactions:

$C_2H_2 + CH \rightarrow c\text{-}C_3H_2 + H$ **R095**

$C_2 + C_3H_8 \rightarrow c\text{-}C_3H_2 + C_2H_6$

LOSS

Radical-molecule reactions:

$c\text{-}C_3H_2 + CH_3 \rightarrow C_2H_3 + C_2H_2$ **R105**

$c\text{-}C_3H_2 + H + M \rightarrow C_3H_3 + M^*$ **R106**

$c\text{-}C_3H_2 + C_2H_3 \rightarrow C_3H_3 + C_2H_2$ **R107**