

# Directed Rectangle-Visibility Graphs have Unbounded Dimension

Kathleen Romanik  
DIMACS Center for Discrete Mathematics\*  
and Theoretical Computer Science  
Rutgers, The State University of New Jersey  
P.O. Box 1179, Piscataway, NJ 08855-1179  
romanik@dimacs.rutgers.edu

## Abstract

Visibility representations of graphs map vertices to sets in Euclidean space and express edges as visibility relations between these sets. One visibility representation in the plane that has been studied is one in which the vertices of the graph map to closed isothetic rectangles and the edges are expressed by horizontal or vertical visibility between the rectangles. Two rectangles are only considered to be visible to one another if there is a non-zero width horizontal or vertical band of sight between them. A graph that can be represented in this way is called a *rectangle-visibility graph*.

A rectangle-visibility graph can be directed by directing all edges towards the positive  $x$  and  $y$  directions, which yields a directed acyclic graph. A directed acyclic graph  $G$  has *dimension*  $d$  if  $d$  is the minimum integer such that the vertices of  $G$  can be ordered by  $d$  linear orderings,  $<_1, \dots, <_d$ , and for vertices  $u$  and  $v$  there is a directed path from  $u$  to  $v$  if and only if  $u <_i v$  for all  $1 \leq i \leq d$ . In this note we show that the dimension of the class of directed rectangle-visibility graphs is unbounded.

## 1 Rectangle-Visibility Graphs

The problem of determining a *visibility representation* of a graph, where the vertices of the graph map to sets in Euclidean space and the edges are expressed as visibility relations between these sets, has been widely studied (see [BETT93] for a survey). One representation in the plane that has been studied [Wis89, DH94] maps each vertex of the graph to a closed isothetic rectangle in  $E^2$  and each edge to a horizontal or vertical band of sight between two rectangles.

More formally, consider an arrangement of closed rectangles in  $E^2$  such that the sides of the rectangles are parallel to the axes and the rectangles are pairwise disjoint except

---

\*DIMACS is an NSF Science and Technology Center, funded under contract STC-88-09648, and also receives support from the New Jersey Commission on Science and Technology.

possibly at their boundaries. Two rectangles  $R_i$  and  $R_j$  are  $\epsilon$ -visible if there is a non-degenerate rectangle  $R_{ij}$  with two opposite sides that are subsets of the boundaries of  $R_i$  and  $R_j$ , and  $R_{ij}$  intersects no other rectangle. Such an arrangement is a *rectangle-visibility representation* a graph  $G = (V, E)$ . A graph admits such a representation provided that the following hold:

- There exists a 1-1 onto correspondence between the rectangles and the vertices in  $V$ .
- Vertices  $v_i$  and  $v_j$  are adjacent in  $G$  if and only if their corresponding rectangles  $R_i$  and  $R_j$  are  $\epsilon$ -visible.

Rectangle-visibility graphs are an extension of *bar-visibility graphs*, which were defined independently by Wismath [Wis85] and Tamassia and Tollis [TT86]. In this representation vertices map to closed, disjoint, horizontal line segments in the plane, and two vertices are adjacent in the graph if and only if their corresponding segments are  $\epsilon$ -visible in the vertical direction.

The class of all bar-visibility graphs was completely characterized by both Wismath [Wis85] and Tamassia and Tollis [TT86]. They independently proved that a graph has a bar-visibility representation if and only if it has a planar embedding such that all cut vertices lie on the external face.

The class of all rectangle-visibility graphs has not been completely characterized. However, Wismath [Wis89] proved that all planar graphs have a rectangle-visibility representation. Also, Dean and Hutchinson [DH94] proved that a complete bipartite graph  $K_{p,q}$  has rectangle-visibility representation if and only if  $p \leq 4$ .

## 2 Dimension of Directed Acyclic Graphs

A directed acyclic graph  $G$  has *dimension*  $d$  if  $d$  is the minimum integer such that the vertices of  $G$  can be ordered by  $d$  linear orderings,  $<_1, \dots, <_d$ , and for vertices  $u$  and  $v$  there is a directed path from  $u$  to  $v$  if and only if  $u <_i v$  for all  $1 \leq i \leq d$  [Tro92]. A class  $\mathcal{G}$  of graphs has dimension  $d$  if  $d$  is the largest dimension of any graph in  $\mathcal{G}$ .

A bar-visibility representation of a graph can be directed by directing all edges towards the positive  $y$  direction. It has been shown ([BT88, RU88]) that any graph with a directed bar-visibility representation has dimension at most two. A rectangle-visibility representation of a graph can be directed by directing all edges towards the positive  $x$  and  $y$  direction, yielding a directed acyclic graph.

Let us denote by  $K_{n,n} - M$  a complete bipartite graph with a perfect matching removed, where  $n$  is the size of both partitions. Note that both partitions must have the same size for there to be a perfect matching. It is well known that the directed  $K_{n,n} - M$ , where all edges are directed from one partition to the other, has dimension  $n$ .

Since a directed acyclic graph can be used to represent a partial order, work done by Rival and Urrutia [RU88, RU92] on representing partially ordered sets by moving convex objects in space is related to our study of the dimension of rectangle-visibility graphs.

### 3 Unbounded Dimension of Rectangle-Visibility Graphs

We now show that the dimension of the class of directed rectangle-visibility graphs is unbounded. We show this by giving a class of graphs  $\mathcal{G} = \{G_n \mid n \geq 1\}$  such that the dimension of  $G_n$  is at least  $n$ , and then giving a directed rectangle-visibility representation of  $G_n$ .

The directed graph  $G_n = (V, E)$  that we construct is similar to  $K_{n,n} - M$ , except that the edges are replaced by directed paths. It has  $2n + 3n(n - 1)$  vertices defined as follows:

$$V = \{a_1, \dots, a_n, e_1, \dots, e_n\} \cup \{b_{i,j}, c_{i,j}, d_{i,j} \mid 1 \leq i, j \leq n, i \neq j\}$$

The vertices  $a_1, \dots, a_n$  and  $e_1, \dots, e_n$  correspond to the two partitions of  $K_{n,n} - M$ . The following is a description of the edges of  $G_n$ :

- Each vertex  $a_i$  is a source and has edges  $\{(a_i, b_{i,j}) \mid j \neq i\}$  coming out of it.
- The  $b_{i,j}, c_{i,j}, d_{i,j}$  vertices are connected by edges  $(b_{i,j}, c_{i,j})$  and  $(c_{i,j}, d_{i,j})$ .
- Each vertex  $e_j$  is a sink and has edges  $\{(d_{i,j}, e_j) \mid j \neq i\}$  going into it.

See Figure 1 for an illustration of the subgraph of  $G_n$  with source vertex  $a_i$ . In graph  $G_n$  each vertex  $a_i$  has a directed path to each  $e_j$ , where  $j \neq i$ , but there is no path from  $a_i$  to  $e_i$ .

**Lemma 3.1** *Graph  $G_n$  has dimension at least  $n$ .*

**Proof:** We consider only the relative order of the  $a_i$  and  $e_i$  vertices. Since there is a directed path from  $a_i$  to  $e_j, j \neq i$ ,  $a_i$  must appear before  $e_j, j \neq i$  (i.e.  $a_i < e_j$ ) in each linear ordering of the vertices. Since there is no path from  $a_i$  to  $e_i$ ,  $e_i$  must appear before  $a_i$  (i.e.  $e_i < a_i$ ) in some linear ordering of the vertices. Consider a linear ordering  $<_l$  in which  $e_i <_l a_i$ . For all other  $a_j, j \neq i$ , we must have  $a_j <_l e_i$ , and for all other  $e_j, j \neq i$ , we must have  $a_i <_l e_j$ . Thus in the ordering  $<_l$ , no other pair  $a_j, e_j$  can be reversed. Since each pair  $a_j, e_j$  must be reversed in some ordering, this requires at least  $n$  linear orderings.

□

We now describe a directed rectangle-visibility representation for  $G_n$ . The rectangles for the  $a_i$  vertices are in a staircase arrangement, as are the rectangles for the  $b_{i,j}$  vertices,




Figure 1: Subgraph of  $G_n$  with source vertex  $a_i$

the  $d_{i,j}$  vertices and the  $e_j$  vertices. All of the  $(a_i, b_{i,j})$  edges and  $(c_{i,j}, d_{i,j})$  edges are horizontal, and the rest of the edges are vertical. Figure 2 illustrates the construction for  $G_4$ . It is easy to see how this construction can be extended for any  $n \geq 1$ . Using this rectangle-visibility representation of  $G_n$  we get the following theorem.

**Theorem 3.1** *The dimension of the class of directed rectangle-visibility graphs is unbounded.*

## References

- [BETT93] Giuseppe Di Battista, Peter Eades, Roberto Tamassia, and Ioannis G. Tollis. Algorithms for Automatic Graph Drawing: An Annotated Bibliography. Technical report, Department of Computer Science, Brown University, 1993.
- [BT88] Giuseppe Di Battista and Roberto Tamassia. Algorithms for Plane Representations of Acyclic Digraphs. *Theoretical Computer Science*, 61:175–198, 1988.
- [DH94] Alice M. Dean and Joan P. Hutchinson. Rectangle-Visibility Representations of Bipartite Graphs. In Roberto Tamassia and Ioannis Tollis, editors, *Graph Drawing '94*, pages 159–166, Princeton, NJ, October 10–12 1994.




Figure 2: Rectangle-visibility representation for graph  $G_n$

- [RU88] Ivan Rival and Jorge Urrutia. Representing Orders by Translating Convex Figures in the Plane. *Order* 4, pages 319–339, 1988.
- [RU92] Ivan Rival and Jorge Urrutia. Representing Orders by Moving Figures in Space. *Discrete Mathematics*, 109:255–263, 1992.
- [Tro92] William T. Trotter. *Combinatorics and Partially Ordered Sets: Dimension Theory*. Johns Hopkins University Press, Baltimore, MD, 1992.
- [TT86] Roberto Tamassia and Ioannis G. Tollis. A Unified Approach to Visibility Representations of Planar Graphs. *Discrete Computational Geometry*, 1:321–341, 1986.
- [Wis85] Stephen K. Wismath. Characterizing Bar Line-of-Sight Graphs. In *Proceedings of the First Annual Symposium on Computational Geometry*, pages 147–152, Baltimore, MD, June 5-7 1985. ACM Press.
- [Wis89] Stephen K. Wismath. *Bar-Representable Visibility Graphs and a Related Flow Problem*. PhD thesis, University of British Columbia, Vancouver, BC, 1989.

