

# Human Mobility, AI assistants, and urban emissions: an insidious triangle

Luca Pappalardo<sup>1</sup>, Matteo Böhm<sup>2</sup>, Giuliano Cornacchia<sup>1,3</sup>, Giovanni Mauro<sup>1,3,4</sup>,  
Dino Pedreschi<sup>3</sup> and Mirco Nanni<sup>1,\*</sup>

<sup>1</sup>*Institute of Information Science and Technologies, National Research Council of Italy (ISTI-CNR), Pisa, Italy*

<sup>2</sup>*Department of Computer, Control and Management Engineering, Sapienza University of Rome, Rome, Italy*

<sup>3</sup>*Department of Computer Science, University of Pisa, Pisa, Italy*

<sup>4</sup>*IMT Lucca, Italy*

## Abstract

Transportation remains a significant contributor to greenhouse gas emissions, with a substantial proportion originating from road transport and passenger travel in particular. Today, the relationship between transportation and urban emissions is even more complex, given the increasingly prevalent role and the pervasiveness of AI-based GPS navigation systems such as Google Maps and TomTom. While these services offer benefits to individual drivers, they can also exacerbate congestion and increase pollution if too many drivers are directed onto the same route. In this article, we provide two examples from our research group that explore the impact of vehicular transportation and mobility-AI-based applications on urban emissions. By conducting realistic simulations and studying the impact of GPS navigation systems on emissions, we provide insights into the potential for mitigating transportation emissions and developing policies that promote sustainable urban mobility. Our examples demonstrate how vehicle-generated emissions can be reduced and how studying the impact of GPS navigation systems on emissions can lead to unexpected findings. Overall, our analysis suggests that it is crucial to consider the impact of emerging technologies on transportation and emissions, and to develop strategies that promote sustainable mobility while ensuring the optimal use of these tools.

## Keywords

Human Mobility, Climate Change, GHG emissions, Social AI

## 1. Introduction

Assessing Greenhouse Gas (GHG) emissions and air pollution is essential to mitigating climate change and promoting human health. Among various sources of emissions, transportation ones significantly increased since 1970, with 11.9% of global GHG emissions in 2016 originating from road transport, 60% of which from passenger travel [1, 2]. Additionally, transportation emissions contribute to non-CO<sub>2</sub> pollutants such as nitrogen oxides, ozone, particulate matter, and volatile organic compounds, significantly impacting climate change and threatening human health [1]. Achieving Sustainable Development Goals (SDGs) by 2030 requires urgent action towards reducing cities' per capita environmental impact [3].

When examining the impact of transportation on urban emissions and welfare, it is essential to consider the impact of Artificial Intelligence (AI) on human mobil-

ity. GPS navigation systems, such as Google Maps and TomTom, have become ubiquitous features in transportation and offer significant benefits to individual drivers. However, they can also create congestion and increased pollution when too many drivers are directed on the same route [4, 5, 6]. The town of Leonia in New Jersey is a notable example of this phenomenon. In 2017, navigation apps repeatedly directed drivers onto Leonia's narrow, hilly streets, causing significant congestion [6]. As a result, the police had to close dozens of streets to non-residents during rush hour periods, effectively taking most of the town out of circulation for popular traffic apps. The unintended consequences of well-intentioned navigation apps pose a significant challenge to transportation planning and require careful consideration in urban policy development.

This article presents examples of how vehicular transportation and mobility-AI-based applications impact urban emissions. In Section 2, we examine the spatial patterns of vehicle-generated emissions and how to design more efficient emission reduction scenarios through realistic simulations. In Section 3, we demonstrate how studying the impact of GPS navigation systems on emissions can reveal unexpected results. By evaluating the impact of mobility and AI-based applications on emissions, we aim to provide insights into the potential for mitigating transportation emissions and developing poli-

*Ital-IA 2023: 3rd National Conference on Artificial Intelligence, organized by CINI, May 29–31, 2023, Pisa, Italy*

\*Corresponding author.

✉ luca.pappalardo@isti.cnr.it (L. Pappalardo);

bohm@diag.uniroma1.it (M. Böhm);

giuliano.cornacchia@phd.unipi.it (G. Cornacchia);

giovanni.mauro@phd.unipi.it (G. Mauro); dino.pedreschi@unipi.it

(D. Pedreschi); mirco.nanni@isti.cnr.it (M. Nanni)



© 2022 Copyright for this paper by its authors. Use permitted under Creative Commons License Attribution 4.0 International (CC BY 4.0).

CEUR Workshop Proceedings (CEUR-WS.org)

cies that promote sustainable urban mobility.

## 2. Understanding urban emissions

Existing methods to measure vehicles' emissions vary widely in their level of detail and generalizability. Some methods rely on small samples of vehicles with high spatio-temporal resolutions but limited generalizability due to their sample size. For example, particulate sensors and portable emissions measurement systems provide accurate emissions measurements in real-world driving conditions but are limited in scope [7, 8, 9]. In contrast, studies using odometer readings provide estimates for entire regions but lack instantaneous speed and acceleration data [10, 11, 12, 13].

GPS traces offer a trade-off between these two extremes, providing high spatio-temporal resolution and the ability to estimate emissions using microscopic models while covering a representative fraction of the vehicle fleet [14]. Several studies have used GPS traces to investigate the relationship between emissions and the urban environment, vehicle miles travelled and fuel consumption, trip rates and travel mode choice, and more [15, 16, 17, 18, 19]. Overall, using GPS traces provides a valuable tool for understanding the impact of vehicles on the environment and implementing strategies to reduce emissions.

Despite the variety of literature, it remains unclear what statistical patterns characterize the distribution of emissions per vehicle and road, how these distributions change over time and space, and how this information can be used to simulate emission reduction scenarios. While studies have reported that the distribution of emissions from on-road remote sensing sites across vehicles is skewed [20, 21], this finding has been questioned due to the limitations of this type of measurement [22].

### 2.1. Emissions patterns

In a recent study [23], we analyse anonymous GPS trajectories describing 423,018 trips from 16,715 private light-duty vehicles moving in Greater London, Rome, and Florence throughout January 2017 to compute vehicles' emissions. The trajectories are produced by onboard GPS devices that automatically turn on when the vehicle starts, transmitting a point every minute to the server via a GPRS connection.

We develop a methodological framework to compute vehicles' emissions from their raw GPS trajectories, and we use a microscopic emissions model that estimates the vehicles' instantaneous emissions of carbon dioxide ( $\text{CO}_2$ ), nitrogen oxides ( $\text{NO}_x$ ), particulate matter (PM), and volatile organic compounds (VOC) from speed, acceleration, and fuel type.

We find that a few "gross polluters" are responsible for a tremendous amount of emissions in all three cities, and most vehicles emit significantly less. Indeed, the distribution of emissions per vehicle is associated with a Gini coefficient higher than 0.55 for all the cities and pollutants. The top 10% of gross polluters in Florence, Rome, and London are responsible for 47.5%, 50.5%, and 38.5% of the total  $\text{CO}_2$  emitted during the month, respectively. The study also finds that the distributions of  $\text{CO}_2$  emissions per vehicle of Rome and Florence are well approximated by a truncated power law, while a stretched exponential well approximates London's distribution. This pattern is consistent for other pollutants as well. Similarly, a few "grossly polluted roads" suffer from a significant quantity of emissions, but most suffer significantly less. The distribution of emissions per road is associated with a Gini coefficient higher than 0.64 for all the cities and pollutants, and a truncated power law well approximates it. Figure 1 shows the entire road network of Greater London, with the emissions on each road normalised by the road length to better highlight the differences between the roads.


### 2.2. Simulation scenarios

We study how electrifying a certain share of vehicles would change the emissions on the three cities' roads. In this setting, even if a vehicle's electrification changes its driver's mobility behaviour, it would not create any emissions.

We find that the electrification of just the top 1% of gross polluters would reduce emissions as much as electrifying 10% of random vehicles. In contrast, the percentage reduction of the overall emissions grows almost linearly when the share of electric vehicles is chosen randomly. A Generalised Logistic Function (GLF) approximates the growth rate when the gross polluters are electrified first. The model gives  $R^2 = 0.99$  for Rome, and similar results hold for Florence. In Greater London, the growth starts slowly: there are fewer vehicles with high emissions levels, and electrifying the most polluting vehicles is slightly less effective in reducing emissions than in the other two cities [23].

Given the increasing importance of remote working [25, 26], we also simulate the impact of a massive shift to remote working on reducing vehicle emissions. We assume that this working style eliminates commuting trips. We then perform a simulation in which a growing share of these commuters become home workers, i.e., they no longer travel between their home and work locations (detected from GPS traces).

We find that emissions reduction is more effective when the home workers are gross polluters: remote working for the top 1% gross polluters leads to the same reduction reached if they were  $\approx 4\%$  random vehicles.



**Figure 1:** (a) CO<sub>2</sub> emissions (expressed as grams per metre of road emitted during January 2017) on the road network of Greater London. The roads are coloured according to the level of emissions in a gradient ranging from yellow (low emission) to red (high emission). The road network was plotted with the Python library OSMnx [24]. (b) Plot, on the log–log scale, showing  $P(X > x)$ , i.e., the complementary cumulative distribution function (CCDF; black dots) of the CO<sub>2</sub> emissions per vehicle, together with the best fit (red curve), in Greater London. (c) Plot, on the log–log scale, showing the CCDF (black dots) of the CO<sub>2</sub> emissions per road, together with the best fit (red curve), in Greater London.

Again, a GLF fits emissions reduction when the gross polluters become home workers. Overall, these results demonstrate that targeting specific profiles of vehicles can significantly improve emission reduction policies.

### 3. Understanding impact of AI

According to preliminary research, the influence of navigation applications on the urban environment is a topic that remains unclear and incomplete, as existing studies produce inconsistent and sporadic outcomes [27, 28]. On the one hand, these apps may contribute to mitigating CO<sub>2</sub> emissions [29]. On the other hand, their usage may increase population exposure to pollution in highly populated regions [30]. Real-time navigation apps provide drivers with optimal routes to reach their destinations, considering the current traffic conditions. However, despite their undeniable practicality, online navigation applications can in principle generate several issues in urban traffic [5, 4]. These apps are usually optimized to minimize individual drivers’ travel time without considering the collective impact of the aggregated drivers’ behaviour on the city’s overall traffic situation. For example, these apps may not factor in whether the recommended routes could handle the additional traffic generated by the app or whether this traffic could pose a risk to safety or lead to further pollution. The impact of

a driver’s routing choice cannot be evaluated in isolation because it depends on the concurrent choices of other drivers in the city. If too many drivers select the same “eco-friendly” route, the route may become congested, reducing its eco-friendliness. Therefore, a better understanding of the impact of individual routing choices on the urban environment is necessary.

In a recent paper [31], we designed a simulative framework – TrafficCO<sub>2</sub> – using the state-of-the-art traffic simulator SUMO [32] to create realistic simulations of traffic under different settings. The simulations were conducted in the city of Milan assuming that vehicles would either adhere to the directions of commercial navigation systems – OpenStreetMap (OSM) and TomTom (TT) – or follow a randomised deviation from the fastest route that emulates the unpredictability and irrationality of human drivers.

We varied the percentage of vehicles of the fleet circulating in Milan that followed a navigation app’s suggestions, in order to assess the impact of the rate of routed vehicles on the urban environment. We found that the greater the number of vehicles following the navigation app’s suggestion, the higher the total CO<sub>2</sub> emissions in the city: blindly following the recommendations of a navigation app, which are optimised from an individual standpoint, can lead to traffic congestion in some areas of the city, thus leading to spatial polarisation which results in increased travel time and emissions. Conversely,

taking “noisy” routes increases the diversity of travelled paths, resulting in a better distribution of traffic on the road network and a decreased travel time and emissions. The fraction of routed vehicles also influenced the spatial distribution of emissions in the city, with more emissions concentrating on the external ring road when more vehicles were routed.

Our results also suggested that introducing randomness into the path generation and suggestion phases could be a solution to avoid suggesting only the optimal paths: route perturbation was beneficial, resulting in shorter travel times and lower emissions in the city.

The study acknowledged that the situation in the real world is more complex, with multiple navigation apps coexisting simultaneously, each with its heuristics and representation of urban reality. The evidence suggests a need for algorithms that can exploit social and collective dimensions while simultaneously meeting individual needs. This challenge requires shifting from an individual to a collaborative and social paradigm, where the choices of non-rational or AI-assisted agents who exploit the system and their impact on the whole society are considered.

## 4. Conclusion

Urban traffic is a complex system where individual satisfaction is intimately bound to collective happiness, e.g., traffic is smooth for an individual if it is so for everyone. A system where individual interests go hand in hand with collective happiness. There is plenty of room for a better understanding of the impact of individual routing choices on the urban environment, as well as for studying how to design platform architectures and routing recommendations that influence citizens’ behaviour towards better aggregated outcomes. The challenge is to turn collective goals into an optimization target, whereby users transparently understand and accept recommendations, which can be individually sub-optimal but produce a more efficient collective outcome. Using (social) norms (such as minimising CO<sub>2</sub> emissions) as targets for collective goals without using them as global constraints might be a way to achieve this goal.

## Acknowledgments

This work has been partially supported by EU project H2020 Humane-AI-net G.A. 952026, EU project H2020 SoBigData++ G.A. 871042, and by the CHIST-ERA grant CHIST-ERA-19-XAI-010, by MUR (grant No. not yet available), FWF (grant No. I 5205), EPSRC (grant No. EP/V055712/1), NCN (grantNo. 2020/02/Y/ST6/00064), ETAg (grant No. SLTAT21096), BNSF (grant No. KII-06-OOO2/5).

## References

- [1] Intergovernmental Panel on Climate Change, Climate Change 2014: Mitigation of Climate Change: Working Group III Contribution to the IPCC Fifth Assessment Report, Cambridge University Press, 2015, p. 599–670. doi:10.1017/CB09781107415416.014.
- [2] H. Ritchie, Sector by sector: where do global greenhouse gas emissions come from?, 2020. Available at <https://ourworldindata.org/ghg-emissions-by-sector>.
- [3] United Nations General Assembly, Transforming our world: the 2030 Agenda for Sustainable Development, Technical Report, 2015. Accessed: 2021-02-23.
- [4] J. Macfarlane, Your navigation app is making traffic unmanageable, *IEEE Spectrum* (2019) 22–27.
- [5] S. Siuhi, J. Mwakalonge, Opportunities and challenges of smart mobile applications in transportation, *Journal of Traffic and Transportation Engineering* 3 (2016) 582–592.
- [6] L. W. Foderaro, Navigation apps are turning quiet neighborhoods into traffic nightmares, *The New York Times* (2017).
- [7] P. deSouza, A. Anjomshoa, F. Duarte, R. Kahn, P. Kumar, C. Ratti, Air quality monitoring using mobile low-cost sensors mounted on trash-trucks: Methods development and lessons learned, *Sustainable Cities and Society* 60 (2020) 102239. doi:<https://doi.org/10.1016/j.scs.2020.102239>.
- [8] H. S. Chong, S. Kwon, Y. Lim, J. Lee, Real-world fuel consumption, gaseous pollutants, and CO<sub>2</sub> emission of light-duty diesel vehicles, *Sustainable Cities and Society* 53 (2020) 101925. doi:<https://doi.org/10.1016/j.scs.2019.101925>.
- [9] J. M. Luján, V. Bermúdez, V. Dolz, J. Monsalve-Serrano, An assessment of the real-world driving gaseous emissions from a euro 6 light-duty diesel vehicle using a portable emissions measurement system (pems), *Atmospheric Environment* 174 (2018) 112 – 121. doi:<https://doi.org/10.1016/j.atmosenv.2017.11.056>.
- [10] T. Chatterton, J. Barnes, R. E. Wilson, J. Anable, S. Cairns, Use of a novel dataset to explore spatial and social variations in car type, size, usage and emissions, *Transportation Research Part D: Transport and Environment* 39 (2015) 151–164. doi:<https://doi.org/10.1016/j.trd.2015.06.003>.
- [11] S. R. Kancharla, G. Ramadurai, Incorporating driving cycle based fuel consumption estimation in green vehicle routing problems, *Sustainable Cities and Society* 40 (2018) 214–221. doi:<https://doi.org/10.1016/j.scs.2018.04.016>.
- [12] A. Choudhary, S. Gokhale, Urban real-world driv-

- ing traffic emissions during interruption and congestion, *Transportation Research Part D: Transport and Environment* 43 (2016) 59–70. doi:<https://doi.org/10.1016/j.trd.2015.12.006>.
- [13] F. Zheng, J. Li, H. van Zuylen, C. Lu, Influence of driver characteristics on emissions and fuel consumption, *Transportation Research Procedia* 27 (2017) 624–631. doi:<https://doi.org/10.1016/j.trpro.2017.12.142>, 20th EURO Working Group on Transportation Meeting, EWGT 2017, 4-6 September 2017, Budapest, Hungary.
- [14] L. Pappalardo, S. Rinzivillo, Z. Qu, D. Pedreschi, F. Giannotti, Understanding the patterns of car travel, *European Physical Journal: Special Topics* 215 (2013). doi:10.1140/epjst/e2013-01715-5.
- [15] M. Nyhan, S. Sobolevsky, C. Kang, P. Robinson, A. Corti, M. Szell, D. Streets, Z. Lu, R. Britter, S. R. Barrett, C. Ratti, Predicting vehicular emissions in high spatial resolution using pervasively measured transportation data and microscopic emissions model, *Atmospheric Environment* 140 (2016) 352 – 363. doi:<https://doi.org/10.1016/j.atmosenv.2016.06.018>.
- [16] J. Liu, K. Han, X. M. Chen, G. P. Ong, Spatial-temporal inference of urban traffic emissions based on taxi trajectories and multi-source urban data, *Transportation Research Part C: Emerging Technologies* 106 (2019) 145 – 165. doi:<https://doi.org/10.1016/j.trc.2019.07.005>.
- [17] C. K. Gatley, L. R. Hutyra, S. Peterson, I. Sue Wing, Urban emissions hotspots: Quantifying vehicle congestion and air pollution using mobile phone gps data, *Environmental Pollution* 229 (2017) 496–504. doi:<https://doi.org/10.1016/j.envpol.2017.05.091>.
- [18] J. Chen, W. Li, H. Zhang, W. Jiang, W. Li, Y. Sui, X. Song, R. Shibasaki, Mining urban sustainable performance: Gps data-based spatio-temporal analysis on on-road braking emission, *Journal of Cleaner Production* 270 (2020) 122489. doi:<https://doi.org/10.1016/j.jclepro.2020.122489>.
- [19] Y. Sui, H. Zhang, X. Song, F. Shao, X. Yu, R. Shibasaki, R. Sun, M. Yuan, C. Wang, S. Li, Y. Li, Gps data in urban online ride-hailing: A comparative analysis on fuel consumption and emissions, *Journal of Cleaner Production* 227 (2019) 495 – 505. doi:<https://doi.org/10.1016/j.jclepro.2019.04.159>.
- [20] P. Guenther, G. Bishop, J. Peterson, D. Stedman, Emissions from 200 000 vehicles: a remote sensing study, *Science of The Total Environment* 146-147 (1994) 297 – 302. doi:[https://doi.org/10.1016/0048-9697\(94\)90249-6](https://doi.org/10.1016/0048-9697(94)90249-6).
- [21] C. Brand, B. Boardman, Taming of the few – the unequal distribution of greenhouse gas emissions from personal travel in the uk, *Energy Policy* 36 (2008) 224–238. doi:<https://doi.org/10.1016/j.enpol.2007.08.016>.
- [22] Y. Huang, N. Surawski, Y. S. Yam, C. Lee, J. Zhou, B. Organ, E. Chan, Re-evaluating effectiveness of vehicle emission control programs targeting high-emitters, *Nature Sustainability* 3 (2020). doi:10.1038/s41893-020-0573-y.
- [23] M. Böhm, M. Nanni, L. Pappalardo, Gross polluters and vehicle emissions reduction, *Nature Sustainability* (2022). doi:10.1038/s41893-022-00903-x.
- [24] G. Boeing, Osmnx: New methods for acquiring, constructing, analyzing, and visualizing complex street networks, *Computers, Environment and Urban Systems* 65 (2017) 126 – 139. doi:<https://doi.org/10.1016/j.compenvurbsys.2017.05.004>.
- [25] L. Vyas, N. Butakhieo, The impact of working from home during covid-19 on work and life domains: an exploratory study on hong kong, *Policy Design and Practice* (2020) 1–18.
- [26] L. Nagel, The influence of the covid-19 pandemic on the digital transformation of work, *International Journal of Sociology and Social Policy* (2020).
- [27] E. Ericsson, H. Larsson, K. Brundell-Freij, Optimizing route choice for lowest fuel consumption – potential effects of a new driver support tool, *Transportation Research Part C: Emerging Technologies* 14 (2006) 369–383. doi:<https://doi.org/10.1016/j.trc.2006.10.001>.
- [28] Z. Samaras, L. Ntziachristos, S. Toffolo, G. Magra, A. Garcia-Castro, C. Valdes, C. Vock, W. Maier, Quantification of the effect of its on co2 emissions from road transportation, *Transportation Research Procedia* 14 (2016) 3139–3148. doi:<https://doi.org/10.1016/j.trpro.2016.05.254>.
- [29] N. Arora, T. Cabannes, S. G. Subramaniam, Y. Li, P. McAfee, M. Nunkesser, C. Osorio, A. Tomkins, I. Tsogsuren, Quantifying the sustainability impact of google maps: A case study of salt lake city, 2021.
- [30] F. Perez-Prada, A. Monzón, C. Valdés, Managing traffic flows for cleaner cities: The role of green navigation systems, *Energies* 10 (2017) 791. doi:10.3390/en10060791.
- [31] G. Cornacchia, M. Böhm, G. Mauro, M. Nanni, D. Pedreschi, L. Pappalardo, How routing strategies impact urban emissions, in: *Proceedings of the 30th International Conference on Advances in Geographic Information Systems, SIGSPATIAL '22*, Association for Computing Machinery, New York, NY, USA, 2022. doi:10.1145/3557915.3560977.
- [32] P. A. Lopez, E. Wiessner, M. Behrisch, L. Bieker-Walz, J. Erdmann, Y.-P. Flotterod, R. Hilbrich, L. Lucken, J. Rummel, P. Wagner, Microscopic traffic simulation using SUMO, 2018.