

Cite this: *Dalton Trans.*, 2024, **53**, 9887Alkyl backbone variations in common β -diketiminate ligands and applications to *N*-heterocyclic silylene chemistry†‡

Connor Bourne, Huanhuan Dong, Katharine McKain, Lena C. Mayer, Aidan P. McKay, David B. Cordes, Alexandra M. Z. Slawin and Andreas Stasch *

We report the extension of the common β -diketimine proligand class, $^{R_{Ar}}nacnacH$ ($HC(RCNAr)_2H$), where R is an alkyl group such as Et or *i*Pr, plus Ph, and Ar is a sterically demanding aryl substituent such as Dip = 2,6-diisopropylphenyl, Dep = 2,6-diethylphenyl, Mes = 2,4,6-trimethylphenyl or mesityl, Xyl = 2,6-dimethylphenyl, *via* one-pot condensation procedures. When a condensation reaction is carried out using the chemical dehydrating agent PPSE (polyphosphoric acid trimethylsilylester), β -diketiminate phosphorus(v) products such as $(^{iPrMes}nacnac)PO_2$ can also be obtained, which can be converted to the respective proligand $^{iPrMes}nacnacH$ *via* alkaline hydrolysis. The $^{R_{Ar}}nacnacH$ proligands can be converted to their alkali metal complexes with common methods and we have found that deprotonation of $^{iPrDip}nacnacH$ is significantly more sluggish than that of related β -diketimines with smaller backbone alkyl groups. The basicity of the $^{R_{Ar}}nacnac^-$ anions can play a role in the success of their salt metathesis chemistry and we have prepared and structurally characterised the $^{EtDip}nacnac$ -derived silicon(II) compounds $(^{EtDip}nacnac)SiBr$ and $(^{EtDip}nacnac)Si$, where $^{EtDip}nacnac^-$ is the deprotonated variant $MeCHC(NDip)CHC(NDip)Et$.

Received 2nd May 2024,
Accepted 20th May 2024

DOI: 10.1039/d4dt01298d

rsc.li/dalton

1. Introduction


β -Diketiminato (BDI) ligands, or “nacnac’s” (Fig. 1) from the acetylacetonone (acacH) analogy, are a highly popular ligand class that has been employed in complexes from all parts of the periodic table and for a wide variety of applications.^{1–16} Their general ease of synthesis, including sterically demanding variants, and their generally robust nature has made these ligands a go-to choice where *N,N'*-chelating monoanionic spectator ligands are desirable. For example, these ligands have been successfully and widely employed for a range of reactive low oxidation state metal complexes from across the periodic table.^{2,9,10,12} Their structure allows the tuning of steric and electronic factors,^{1,12} for example *via* varying the R and Ar groups in the symmetric variants shown in Fig. 1A. Regarding electronic factors, even fully fluorinated β -diketimines have been prepared and assessed.^{17,18} In many fields and applications where steric factors are a dominant influence, *e.g.*, in

s-block chemistry,^{8–10} variation of predominantly the *N*-aryl groups, but also the substituents (R) in the ligand backbone (see Fig. 1A and B) have been focused on. In some cases, small steric changes can effect large differences on product outcomes.¹⁹ Most nacnac variants in the literature employ the readily available parent “acac” backbone fragment, and these are typically cheap and easy to synthesise. Although nacnac’s

EaStCHEM School of Chemistry, University of St Andrews, North Haugh, St Andrews, KY16 9ST, UK. E-mail: as411@st-andrews.ac.uk

†Electronic supplementary information (ESI) available: Experimental section, NMR spectra, X-ray diffraction. CCDC 2352325–2352336. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d4dt01298d>

‡The research data (NMR spectroscopy) supporting this publication can be accessed at <https://doi.org/10.17630/9869f43a-be7d-4c25-8bdf-de7ad99a39b5>.

Fig. 1 $^{R_{Ar}}nacnacH$ species.


generally serve as robust monoanionic spectator ligands, the backbone unit can engage in various side reactions owing to the electrophilic, nucleophilic and CH acidic properties of various parts of the acac-derived backbone, see Fig. 1B.⁵ In addition, other decomposition reactions are possible, including the extrusion of an ArN (*e.g.* DipN) unit from the ligand scaffold.⁵ A range of metal coordination modes other than the most common *N,N*-chelating one (Fig. 1B) can be observed,^{1,5} not unrelated to the structures of the main tautomers of ^{RAr}nacnacH species (Fig. 1C). The delocalised free β -diketiminato unit can form various isomers (*E/Z*) that are accessible within a low range of energies.²⁰ Ligand backbone units that allow wider electronic delocalisation, *e.g.*, those with aryl groups, can induce some non-innocent ligand behaviour and are prone to reduction to a radical dianionic ligand unit in some instances.^{1,4,5} Overall, it is surprising that only few studies have placed more emphasis on the ligand backbone substitution.^{4,5} Other common backbone modifications are, for example, *t*Bu₂-backbone substituted systems, such as ^{tBuDip}nacnacH.²¹ These bulky substituents provide a significant steric effect and also have an electronic impact on the nature of the ligand system.

A common method for the synthesis of nacnac ligands is a simple condensation reaction between a β -diketone and two equivalents of anilines or primary amines,^{1,21–24} although for some examples, the activation of the diketone *via* a Meerwein salt has been employed.¹ For the construction of bulky backbone variants like ^{tBuDip}nacnacH,²¹ ^{tBuMes}nacnacH²¹ and ^{PhDip}nacnacH,²⁵ however, two ArN-containing halves have to be synthesised followed by linking these in a C–C bond formation reaction to build the backbone unit in a multistep approach. Aside from hydrocarbyl groups for backbone substituents (R), introduction of secondary amino groups in these positions affords bisamidines that act as electronically modified, electron-rich “*N*-nacnac” ligands.^{26,27} We have very recently introduced a one-pot condensation method to prepare ^{iPrDip}nacnacH²⁸ and this work expands on this backbone modification. We also consider ^{EtAr}nacnacH proligands which have been sporadically mentioned in the literature,^{29–32} but in some cases without much synthetic or spectroscopic detail.

2. Results and discussion

Ligand synthesis

Installing two ethyl groups into the backbone unit, *i.e.*, forming ^{EtAr}nacnacH proligands, can be achieved by simply modifying a common acid-catalysed condensation reaction^{22,32} according to Scheme 1. This requires stoichiometric *para*-toluenesulfonic acid hydrate (*p*TsOH·H₂O), an arene solvent such as toluene and heating under reflux with the use of a Dean–Stark trap to remove water from the condensation reaction system. For sterically more demanding Ar = Dip the use of xylene for up to two days has been advantageous. We recommend carrying out reactions that require high temperatures or long reaction times under nitrogen atmosphere to limit



Scheme 1 Synthesis of proligands 1.

side-reactions. A typical alkaline workup with aqueous sodium carbonate solution and dichloromethane, followed by treatment with methanol afforded crystalline crops of ^{EtDip}nacnacH **1a** (65%), ^{EtDep}nacnacH **1b** (48%), ^{EtMes}nacnacH **1c** (72%), and ^{EtXyl}nacnacH **1d** (75%) in moderate to good isolated yields and higher *in situ* yields.

Moving to isopropyl-substituted targets, *i.e.*, to ^{iPrAr}nacnacH, we first tested simple condensation protocols for the synthesis of the *N*-mesityl substituted system, *i.e.*, to afford ^{iPrMes}nacnacH **2c**. Reactions of two equivalents of 2,4,6-trimethylaniline (mesidine, MesNH₂), one equivalent of *p*TsOH·H₂O, in refluxing xylene (mixture of isomers, boiling range. *ca.* 136–140 °C) with a Dean–Stark trap under nitrogen atmosphere for 16 h afforded predominantly the monosubstituted β -ketoimine species *iPrC*(NHMes)CHC(=O)*iPr* **3c** as a colourless crystalline material after workup (Fig. 2). The crude product from this reaction did show NMR resonances of low integration for the desired ^{iPrMes}nacnacH **2c**, and thus a more forcing reaction setup with longer reaction times was employed using 4.5 equivalents of MesNH₂, 2 equivalents of *p*TsOH·H₂O, and reflux in xylene under nitrogen for 6–7 days. This was followed by an alkaline aqueous work-up and distilling off of excess MesNH₂ under vacuum to afford crude **2c** as a light-brown oil. The crude product contained small quantities of crystalline *iPrC*(NHMes)CHC(=O)*iPr* **3c** as a side product which could be separated off *via* column chromatography to afford **2c** in 47% isolated yield (Scheme 2). In one instance, compound **2c** crystallised after standing at room temperature for months to a few large colourless crystals, and was structurally characterised (Fig. 3).





Fig. 2 Molecular structure of *iPrC*(NHMe)CHC(=O)*iPr* **3c** (30% thermal ellipsoids). Hydrogen atoms except H1 are omitted for clarity. Selected bond lengths (Å) and angles (°): O5–C4 1.247(2), N1–C2 1.343(2), N1–C6 1.438(2), C2–C3 1.386(3), C3–C4 1.420(3); N1–C2–C3 120.49(17), C2–N1–C6 127.07(16), C2–C3–C4 123.50(17), O5–C4–C3 123.34(17).





Scheme 2 Synthesis of $i\text{Pr}^{\text{Mes}}\text{nacnacH}$ **2c** and **3c**.

It is evident that the method leading to **2c** would be difficult to translate to derivatives with sterically more demanding aryl groups and led us to investigate chemical dehydrating agents with the aim of a one-step condensation protocol. For this, we explored the related powerful acidic dehydrating agent polyphosphoric acid trimethylsilylester, PPSE.^{33,34} PPSE is prepared by adding phosphorus pentoxide to hexamethyldisiloxane in dichloromethane, followed by a reflux period and removal of all volatiles. The honey-like residue is then treated with the respective substituted aniline and diketone (2,6-dimethyl-3,5-heptanedione) and heated to 170 °C for approximately 16–24 hours. For workup, the reaction mixture is cooled to only *ca.* 90–95 °C to prevent it from solidifying, which impedes the neutralisation, and is slowly and carefully (caution!) treated with aqueous sodium hydroxide (NaOH) solution, that can be introduced *via* the reflux condenser, to quench the mixture and bring it to a basic pH value. After further cooling, extraction with dichloromethane yielded the crude products that can in most cases be precipitated using methanol. This afforded $i\text{Pr}^{\text{Dip}}\text{nacnacH}$ **2a** (86%), $i\text{Pr}^{\text{Dep}}\text{nacnacH}$ **2b** (53%), and $i\text{Pr}^{\text{Mes}}\text{nacnacH}$ **2c** (51%) in good (**2a**, **2b**) or moderate (**2c**) isolate yields (Scheme 3). The yield for **2a** has been slightly optimised from our initial report²⁸ using this method by gradually lowering the stoichiometric amount of PPSE for the synthesis, and similar optimisations could be envisaged for the preparations of **2b** and **2c**. For the synthesis of $i\text{Pr}^{\text{Mes}}\text{nacnacH}$ **2c**, another main product was produced alongside **2c**, which was characterised as the

β -diketiminato phosphorus(v) oxide ($i\text{Pr}^{\text{Mes}}\text{nacnac}$) PO_2 **4c** in approximately 29% yield (Scheme 3). Thus, **2c** and **4c** were often afforded in approximately similar quantities. The compounds can be easily separated by adding *n*-hexane to the crude product which precipitates **4c**, see Fig. 4 for the molecular structure, and yielded highly soluble **2c**. Workups from syntheses of $i\text{Pr}^{\text{Dep}}\text{nacnacH}$ **2b** also contained varying quantities of ($i\text{Pr}^{\text{Dep}}\text{nacnac}$) PO_2 **4b**, according to NMR spectroscopy that were difficult to separate by extracting into different solvents or by fractional crystallisation. No related species was found alongside the synthesis of $i\text{Pr}^{\text{Dip}}\text{nacnacH}$ **2a**. Compound **4c** is air stable and a rare β -diketiminato phosphorus compound that is *N,N'*-chelated, *vide infra*. We eventually found that stirring ($i\text{Pr}^{\text{Mes}}\text{nacnac}$) PO_2 **4c** in an aqueous sodium hydroxide solution and dichloromethane mixture, *i.e.*, similar to the workup conditions, deprotects **4c** to afford the proligand **2c**, which can thus be obtained in high yield overall. It appears that the β -diketiminato phosphorus(v) oxide species ($i\text{Pr}^{\text{Ar}}\text{nacnac}$) PO_2 **4** are most stable for the smaller aryl groups (Mes) whereas the bulkier ones (Dip, Dep) more readily hydrolyse under the workup conditions. Proligand **1a** could also be prepared using the PPSE method, but without optimising the conditions, we found no improvement in yield compared to the standard condensation reaction (Scheme 1).

To investigate how widely PPSE can be used for the one-step condensation synthesis of bulky β -diketimine proligands, we used the same method that successfully forms compound **2**, but using 2,2,6,6-tetramethyl-3,5-heptanedione instead of 2,6-dimethyl-3,5-heptanedione as the diketone with 2,6-diisopropylaniline to attempt the synthesis of $t\text{Bu}^{\text{Dip}}\text{nacnacH}$ ²¹ in a one-step procedure. This, however, afforded a mixture of products after workup, did not yield detected quantities of $t\text{Bu}^{\text{Dip}}\text{nacnacH}$, and the conditions were deemed unsuitable to convert this bulkier diketone. The method was, however, suitable to convert dibenzoylmethane (1,3-diphenylpropane-1,3-dione) with 2,6-diisopropylaniline, DipNH_2 , to the known proligand $\text{Ph}^{\text{Dip}}\text{nacnacH}$ **5**,²⁵ in 56% isolated yield (unoptimised, Scheme 4). To test if the method is suitable to easily install bulky tertiary alkyl groups into the *N*-positions of a β -diketimine proligand, a reaction between 2,6-dimethyl-3,5-heptanedione, two equivalents of 1-adamantylamine (1-aminoadamantane) and the respective quantities of PPSE under the usual conditions was conducted. After workup, however, a product mixture was obtained that contained large quantities of 1-aminoadamantane and this system was not further studied.





Fig. 3 Molecular structure of $i\text{Pr}^{\text{Mes}}\text{nacnacH}$, **2c** (30% thermal ellipsoids). Hydrogen atoms except H1 are omitted for clarity. Selected bond lengths (Å) and angles (°): N1–C2 1.354(2), N5–C4 1.299(2), C2–C3 1.373(3), C3–C4 1.438(3); C2–N1–C6 127.68(17), N1–C2–C3 119.33(18), C2–C3–C4 125.27(18), N5–C4–C3 119.34(17).





Scheme 3 Synthesis of $(iPr^{Ar})nacnacH$ **2** and $(iPr^{Ar})nacnac)PO_2$ **4**.





Fig. 4 Molecular structure of $(iPr^{Mes})nacnac)PO_2$, **4c** (30% thermal ellipsoids). Hydrogen atoms are omitted for clarity. Selected bond lengths (Å) and angles ($^\circ$): P1–O1 1.4712(16), P1–O2 1.4720(17), P1–N5 1.7518(17), P1–N1 1.7548(17), N1–C2 1.356(3), N5–C4 1.356(3), C2–C3 1.387(3), C3–C4 1.390(3); O1–P1–O2 121.67(10), N5–P1–N1 98.47(8), O1–P1–N1 108.54(9), O2–P1–N1 108.50(9), O1–P1–N5 109.55(9), O2–P1–N5 107.62(9).



Scheme 4 Synthesis of $PhDip)nacnacH$ **5**.

The β -diketimine proligands $EtDip)nacnacH$, **1a** (Fig. S61[†]), $EtDep)nacnacH$, **1b** (Fig. S62[†]), $EtXyl)nacnacH$, **1d** (Fig. S63[†]), $iPrDep)nacnacH$, **2b** (Fig. S64[†]), and $iPr^{Mes})nacnacH$, **2c** (Fig. 3), were structurally characterised and show the expected overall structures and geometrical features, see the ESI[†] for details. Across the structures, the level of localised *versus* delocalised bonding in the conjugated backbone unit that can be observed varies and is comparable to previously characterised examples, *e.g.* **2a**,²⁸ but some disorder or poorly ordered features are often present and thus these structures are not discussed in detail. The mesityl compounds $iPrC(NHMe)CHC(=O)iPr$ **3c**, $iPr^{Mes})nacnacH$ **2c**, and $(iPr^{Mes})nacnac)PO_2$ **4c** (Fig. 1–3) are shown here with representative bond lengths and angles. Some localisation of bonds of the ligand backbone unit can be inferred from the structure of **2c**, the well-defined tautomeric form of **3c** is observed, and the geometrical features are unremarkable. Similarly, the NMR spectroscopic data of compounds **1**, **2**, and **3c** are as expected. The three discussed $RDip)nacnacH$ proligands, for R = Me, Et, iPr, were exclusively observed by NMR spectroscopy in solution in their tautomeric


enamine form, *i.e.*, with a highly downfield shifted NH 1H NMR resonance near δ 12 ppm. The $tBuDip)nacnacH$ proligand, on the other hand, shows the dominant diimine form in solution with some percentage present in the enamine form (Fig. 1C).²¹ This may be explained by considering the basicity of the deprotonated fragments, $RDip)nacnac^-$. In these, the HOMO is typically predominantly associated with the p-orbital on the (delocalised) backbone CH unit^{1,18} (“enamine form”) which is likely destabilised by the positive inductive effect of the electron-rich *tert*-butyl groups on the neighbouring iminyl groups. Thus, this tautomer may become less stable, the CH backbone unit becomes the preferred position for protonation and the diimine form of the proligand becomes energetically favourable.

Compound **4c** shows the phosphorus centre to be N,N' -chelated as part of a six-membered ring with P=O bonds (P1–O1 1.4712(16), P1–O2 1.4720(17)) that are of similar lengths to those found in a few related species.^{35–37} The N–P–N angle ($98.47(8)^\circ$) is narrow due to the constraints of the heterocycle and, accordingly, the O–P–O angle ($121.67(10)^\circ$) is quite obtuse. The introduction of phosphorus centres into a classical β -diketiminate system^{38–45} is relatively rare and often met with unexpected product outcomes displaying different ligand–P connectivities, for example, when targeted *via* salt metathesis of β -diketiminate alkali metal complexes with phosphorus(III) halides. Often, the phosphorus centre forms a heterocycle with the β -diketiminate involving an activated backbone methyl group or simply bonds to the γ -carbon. N,N' -bound β -diketiminate phosphorus systems are rare^{42,43} and both control and prediction of the product type can be difficult. Thus, the convenient one-step formation of **4c** offers a new synthetic entry route.

Proligand deprotonation: conversion to alkali metal complexes

Alkali metal complexes of sterically demanding β -diketiminate ligands^{20,21,46–52} are a common synthon in β -diketiminate chemistry because they are generally easy to obtain and undergo wide salt metathesis chemistry to coordinate the ligand entity to a wide range of elements.^{1–16} *s*-block organometallics such as *n*-butyllithium and benzyl potassium are routinely employed to deprotonate β -diketimines to alkali metal complexes in coordinating or hydrocarbon solvents. These general methods can also be used for the $RAr)nacnacH$ compounds described herein (Scheme 5), and more details are provided on reactions of Dip-containing variants $EDip)nacnacH$ **1a** and $iPrDip)nacnacH$ **2a**. As widely performed, lithiations of **1a**





Scheme 5 Formation of β -diketiminato alkali metal complexes.

and **2a**, and the other presented proligands, with *n*-butyllithium can be generally carried out in hydrocarbon solvents (toluene, benzene, *n*-hexane), coordinating solvents (THF, diethylether) or solvent mixtures. Reactions of **1a** or **2a**, respectively, with benzyl potassium are typically carried out by stirring a suspension of a 1:1 mixture in toluene overnight. Often, alkali metal bis(trimethylsilyl)amide complexes, $[M\{N(SiMe_3)_2\}]$, M = Li, Na, K, have been used for this type of conversion as well. We have studied the reaction between $^{RDip}nacnacH$ and one equivalent of $[Na\{N(SiMe_3)_2\}]$ in deuterated benzene on a small scale followed by 1H NMR spectroscopy (Scheme 6 and Fig. S53, S55, S56[†]). $^{MeDip}nacnacH$ reacts slowly with $[Na\{N(SiMe_3)_2\}]$ at room temperature (e.g., ca. 40% conversion after 16 hours) or rapidly at 60 °C. $^{EtDip}nacnacH$ **1a** shows a slower reaction with $[Na\{N(SiMe_3)_2\}]$ at room temperature and only slowly converts to $[(^{EtDip}nacnac)Na]$ **7b** over several hours at 60 °C, e.g., ca. 60% conversion after 5 hours and full conversion overnight. For the same reaction with $^{iPrDip}nacnacH$ **2a**, however, we find only very sluggish conversion at 60 °C and this system required prolonged heating, for example, around five days to afford full conversion to $[(^{iPrDip}nacnac)Na]$ **7a**. We next investigated the reaction between $^{RDip}nacnacH$ and ten equivalents of $[K\{N(SiMe_3)_2\}]$ in deuterated benzene more quantitatively under pseudo-first order reaction conditions and found rate constants of $k'_{MeDip} = 3.0 \times 10^{-2} \text{ min}^{-1}$ at room temperature, $k'_{1a} = 7.6 \times 10^{-3} \text{ min}^{-1}$ at room temperature, and $k'_{2a} = 2.5 \times 10^{-3} \text{ min}^{-1}$ at 60 °C (Fig. S57–S60[†]). These differences highlight the significant effect of the ligand backbone, where $^{MeDip}nacnacH$ is deprotonated approximately four times faster than **1a**, and **1a** is deprotonated approximately three times faster at room temperature compared with **2a** at 60 °C. In stoichiometric reactions, **2a** can be more quickly deprotonated with the smaller and stronger amide base $[Li(NEt_2)]$ in deuterated benzene to $[(^{iPrDip}nacnac)Li]$ **6a** after heating to 60 °C for two days. Thus, the acidity of all the studied proligands $^{RDip}nacnacH$, R = Me, Et, *i*Pr, is higher than that of $HN(SiMe_3)_2$ (pK_a value 25.8).^{5,3} It is tempt-

ing to suggest that in addition to there being different kinetic barriers for the deprotonation of $^{RDip}nacnacH$ with $[Na\{N(SiMe_3)_2\}]$, likely for steric reasons, the lower acidity of **2a** in the series also leads to the markedly slower deprotonation reaction at 60 °C and a more basic and less thermodynamically favoured sodium complex (**7a**) when compared with reactions of $^{MeDip}nacnacH$ and **1a**. As part of this study, we generally found that $^{EtAr}nacnac$ -compounds show a higher solubility in aromatic or aliphatic solvents compared with $^{iPrAr}nacnac$ -compounds and this can play a role in their further transformations.

As part of the study, NMR data was acquired for a series of species, see Scheme 5, and the molecular structures of $[(^{iPrDip}nacnac)Li]$ **6a** (Fig. 5), $[(^{iPrDip}nacnac)Li(OEt_2)]$ **6a(OEt₂)** (Fig. S71[†]), and $[(^{EtDip}nacnac)K(C_6H_6)]$ **8b(C₆H₆)** (Fig. 6), were structurally characterised. Complex **6a** crystallised as a weakly-interacting dimer, highly similar to the structure of $[(^{MeDip}nacnac)Li]$. For the latter, a weakly-bound dodecameric isomer has been characterised as well.^{2,3} The molecular struc-





Fig. 5 Molecular structure of $[(^{iPrDip}nacnac)Li]$ **6a** (30% thermal ellipsoids). Hydrogen atoms are omitted for clarity, and weak Li...C interactions are shown with dashed bonds. Selected bond lengths (Å) and angles (°): N1–Li1 1.886(4), N5–Li1 1.893(4), Li1...C28' 3.199(5); N1–Li1–N5 100.2(2).



Scheme 6 Formation of β -diketiminato alkali metal complexes via alkali metal bis(trimethylsilyl)amides.






Fig. 6 Molecular structure of $[(^{\text{EtDip}}\text{nacnac})\text{K}(\text{C}_6\text{H}_6)]$ **8b** (C_6H_6) (30% thermal ellipsoids). Hydrogen atoms are omitted for clarity, and weak $\text{K}\cdots\text{C}$ interactions are shown with dashed bonds. Selected bond lengths (\AA) and angles ($^\circ$): K1-N1 2.6979(12), K1-N5 2.6613(12), $\text{K1}\cdots\text{H25}'$ 2.831, $\text{K1}\cdots\text{C25}'$ 3.2747(16); N5-K1-N1 70.67(3).

ture of **6a**(OEt_2) also mimics the previously characterised $[(^{\text{MeDip}}\text{nacnac})\text{Li}(\text{OEt}_2)]$.²³ In complex **8b**(C_6H_6), the K^+ ion is N,N' -chelated by the nacnac ligand and approximately η^4 -coordinated by a benzene molecule and shows a contact to a *para*-CH unit of a Dip-substituent from a neighbouring molecule leading to weak one-dimensional contacts in the lattice. In benzene-free $[(^{\text{MeDip}}\text{nacnac})\text{K}]$, the “completion” of the potassium coordination sphere is achieved by $\text{K}\eta^5$ -coordination to a neighbouring Dip-substituent.^{46,47}


Synthesis of β -diketiminato-derived silylene compounds

A case where the facile activation of the β -diketiminato ligand backbone plays a crucial role in product stability and compound reactivity is in low oxidation state silicon chemistry.^{54–59} Driess and co-workers have introduced a reactive two-coordinate silylene based on the dideprotonated β -diketimine $\text{MeDip}_2\text{nacnacH}$, *i.e.*, $(^{\text{MeDip}}\text{nacnac})\text{Si}$, where $\text{MeDip}_2\text{nacnac}'$ is $\text{H}_2\text{CC}(\text{NDip})\text{CHC}(\text{NDip})\text{Me}$.^{60,61} Related to this, the groups of Driess⁶² and Aldridge⁶³ have recently introduced and studied the chemistry of rare β -diketiminato chlorosilylene species and ligand backbone modifications were instrumental in accessing the desired target compounds.

Reacting $[(^{\text{iPrDip}}\text{nacnac})\text{Li}]$ **6a** with Roesky, Stalke and co-workers' N -heterocyclic carbene adduct IDipSiCl_2 ($\text{IDip} = \text{IPr} = \{\text{HC}(\text{NDip})\}_2\text{C}$)⁶⁴ as a SiCl_2 source in an attempted salt metathesis reaction in deuterated benzene afforded large quantities of proligand **2a** alongside uncoordinated IDip , as judged by ^1H NMR spectroscopy. In this reaction, we suspected that the backbone CH on the IDip ligand may be activated *via* an abnormal carbene-species^{65–67} with the highly basic ligand system using **6a**. Thus, we attempted a similar salt metathesis reaction of **6a** using the imidazolynilidene-based N -heterocyclic carbene adduct SIDipSiBr_2 ($\text{SIDip} = \text{SIPr} = \{\text{H}_2\text{C}(\text{NDip})\}_2\text{C}$)⁶⁸ introduced by Filippou and co-workers. In the latter, the saturated N -heterocyclic carbene backbone appears to be more inert towards deprotonation reactions in general. This reaction, however, also formed proligand **2a** and uncoordinated SIPr as part of the reaction mixture. Switching to a

similar reaction of the less basic $[(^{\text{EtDip}}\text{nacnac})\text{Li}]$ **6b** with SIDipSiBr_2 in deuterated benzene which was monitored by ^1H NMR spectroscopy saw the starting materials get consumed and showed the appearance of NMR resonances for a main new β -diketiminato-containing product of anticipated symmetry plus those for uncoordinated SIDip . Performing the reaction on a slightly larger scale at room temperature afforded $(^{\text{EtDip}}\text{nacnac})\text{SiBr}$ **9** in around 43% isolated yield (Scheme 7), which could also be structurally characterised, see Fig. 7. The overall molecular structure of **9** is similar to those of β -diketiminato chlorosilylenes introduced by Driess⁶² and Aldridge.⁶³ In **9**, the β -diketiminato-Si-Br angle is quite acute; $83.81(10)^\circ$ for the $\text{Br1-Si1}\cdots\text{C3}_{\text{backbone}}$ angle and $98.75(10)^\circ$ when using the $\text{Br1-Si1-(N1}\cdots\text{N5-midpoint)}$ angle. The Si-Br distance in **9** (2.405(2) \AA) is similar to that found in a diimino-phosphinate silicon(II) bromide (2.4545(10) \AA)⁶⁹ whereas the one present in a related amidinate silicon(II) bromide⁷⁰ is too disordered for reliable comparison. In solution, compound **9** shows four doublets and two septets for the protons of the isopropyl groups by ^1H NMR spectroscopy in accordance with its expected symmetry showing different environments above and below the nacnac-Si-plane. A $^{29}\text{Si}\{^1\text{H}\}$ NMR resonance of δ -7.4 ppm was found for compound **9** that lies in the expected range (*cf* δ -10.7 ppm (ref. 62), δ 2 ppm (ref. 63)).

To test the robustness of the ligand system, we treated $(^{\text{EtDip}}\text{nacnac})\text{SiBr}$ **9** with the strong amide base $[\text{KN}\{\text{N}(\text{SiMe}_3)_2\}]$ in aromatic solvents at room temperature and the reaction led



Scheme 7 Synthesis of $(^{\text{EtDip}}\text{nacnac})\text{SiBr}$ **9**.







Fig. 7 Molecular structure of $(^{\text{EtDip}}\text{nacnac})\text{SiBr}$ **9** (30% thermal ellipsoids). Hydrogen atoms and the minor component of disorder are omitted for clarity. Selected bond lengths (Å) and angles ($^{\circ}$): Br1–Si1 2.405(2), Si1–N1 1.8406(12), Si1–N5 1.8652(12), N1–C2 1.3505(18), C2–C3 1.383(2), C3–C4 1.399(2), N5–C4 1.3323(18); N1–Si1–N5 94.53(6), N1–Si1–Br1 97.20(9), N5–Si1–Br1 94.67(8), C2–C3–C4 125.50(14), Br1–Si1...C3 83.81(10), Br1–Si1–(N1...N5-midpoint) 98.75(10).

to formal HBr elimination and formation of $(^{\text{EtDip}}\text{nacnac})\text{Si}$ **10**, where $^{\text{EtDip}}\text{nacnac}'$ is the backbone-deprotonated “divalent” ligand variant $\text{MeCHC}(\text{NDip})\text{CHC}(\text{NDip})\text{Et}$, *i.e.*, an analogue of Driess’ silylene, plus KBr and $\text{HN}(\text{SiMe}_3)_2$ (Scheme 8). This demonstrates that the backbone positions in $^{\text{EtDip}}\text{nacnac}$ compounds can still be deprotonated, but likely less easily than those of $^{\text{MeDip}}\text{nacnac}$ derivatives. Compound **10** (Fig. 8) crystal-



Scheme 8 Synthesis of $(^{\text{EtDip}}\text{nacnac})\text{Si}$ **10**.




Fig. 8 Molecular structure of $(^{\text{EtDip}}\text{nacnac})\text{Si}$ **10** (30% thermal ellipsoids). Hydrogen atoms and the minor component disorder are omitted for clarity. Selected bond lengths (Å) and angles ($^{\circ}$): Si1–N1 1.7358(12), Si1–N1' 1.7357(12); N1–Si1–N1' 99.23(8).

lised with half a molecule in the asymmetric unit and thus the backbone unit is disordered by symmetry and individual bond lengths in that part of the molecule will not be commented on. The silicon(II) centre is part of a planar six-membered ring formed by bonding to both nitrogen atoms of the $^{\text{EtDip}}\text{nacnac}'$ ligand. The average Si–N distance in **10** (1.736 Å) is marginally longer than the average Si–N distance in Driess’ silylene ($^{\text{MeDip}}\text{nacnac}'\text{Si}$ (1.708 Å),⁶⁰ and as expected significantly shorter than those in the three coordinate silicon(II) compound **9** (1.853 Å mean). There is an absence of any weak long-range Si...Si contacts in the solid state structures of both **10** and **9**. NMR spectra of **10** are broadly comparable to those of related ($^{\text{MeDip}}\text{nacnac}'\text{Si}$) showing four doublets and two septets for the protons of the isopropyl groups (^1H NMR spectroscopy) as expected for a planar central unit with two different molecular halves plus resonances that support the backbone deprotonation, *i.e.*, a quartet (1 H) at δ 3.84 ppm and an associated doublet (3 H) at δ 1.62 ppm. The ^{29}Si NMR resonance of **10** (δ 87.9 ppm) is very close to that of ($^{\text{MeDip}}\text{nacnac}'\text{Si}$) (δ 88.4 ppm).⁶⁰

3. Conclusions

In this work we have presented the synthesis of a range of β -diketimine proligands, $^{\text{RAR}}\text{nacnacH}$, where the backbone substituents (R) are Et or *i*Pr and the *N*-aryl groups (Ar) are Dip, Dep, Mes or Xyl. Derivatives with R = Et can be synthesised by a common condensation type protocol, whereas for derivatives with R = *i*Pr, a powerful dehydrating agent, PPSE, was used in a one-pot procedure. The latter also afforded the phosphorus(V) β -diketiminate compound ($^{\text{iPrMes}}\text{nacnac})\text{PO}_2$ **4c** which could be hydrolysed to $^{\text{iPrMes}}\text{nacnacH}$ **2c** under aqueous alkaline conditions and likely presents the initial product formed during the assembly reaction in PPSE before workup. The β -diketimine proligands $^{\text{RAR}}\text{nacnacH}$ can be converted to alkali metal complexes using common reagents such as *n*-butyllithium and benzyl potassium. Reactions of $^{\text{RDip}}\text{nacnacH}$ with sodium or potassium bis(trimethylsilyl)amide highlight significant differences in reaction rates for the deprotonation reaction and show one aspect of an influence of the alkyl substitution in the backbone unit. Furthermore, the different basicities and/or minor differences in steric profile of the ligands $^{\text{RDip}}\text{nacnac}^-$ play a role in salt metathesis of their alkali metal complexes. For $[(^{\text{iPrDip}}\text{nacnac})\text{Li}]$ **6a** side-reactions such as proton abstraction were observed likely due to the high basicity, whereas less basic $[(^{\text{EtDip}}\text{nacnac})\text{Li}]$ **6b** could be used in a salt metathesis approach with SiDipSiBr_2 to afford the rare β -diketiminate silicon(II) bromide compound ($^{\text{EtDip}}\text{nacnac})\text{SiBr}$ **9**. Nevertheless, despite the protected backbone unit in **9**, HBr can be eliminated using $[\text{KN}\{\text{N}(\text{SiMe}_3)_2\}]$ to afford the analogue of Driess’ silylene, ($^{\text{EtDip}}\text{nacnac}'\text{Si}$) **10**. Now, a range of backbone R groups including Me, Et, *i*Pr, *t*Bu and Ph can be easily introduced to popular $^{\text{RAR}}\text{nacnacH}$ proligands and all except R = *t*Bu are accessible *via* a one-step protocol. This allows researchers to choose suitable nacnac- or BDI-ligands



from a wider pool based on steric profile, basicity, solubility, and ligand robustness for their choice of application.

Conflicts of interest

There are no conflicts of interest to declare.

Acknowledgements

We are grateful to the EPSRC DTG (EP/R513337/1), the China Scholarship Council, and the School of Chemistry at the University of St Andrews for support.

References

- L. Bourget-Merle, M. F. Lappert and J. R. Severn, *Chem. Rev.*, 2002, **102**, 3031–3065.
- Y. C. Tsai, *Coord. Chem. Rev.*, 2012, **256**, 722–758.
- E. Glöckler, S. Ghosh and S. Schulz, *Polym. Rev.*, 2023, **63**, 478–514.
- S. Ito and Y. Morimoto, β -Diketiminates as Redox Non-innocent Supporting Ligands in Coordination Chemistry, in *Chemical Science of π -Electron Systems*, ed. T. Akasaka, *et al.*, Springer, Japan 2016.
- C. Camp and J. Arnold, *Dalton Trans.*, 2016, **45**, 14462–14498.
- R. Kretschmer, *Chem. – Eur. J.*, 2020, **26**, 2099–2119.
- R. B. Ferreira and L. J. Murray, *Acc. Chem. Res.*, 2019, **52**, 447–455.
- S. P. Sarish, S. Nembenna and H. W. Roesky, *Acc. Chem. Res.*, 2011, **44**, 157–170.
- C. Jones, *Nat. Rev. Chem.*, 2017, **1**, 0059.
- B. Rösch and S. Harder, *Chem. Commun.*, 2021, **57**, 9354–9365.
- L. S. Vasil'ev, S. V. Baranin and I. V. Zavarzin, *Russ. Chem. Bull., Int. Ed.*, 2017, **66**, 1398–1418.
- M. Zhong, S. Sinhababu and H. W. Roesky, *Dalton Trans.*, 2020, **49**, 1351–1364.
- C. Chen, S. M. Bellows and P. L. Holland, *Dalton Trans.*, 2015, **44**, 16654–16670.
- R. L. Webster, *Dalton Trans.*, 2017, **46**, 4483–4498.
- S. Hohloch, B. M. Kriegel, R. G. Bergman and J. Arnold, *Dalton Trans.*, 2016, **45**, 15725–15745.
- D. Zhu and P. H. M. Budzelaar, *Dalton Trans.*, 2013, **42**, 11343–11354.
- K. Huse, C. Wölper and S. Schulz, *Eur. J. Inorg. Chem.*, 2018, 3472–3480.
- K. Huse, H. Weinert, C. Wölper and S. Schulz, *Dalton Trans.*, 2020, **49**, 9773–9780.
- R. Lalrempuia, C. E. Kefalidis, S. J. Bonyhady, B. Schwarze, L. Maron, A. Stasch and C. Jones, *J. Am. Chem. Soc.*, 2015, **137**, 8944–8947.
- H. Videa and A. J. Martínez-Martínez, *Dalton Trans.*, 2023, **52**, 13058–13062.
- P. H. M. Budzelaar, A. B. van Oort and A. G. Orpen, *Eur. J. Inorg. Chem.*, 1998, 1485–1494.
- D. J. Mindiola, P. L. Holland and T. H. Warren, Complexes of Bulky β -Diketiminates Ligands, in *Inorganic Syntheses*, ed. T. B. Rauchfuss, John Wiley & Sons, 2010, vol. 35, pp. 1–55.
- M. Stender, R. J. Wright, B. E. Eichler, J. Prust, M. M. Olmstead, H. W. Roesky and P. P. Power, *J. Chem. Soc., Dalton Trans.*, 2001, 3465–3469.
- I. El-Zoghbi, A. Ased, P. O. Oguadinma, E. Tchirioua and F. Schaper, *Can. J. Chem.*, 2010, **88**, 1040–1045.
- M. Arrowsmith, M. R. Crimmin, M. S. Hill and G. Kociok-Köhn, *Dalton Trans.*, 2013, **42**, 9720–9726.
- D. C. H. Do, A. Keyser, A. V. Protchenko, B. Maitland, I. Pernik, H. Niu, E. L. Kolychev, A. Rit, D. Vidovic, A. Stasch, C. Jones and S. Aldridge, *Chem. – Eur. J.*, 2017, **23**, 5830–5841.
- M. A. Land, B. Huo, K. N. Robertson, K. E. O. Ylijoki, P. T. K. Lee, J. Areephong, D. Vidovic and J. A. C. Clyburne, *Dalton Trans.*, 2018, **47**, 10195–10205.
- S. Burnett, C. Bourne, A. M. Z. Slawin, T. van Mourik and A. Stasch, *Angew. Chem., Int. Ed.*, 2022, **61**, e202204472.
- M. Gao, J. Wang, H. Liu, J. Ma, X. Cai, C. Li, J. Hu, X. Zhang, X. Li, Z. Zhang, R. Duan, L. Yang and C. Ma, *US patent*, US2017/0073441A1, 2017.
- W. Liu, Y. Ding, D. Jin, Q. Shen, B. Yan, X. Ma and Z. Yang, *Green Chem.*, 2019, **21**, 3812–3815.
- M. C. Eaton, B. J. Knight, R. Brahmī, R. B. Ferreira, V. J. Catalano, A. L. Rheingold, I. Ghiviriga and L. J. Murray, *J. Org. Chem.*, 2020, **85**, 13579–13588.
- N. Demidov, M. Grebogi, C. Bourne, A. P. McKay, D. B. Cordes and A. Stasch, *Molecules*, 2023, **28**, 7569.
- S. E. López, J. Restrepo and J. Salazar, *J. Chem. Res.*, 2007, 497–502.
- K. Yamamoto and H. Watanabe, *Chem. Lett.*, 1982, 1225–1228.
- A. E. Sheshenev, E. V. Boltukhina, A. A. Grishina, I. Cisarova, I. M. Lyapkalo and K. K. (Mimi) Hii, *Chem. – Eur. J.*, 2013, **19**, 8136–8143.
- H. Kalita, W.-Z. Lee and M. Ravikanth, *Inorg. Chem.*, 2014, **53**, 9431–9438.
- A. Hinz, R. Kuzora, U. Rosenthal, A. Schulz and A. Villinger, *Chem. – Eur. J.*, 2014, **20**, 14659–14673.
- P. B. Hitchcock, M. F. Lappert and J. E. Nycz, *Chem. Commun.*, 2003, 1142–1143.
- P. J. Ragogna, N. Burford, M. D'eon and R. McDonald, *Chem. Commun.*, 2003, 1052–1053.
- R. Rahm, S. Espenlaub, U. R. Werz and G. Maas, *Heteroat. Chem.*, 2005, **16**, 437–446.
- D. Vidovic, Z. Lu, G. Reeske, J. A. Moore and A. H. Cowley, *Chem. Commun.*, 2006, 3501–3503.
- Z. Lu, G. Reeske, J. A. Moore and A. H. Cowley, *Chem. Commun.*, 2006, 5060–5061.
- L. A. Lesikar, W. D. Woodul and A. F. Richards, *Polyhedron*, 2007, **26**, 3242–3246.



- 44 Z. Lu, M. Findlater and A. H. Cowley, *Chem. Commun.*, 2008, 184–186.
- 45 N. D. Reddy, A. Jana, H. W. Roesky, P. P. Samuel and C. Schulzke, *Dalton Trans.*, 2010, **39**, 234–238.
- 46 W. Clegg, E. K. Cope, A. J. Edwards and F. S. Mair, *Inorg. Chem.*, 1998, **37**, 2317–2319.
- 47 A. G. Avent, M. R. Crimmin, M. S. Hill and P. B. Hitchcock, *J. Organomet. Chem.*, 2006, **691**, 1242–1250.
- 48 H. Hamaki, N. Takeda, T. Yamasaki, T. Sasamori and N. Tokitoh, *J. Organomet. Chem.*, 2007, **692**, 44–54.
- 49 A. J. Wooles, W. Lewis, A. J. Blake and S. T. Liddle, *Organometallics*, 2013, **32**, 5058–5070.
- 50 R. M. Gault, J. R. Lynch, A. R. Kennedy, J. Barker, J. Reid and R. E. Mulvey, *Inorg. Chem.*, 2021, **60**, 6057–6064.
- 51 B. Rösch, T. X. Gentner, J. Eyselein, J. Langer, H. Elsen and S. Harder, *Nature*, 2021, **592**, 717–721.
- 52 D. D. L. Jones, S. Watts and C. Jones, *Inorganics*, 2021, **9**, 72.
- 53 R. R. Fraser, T. S. Mansour and S. Savard, *J. Org. Chem.*, 1985, **50**, 3232–3234.
- 54 S. Yao, A. Saddington, Y. Xiong and M. Driess, *Acc. Chem. Res.*, 2023, **56**, 475–488.
- 55 Y. Wang and G. H. Robinson, *J. Am. Chem. Soc.*, 2023, **145**, 5592–5612.
- 56 C. Shan, S. Yao and M. Driess, *Chem. Soc. Rev.*, 2020, **49**, 6733–6754.
- 57 S. Fujimori and S. Inoue, *Eur. J. Inorg. Chem.*, 2020, 3131–3142.
- 58 R. S. Ghadwal, R. Azhakar and H. W. Roesky, *Acc. Chem. Res.*, 2013, **46**, 444–456.
- 59 M. Asay, C. Jones and M. Driess, *Chem. Rev.*, 2011, **111**, 354–396.
- 60 M. Driess, S. Yao, M. Brym, C. van Wüllen and D. Lentz, *J. Am. Chem. Soc.*, 2006, **128**, 9628–9629.
- 61 M. Driess, S. Yao, M. Brym and C. van Wüllen, *Angew. Chem., Int. Ed.*, 2006, **45**, 6730–6733.
- 62 Y. Xiong, S. Yao, A. Kostenko and M. Driess, *Dalton Trans.*, 2018, **47**, 2152–2155.
- 63 D. C. H. Do, A. V. Protchenko, M. A. Fuentes, J. Hicks, E. L. Kolychev, P. Vasko and S. Aldridge, *Angew. Chem., Int. Ed.*, 2018, **57**, 13907–13911.
- 64 R. S. Ghadwal, H. W. Roesky, S. Merkel, J. Henn and D. Stalke, *Angew. Chem., Int. Ed.*, 2009, **48**, 5683–5686.
- 65 S. C. Sau, P. K. Hota, S. K. Mandal, M. Soleilhavoup and G. Bertrand, *Chem. Soc. Rev.*, 2020, **49**, 1233–1252.
- 66 V. Nesterov, D. Reiter, P. Bag, P. Frisch, R. Holzner, A. Porzelt and S. Inoue, *Chem. Rev.*, 2018, **118**, 9678–9842.
- 67 S. Würtemberger-Pietsch, U. Radius and T. B. Marder, *Dalton Trans.*, 2016, **45**, 5880–5895.
- 68 A. C. Filippou, O. Chernov and G. Schnakenburg, *Chem. – Eur. J.*, 2011, **17**, 13574–13583.
- 69 S. Takahashi, J. Sekiguchi, K. Nakaya, A. Ishii and N. Nakata, *Inorg. Chem.*, 2022, **61**, 7266–7273.
- 70 H.-X. Yeong, K.-C. Lau, H.-W. Xi, K. H. Lim and C.-W. So, *Inorg. Chem.*, 2010, **49**, 371–373.

