

REVIEW TOPIC OF THE WEEK

The Myth of the “Vulnerable Plaque”

Transitioning From a Focus on Individual Lesions to Atherosclerotic Disease Burden for Coronary Artery Disease Risk Assessment



Armin Arbab-Zadeh, MD, PhD,* Valentin Fuster, MD, PhD†

ABSTRACT

The cardiovascular science community has pursued the quest to identify vulnerable atherosclerotic plaque in patients for decades, hoping to prevent acute coronary events. However, despite major advancements in imaging technology that allow visualization of rupture-prone plaques, clinical studies have not demonstrated improved risk prediction compared with traditional approaches. Considering the complex relationship between plaque rupture and acute coronary event risk suggested by pathology studies and confirmed by clinical investigations, these results are not surprising. This review summarizes the evidence supporting a multifaceted hypothesis of the natural history of atherosclerotic plaque rupture. Managing patients at risk of acute coronary events mandates a greater focus on the atherosclerotic disease burden rather than on features of individual plaques. (J Am Coll Cardiol 2015;65:846–55) © 2015 by the American College of Cardiology Foundation.

Cardiovascular atherosclerotic disease is the leading cause of death in Western industrialized nations and in developing countries (1). Strategies to prevent acute coronary events and their sequelae are among our most important public health priorities (2). Identifying patients at increased risk of acute coronary events who may benefit from intensified preventative measures is a major, ongoing challenge (2). Numerous factors (e.g., dyslipidemia, diabetes, and others), are associated with increased rates of adverse events; however, their hazard rates are too small for accurate individual risk assessments (3). The DIAD (Detection of Ischemia in Asymptomatic Diabetics) trial revealed that after 4.8 years of follow-up, 97% of asymptomatic patients with diabetes mellitus remained free of myocardial infarction and cardiac death (4). Even when combined as comprehensive risk scores (e.g., by the Framingham study [5]), predictive accuracy

is insufficient for adequate individual risk assessments, leading to substantial overtreatment and undertreatment, with associated morbidity and societal costs (6).

The mechanisms leading to adverse events from atherosclerotic disease are clearly more complex than initially assumed, explaining our difficulties in accurately predicting events in individuals (7,8). In addition to the presence, extent, and metabolic activity of atherosclerotic disease, individual adaptations and responses to thrombogenic stimulation from altered vascular function are critical for determining the risk of acute coronary events (7,9). Despite a consensus on the complexity of acute coronary event risk evaluation and the necessity for comprehensive patient assessment (10,11), recent efforts to identify high-risk patients have focused on using advanced imaging methods to detect single “vulnerable” atherosclerotic plaques (12). This narrow focus neglects the

From the *Department of Medicine, Cardiology Division, Johns Hopkins University, Baltimore, Maryland; and the †Mount Sinai Medical Center, Icahn School of Medicine, New York, New York. Dr. Zadeh is supported by grant K23-HL098368 from the National Institutes of Health. Dr. Fuster has reported that he has no relationships relevant to the contents of this paper to disclose. Peter Libby, MD, served as Guest Editor for this paper.

Listen to this manuscript's audio summary by **JACC** Editor-in-Chief Dr. Valentin Fuster.

You can also listen to this issue's audio summary by **JACC** Editor-in-Chief Dr. Valentin Fuster.



complexity of the processes leading to risk and lacks supporting evidence. In this review, we summarize the shift from conceptualizing acute coronary event risk as a simple cause-and-effect principle centered on high-risk plaques to a complex model involving numerous factors.

THE "VULNERABLE PLAQUE" CONCEPT

Pathology studies have demonstrated the common association of acute myocardial infarction with the rupture or erosion of a coronary atherosclerotic plaque (13,14), most frequently a thin-cap fibroatheroma (TCFA), characterized by a large lipid or necrotic core separated from the coronary arterial lumen by a thin membrane cap (15). Thus, the identification of TCFA in humans was assumed to signify a high risk of ensuing acute coronary events, which then might necessitate directed treatment or specific preventative measures (16). Accordingly, enormous efforts have been undertaken to enable the identification of TCFA and other high-risk plaque features in humans. The search term *vulnerable plaque* finds more than 400 current National Institutes of Health research awards totaling more than \$150 million per year (17) and almost 2,000 research papers in the National Library of Medicine database. Although not all of these efforts aim to identify "vulnerable plaques," this topic is clearly central to many investigations involving large amounts of research dollars. Industry has also been keenly interested in developing technologies for the visualization of "vulnerable plaques," with progression of several catheter-based inventions, notably intravascular ultrasound (IVUS)-virtual histology, thermography, infrared spectroscopy, palpography, and optical coherence tomography (OCT) to preclinical or clinical stages (12,18).

LIMITATIONS OF STUDIES SUPPORTING THE HIGH-RISK ATHEROSCLEROTIC PLAQUE CONCEPT

A number of clinical investigations used various imaging tools to identify high-risk atherosclerotic plaque features in order to predict an increased risk of adverse events. In one large, prospective clinical study, PROSPECT (Providing Regional Observations to Study Predictors of Events in the Coronary Tree), rates of adverse cardiac events according to types of coronary atherosclerotic plaque were investigated in more than 600 high-risk patients studied with IVUS-virtual histology (19). Although 596 TCFA were identified, only repeat hospitalization for chest pain was associated with events. This was expected, given


the typically larger luminal encroachment of TCFA compared with pathological intimal thickening (the prevalent type of lesion in the study). However, the risk of myocardial infarction or sudden cardiac death related to these lesions was very low (Figure 1). A similar study using IVUS-virtual histology (VIVA [VH-IVUS in Vulnerable Atherosclerosis]) reported nearly identical findings (20). Studies using OCT revealed greatly detailed plaque characteristics in patients with acute coronary syndromes and other at-risk populations (21). Similar to the information provided by IVUS, studies using OCT suggest that a larger lesion plaque burden might indicate an increased risk of acute coronary events (22). Noninvasive imaging studies of the coronary arteries using computed tomographic angiography reported increased rates of acute coronary syndromes in patients with low-attenuation plaques (presumably high in lipid content) with external remodeling compared with those without such plaques (23–25). Puchner et al. (26) recently reported independent prediction of acute

ABBREVIATIONS AND ACRONYMS

IVUS = intravascular ultrasound

TCFA = thin-cap fibroatheroma

FIGURE 1 Risk of MI or Death Associated With Individual Plaques in the PROSPECT Study



Maximum annualized risks (percent) of myocardial infarction (MI) or cardiovascular (CV) death associated with individual coronary atherosclerotic plaques identified at baseline by intravascular ultrasound-virtual histology in the PROSPECT (Providing Regional Observations to Study Predictors of Events in the Coronary Tree) study are shown (18). The rates are based on the occurrence of 6 events (6 myocardial infarctions and no deaths) after 3.4 years of follow-up among 1,005 coronary arterial sites with pathological intimal thickening and 595 thin-cap atheromas (TCFAs), assuming all events were caused by the respective plaque type, thus representing the worst-case scenario. Considering equal risk among the 3,160 plaques detected at baseline (best-case scenario), the event rate associated with each plaque would be only 0.06% per year. The risk of MI or death associated with individual TCFAs or vulnerable plaques is much smaller than what is conventionally considered high risk, even when maximal risk is assumed.

TABLE 1 Prevalence of Subclinical Coronary Atherosclerotic Plaque Ruptures (Percent) in Patients With Stable Coronary Heart Disease or Healthy Controls and in Patients With Acute Coronary Syndromes

First Author (Ref. #)	Mode of Assessment	Number of Stable CHD Patients or Controls*	Nonculprit Plaque Ruptures (%)	Number of ACS Patients	Nonculprit Plaque Ruptures (%)
Arbustini et al. (30)	Pathology	77*	17	106	58
Cheruvu et al. (38)	Pathology	13*	31	33	45
Rioufol et al. (41)	IVUS	—	—	24	79
Kotani et al. (37)	IVUS	48	6	38	11
Schoenhagen et al. (36)	IVUS	92	4	105	19
Hong et al. (35)	IVUS	113	5	122	17
Tanaka et al. (29)	IVUS	—	—	45	24
Takano et al. (34)	Angioscopy	—	—	327	9
Tanaka et al. (33)	OCT	—	—	45	24
Tian et al. (22)	OCT	—	—	82	10
Shimamura et al. (32)	OCT	191	17	—	—
Total		444		927	
Median			11.5		21.5

Multiple asymptomatic plaque ruptures are more frequently found in patients with ACS than in those with stable CHD or healthy controls, indicating the systemic inflammatory state in the coronary arteries with acute events.

*Healthy controls.

ACS = acute coronary syndrome; CHD = coronary heart disease; IVUS = intravascular ultrasound; OCT = optical coherence tomography.

coronary syndrome using computed tomography to identify similar high-risk characteristics in patients presenting with acute chest pain. Finally, plaque hemorrhage, assessed by magnetic resonance imaging, has been implicated in poor outcomes in patients with cerebrovascular disease (27,28). However, all of these studies claiming independent risk prediction of certain plaque features share the fundamental limitation that the atherosclerotic disease burden was not considered as a potential confounder. These "high-risk" features are conceivably mere markers of more extensive and/or active atherosclerotic disease compared with the control group. Given the overwhelming evidence for disease burden as a powerful predictor of outcome, any additional risk features should be assessed against it before we assume independent risk prediction. Therefore, despite promising results from a number of clinical studies, there is no conclusive evidence for truly independent risk prediction associated with high-risk plaque features. High-risk features may still be valuable as markers for disease burden or activity; however, such value has not been established.

EVIDENCE DIMINISHING THE SIGNIFICANCE OF IDENTIFYING "HIGH-RISK" PLAQUES

Despite our ability to identify atherosclerotic lesions that exhibit vulnerable characteristics using various imaging tools, clinical studies have failed to demonstrate meaningful clinical utility for plaque imaging.

These negative results are explained by the numerous pathological and clinical investigations demonstrating that many (if not most) plaques rupture without clinical syndromes (22,29–38). The percent of patients with subclinical plaque ruptures varies with their risk profile and the sensitivity of assessment methods, ranging from 4% to 79% (Table 1). Plaque rupture and its healing are frequently clinically silent but lead to progressive lumen obstruction (39,40). In lesions with advanced lumen narrowing (>50% stenosis), histopathology almost invariably reveals 1 or more healed subclinical plaque ruptures (39–41). Because healed plaque ruptures can be detected by pathology or imaging only within a certain time frame after the initial plaque disruption, the true rate of asymptomatic plaque ruptures is probably underestimated (40). In most patients with advanced atheroma, plaque rupture and subsequent healing have already occurred (40). Pathology studies show that approximately 10% of the subclinical United States adult population exhibits advanced coronary atherosclerosis (42), so it is reasonable to assume that many millions of persons unknowingly experience plaque ruptures each year.


Several longitudinal imaging studies in humans have demonstrated that plaque morphology changes over a few months, gaining or losing "vulnerable" characteristics (43–45). Using IVUS, Kubo et al. (43) found that 75% of TCFAs transition to thick-cap fibroatheromas or fibrotic plaques within a 12-month interval, presumably secondary to rupture and healing (Figure 2). None of these patients experienced events during this period, providing further evidence of frequent subclinical plaque alterations. A recent study used OCT to confirm the dynamic nature of coronary atherosclerotic disease, demonstrating that TCFAs in various stages of development are highly prevalent in patients with coronary atherosclerotic disease (46). In patients with acute coronary syndromes, plaque ruptures are frequently found apart from the culprit lesions, indicating that vulnerability is disseminated throughout the coronary tree (41). This suggests that detection of a state of vulnerability in a patient (e.g., widespread inflammation) is more important than detection of individual sites of vulnerability. The emerging picture of acute coronary event pathophysiology suggests a widespread, systemic condition with great unpredictability as to which particular lesion will be associated with clinically significant vascular thrombosis (47). Although plaque ruptures and erosions are indeed responsible for most culprit lesions in patients with acute events, because the frequency of subclinical plaque ruptures is vastly underestimated, the assumption

that identifying lesions prone to rupture will prevent acute coronary events was unrealistic. Of the many plaque ruptures occurring in patients with atherosclerotic disease, very few will trigger symptomatic events, rendering it exceedingly difficult to predict adverse outcomes associated with particular lesions. Identifying a single TCFA or other "high-risk plaque," without considering other clinical or imaging characteristics, is unlikely to be of incremental benefit for risk prediction over established factors (e.g., the extent and distribution of atherosclerotic plaque burden), because of the low risk associated with a given individual plaque and the temporal relationship of its vulnerable characteristics.

CURRENT PARADIGM OF ACUTE CORONARY EVENT PATHOPHYSIOLOGY

Over the past few decades, clinical and laboratory investigations have led to a more complex concept of the pathophysiology of acute coronary events, involving numerous processes, many with poorly understood interactions (7,8). Although the occurrence of acute coronary events typically requires alterations of coronary atherosclerotic plaques (rupture or erosion), a thrombosis-promoting milieu is necessary to allow a clinically significant decrease in coronary blood flow and associated myocardial ischemia (7,9). Such a setting appears to result from an unfortunate constellation of prothrombotic features, for example, in patients with increased inflammatory activity and systemic or local suppression of fibrinolytic performance, an extraordinarily large stimulus for thrombosis, vasoconstriction, and/or others (7). The respective contributions of these factors (some hereditary, some environmental) and their temporal relationships necessary to trigger clinically meaningful vascular thrombosis are unknown. Factors favoring thrombosis need to be collectively sufficient to tilt the scale away from localized thrombus and toward extensive vascular thrombosis (**Central Illustration**). Because numerous factors influence the performance of the coagulation system at any given point in time, acute coronary events may arise as result of a "perfect storm" scenario, in which plaque disruption occurs in a specific, thrombosis-promoting setting (7). The risk of an acute coronary event equals the probability of plaque rupture or erosion coinciding with vascular thrombosis-promoting conditions that cannot contain the thrombus in the vascular wall. Frequent plaque ruptures, as with a large, metabolically active atherosclerotic disease burden, increase the chance that a plaque rupture coincides with

FIGURE 2 Changes in TCFA During 12 Months of Follow-Up





Changes in thin-cap fibroatheromas (TCFA) observed with intravascular ultrasound-virtual histology 12 months after baseline imaging. Of 20 TCFA, only 5 remained unchanged, while 15 (75%) lost vulnerable characteristics and revealed thickening of the fibrous cap or transformed into fibrotic plaques. The data (Kubo et al. [43]) demonstrate high metabolic activity in atherosclerotic lesions and the short-lived nature of vulnerable plaque characteristics. None of the 99 patients experienced any events during follow-up; thus, many of the observed changes were likely the result of subclinical plaque ruptures.

a thrombosis-conducive setting. Accordingly, the strongest predictors of adverse events are the magnitude and activity of the coronary atherosclerotic plaque burden and the number of risk factors for a prothrombotic milieu, a concept supported by many clinical studies and epidemiologic data (3,5,48–50).

LESSON FOCUSED VERSUS DISEASE BURDEN FOCUSED RISK ASSESSMENT AND MANAGEMENT

Numerous clinical studies using conventional invasive coronary angiography, IVUS, and cardiac computed tomography (8,48,51) have confirmed the strong relationship between atherosclerotic disease burden and risk for adverse events. Despite capturing only calcified atherosclerotic disease, when compared directly, coronary calcium scoring was equivalent to traditional stenosis assessment for predicting mortality and myocardial infarction in asymptomatic patients (52). Using a comprehensive imaging approach in several vascular beds, the BioImage study recently revealed a high prevalence of atherosclerotic disease in patients categorized as high risk for cardiovascular events on the basis of clinical predictors (53). Halting coronary atherosclerotic disease progression and/or altering the vascular thrombosis-promoting milieu via platelet inhibition and risk factor interdiction are approaches proved to lower myocardial infarction and death rates (54–56). Conversely, meta-analyses have not demonstrated reduced rates of myocardial

CENTRAL ILLUSTRATION Fate of Ruptured Coronary Atherosclerotic Plaques According to Thrombotic Milieus**A****B**


Arbab-Zadeh, A. et al. J Am Coll Cardiol. 2015; 65(8):846–55.

The hypothesized interplay of prothrombotic and thrombosis resisting and containing factors that presumably determine the outcome of a ruptured coronary atherosclerotic plaque is shown. **(A)** In the most common scenario, small thrombus formation associated with plaque rupture is contained and vascular occlusive thrombus is inhibited. **(B)** In the less common scenario of several prothrombotic factors coinciding (e.g., inflammatory state, large lesion plaque burden, vasoconstriction, circadian rheological changes), local thrombosis associated with plaque rupture cannot be contained, and clinically significant vascular thrombosis occurs, triggering an acute coronary syndrome (ACS). The constellation of factors leading to these different outcomes is unknown.

infarction or death with lesion-based treatment (i.e., percutaneous coronary intervention) compared with medical therapy in patients with stable coronary artery disease (57,58). Contradicting some earlier reports (59–62), no benefit was shown, even when selecting patients with hemodynamically significant stenoses. These results confirm that risk is most strongly conveyed by the extent of coronary artery disease, but not necessarily by individual lesions, even when highly obstructive. This supports the controversial idea that severe coronary artery stenoses confer no greater risk of triggering acute coronary events than mild lesions (63). Earlier angiographic studies suggested that most myocardial infarctions arise from mild coronary artery stenoses (64,65), but recent data question this paradigm (66,67). Thrombus material accumulates over several days after a plaque disruption, which may not allow lesion size and a partly organized thrombus to be accurately distinguished (66). Pathology studies in patients who died suddenly found culprit lesions to have an average diameter stenosis of approximately 50%, with no clear relationship between stenosis severity and risk of death (68,69). Conversely, acute coronary death rarely results from lesions with <30% luminal stenosis (66,69). Thus, a certain local plaque volume appears necessary to trigger vascular thrombosis. However, given the lack of benefit with coronary stenting, as well as the large number of obstructive lesions found by imaging and autopsy in patients without symptoms of acute coronary events, stenosis severity is unlikely to substantially alter such risk beyond a particular threshold. Patients with high-grade coronary artery stenoses may conceivably carry an increased risk of myocardial infarction and death because these lesions are markers for advanced atherosclerotic disease in the coronary tree (7). Consistent with this notion, nonobstructive and obstructive coronary artery disease are associated with similar risks of myocardial infarction and death if the former affects a larger number of coronary arterial segments (Figure 3) (70). Overall, strong evidence supports addressing the extent and activity of the atherosclerotic burden and thrombosis-promoting risk factors for improved patient outcomes, but there is no conclusive evidence of incremental risk reduction with lesion-specific treatment.

AREAS OF UNCERTAINTY: INFLUENCE OF STUDY POPULATION

The atherosclerotic disease burden is a powerful predictor of outcomes for patients without history of coronary artery disease, facilitated by the ease of



coronary calcium scanning, which approximates the total coronary atherosclerotic disease volume (71). However, it is infrequently performed in patients with established coronary artery disease; thus, in this population, plaque burden data are more limited. Risk characteristics and the need for assessment may conceivably differ in patients with established coronary artery disease compared with those with history of acute coronary syndromes. Aside from calcium scanning, plaque burden assessment is technically difficult, and most available data were derived from IVUS imaging. Atherosclerotic plaque burden assessment using computed tomographic angiography has recently become feasible, but long-term outcome data are not yet available (72). Aggregate data from IVUS-derived plaque burden assessment reveal strong predictive power for outcomes in patients with established coronary artery disease (73). Conventional angiographic data for atherosclerotic disease burden are similarly predictive of patient outcomes and superior to myocardial ischemia testing in an analysis of the COURAGE (Clinical Outcomes Utilizing Revascularization and Aggressive Drug Evaluation) study (74). Thus, the risk associated with plaque burden applies to both asymptomatic patients without prior cardiac events and those with established, stable

coronary artery disease. Whether lesion-based treatment (aside from the culprit lesion) is beneficial in patients with acute coronary syndrome is unclear. Two recent small clinical studies suggested reduced rates of myocardial infarction and death after percutaneous coronary intervention of nonculprit lesions in patients with ST-segment elevation myocardial infarctions (75,76), in stark contrast to large, aggregate data suggesting the opposite (77). In addition to concerns about the effect of chance, given the small numbers of events in these 2 studies, the contributions of event allocation (e.g., differentiating spontaneous and periprocedural events) and varying levels of expertise at the study centers to these results remain unclear. PROSPECT, conducted in patients with acute coronary syndromes, found low risk of myocardial infarction and death associated with nonculprit lesions within 3 years of follow-up (19). A large clinical study, COMPLETE (Complete vs Culprit-Only Revascularization to Treat Multi-Vessel Disease After Primary PCI for STEMI), is underway to conclusively address the question of benefit of stenting nonculprit lesions in patients with ST-segment elevation myocardial infarction (78).

IS THERE A ROLE FOR INDIVIDUAL PLAQUE IMAGING?

Coronary artery imaging has provided insights into numerous lesion characteristics, but we have yet to identify which are useful for guiding management. Changing patterns of lesion characteristics, resulting from widespread use of lipid-modifying medications, pose an additional challenge (79). There are promising data for characterizing coronary arterial lesions before percutaneous interventions (80,81). Heavily calcified lesions adversely affect the outcome of coronary artery revascularization, and imaging data may help weigh treatment options (82,83). Similarly, knowledge of complex bifurcation lesions before cardiac catheterization may avoid high-risk interventions (80,81). Plaque imaging also may elucidate the effects of drugs on atherosclerotic disease (84). Individual plaque features may have particular significance in specific settings; for example, TCFAs may have different implications in patients with or without known susceptibility to vascular thrombosis. Thus, integration of lesion characteristics with risk factors may be valuable. Currently unknown features of atherosclerotic plaque may conceivably independently herald poor outcome. Advanced imaging techniques (e.g., molecular imaging) may elucidate such features and allow further insights into mechanisms of acute coronary event

pathophysiology (85). To determine truly independent risk prediction, any plaque assessment should be measured against the predictive power of atherosclerotic burden and its metabolic activity.

IMPLICATIONS FOR FUTURE INVESTIGATIONS

Although general morphologic patterns of atherosclerotic disease influence the probability of acute coronary syndromes, they are clearly modified by individual characteristics. Pathology and IVUS studies show that most United States adults older than 50 years have evidence of coronary atherosclerotic disease, but only a minority will experience acute events (42,86). Furthermore, the patterns and morphologic features of atherosclerotic disease appear similar among populations, suggesting that the patient's response to a thrombogenic trigger is critical for determining the probability of events. Traditional risk factors for coronary artery disease (e.g., diabetes, smoking, dyslipidemia) and genetic predisposition modify such responses. Several mutations are associated with increased event hazard, and individualized risk characterization may soon be available (87-89).

We need a better understanding of which combination of imaging information and risk factors yields the most accurate individual risk prediction. Research is needed to investigate mechanisms influencing the coagulation system's response to various internal and external modifiers, both locally and systemically. Specifically, we need to understand and potentially to predict the response of the coagulation system to stimuli occurring with atherosclerotic plaque alterations. Variability in the coagulation system's performance depends on numerous hormonal, dietary, and environmental influences, hampering our ability to predict its function at a given time, for example, when plaque ruptures (90-92). Thus, we must strive for comprehensive risk assessment that integrates specific information on the atherosclerotic plaque burden and systemic factors that increase the risk for disease activity and vascular thrombosis and is tailored to specific patient populations and individual patients. This would enable effective, efficient triaging of patients into treatment categories ranging from continued risk factor control to coronary arterial revascularization.

CONCLUSIONS

Despite major advancements in coronary artery imaging and identification of atherosclerotic lesion morphology associated with rupture, there is no conclusive evidence that individual plaque assessment better predicts acute coronary event risk than established risk factors, such as the extent

and severity of coronary artery disease. Pathology and clinical studies consistently demonstrate that atherosclerotic plaques rupture without clinical symptoms much more frequently than is widely acknowledged, challenging the notion of a close association between plaque rupture and clinical events. Conversely, the atherosclerotic disease burden is a consistent, strong predictor of adverse cardiovascular events and deserves greater attention. Current data suggest that rather than focusing on

individual coronary arterial lesions, we need a comprehensive, integrative approach for identifying and managing patients at risk of adverse cardiovascular events.

REPRINT REQUESTS AND CORRESPONDENCE: Dr. Armin Arbab-Zadeh, Johns Hopkins University, Division of Cardiology, 600 N. Wolfe Street, Blalock 524, Baltimore, Maryland 21287-8222. E-mail: azadeh1@jhmi.edu.

REFERENCES

1. Go AS, Mozaffarian D, Roger VL, et al., for the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics—2014 update: a report from the American Heart Association. *Circulation* 2014; 129:e28–292.
2. Franco M, Cooper RS, Bilal U, et al. Challenges and opportunities for cardiovascular disease prevention. *Am J Med* 2011;124:95–102.
3. Greenland P, Knoll MD, Stamler J, et al. Major risk factors as antecedents of fatal and nonfatal coronary heart disease events. *JAMA* 2003;290:891–7.
4. Young LH, Wackers FJ, Chyun DA, et al., for the DIAD Investigators. Cardiac outcomes after screening for asymptomatic coronary artery disease in patients with type 2 diabetes: the DIAD study: a randomized controlled trial. *JAMA* 2009; 301:1547–55.
5. Wilson PW, D'Agostino RB, Levy D, et al. Prediction of coronary heart disease using risk factor categories. *Circulation* 1998;97:1837–47.
6. Detrano R, Guerci AD, Carr JJ, et al. Coronary calcium as a predictor of coronary events in four racial or ethnic groups. *N Engl J Med* 2008;358: 1336–45.
7. Arbab-Zadeh A, Nakano M, Virmani R, et al. Acute coronary events. *Circulation* 2012;125:1147–56.
8. Bentzon JF, Otsuka F, Virmani R, et al. Mechanisms of plaque formation and rupture. *Circ Res* 2014;114:1852–66.
9. Ueda Y, Ogasawara N, Matsuo K, et al. Acute coronary syndrome: insight from angioscopy. *Circ J* 2010;74:411–7.
10. Naghavi M, Libby P, Falk E, et al. From vulnerable plaque to vulnerable patient: a call for new definitions and risk assessment strategies: part I. *Circulation* 2003;108:1664–72.
11. Naghavi M, Libby P, Falk E, et al. From vulnerable plaque to vulnerable patient: a call for new definitions and risk assessment strategies: part II. *Circulation* 2003;108:1772–8.
12. Fleg JL, Stone GW, Fayad ZA, et al. Detection of high-risk atherosclerotic plaque: report of the NHLBI working group on current status and future directions. *J Am Coll Cardiol Img* 2012;5:941–55.
13. Finn AV, Nakano M, Narula J, et al. Concept of vulnerable/unstable plaque. *Arterioscler Thromb Vasc Biol* 2010;30:1282–92.
14. Libby P. Mechanisms of acute coronary syndromes and their implications for therapy. *N Engl J Med* 2013;368:2004–13.
15. Virmani R, Burke AP, Farb A, et al. Pathology of the vulnerable plaque. *J Am Coll Cardiol* 2006;47 8 Suppl:C13–8.
16. Narula J, Garg P, Achenbach S, et al. Arithmetic of vulnerable plaques for noninvasive imaging. *Nat Clin Pract Cardiovasc Med* 2008; 5 Suppl 2:S2–10.
17. National Institutes of Health. NIH Research Portfolio Online Reporting Tools (RePORT). Available at: <http://report.nih.gov>. Accessed December 4, 2014.
18. Hlatky MA, Douglas PS, Cook NL, et al. Future directions for cardiovascular disease comparative effectiveness research: report of a workshop sponsored by the National Heart, Lung, and Blood Institute. *J Am Coll Cardiol* 2012;60:569–80.
19. Stone GW, Maehara A, Lansky AJ, et al., for the PROSPECT Investigators. A prospective natural-history study of coronary atherosclerosis. *N Engl J Med* 2011;364:226–35.
20. Calvert PA, Obaid DR, O'Sullivan M, et al. Association between IVUS findings and adverse outcomes in patients with coronary artery disease: the VIVA (VH-IVUS in Vulnerable Atherosclerosis) study. *J Am Coll Cardiol Img* 2011;4:894–901.
21. Yonetu T, Bouma BE, Kato K, et al. Optical coherence tomography—15 years in cardiology. *Circ J* 2013;77:1933–40.
22. Tian J, Ren X, Vergallo R, et al. Distinct morphological features of ruptured culprit plaque for acute coronary events compared to those with silent rupture and thin-cap fibroatheroma: a combined optical coherence tomography and intravascular ultrasound study. *J Am Coll Cardiol* 2014;63:2209–16.
23. Motoyama S, Sarai M, Harigaya H, et al. Computed tomographic angiography characteristics of atherosclerotic plaques subsequently resulting in acute coronary syndrome. *J Am Coll Cardiol* 2009;54:49–57.
24. Maurovich-Horvat P, Hoffmann U, Vorpahl M, et al. The napkin-ring sign: CT signature of high-risk coronary plaques? *J Am Coll Cardiol Img* 2010;3:440–4.
25. Otsuka K, Fukuda S, Tanaka A, et al. Napkin-ring sign on coronary CT angiography for the prediction of acute coronary syndrome. *J Am Coll Cardiol Img* 2013;6:448–57.
26. Puchner SB, Liu T, Mayrhofer T, et al. High-risk plaque detected on coronary CT angiography predicts acute coronary syndromes independent of significant stenosis in acute chest pain: Results from the ROMICAT-II trial. *J Am Coll Cardiol* 2014; 64:684–92.
27. Hellings WE, Peeters W, Moll FL, et al. Composition of carotid atherosclerotic plaque is associated with cardiovascular outcome: a prognostic study. *Circulation* 2010;121:1941–50.
28. Saam T, Hetterich H, Hoffmann V, et al. Meta-analysis and systematic review of the predictive value of carotid plaque hemorrhage on cerebrovascular events by magnetic resonance imaging. *J Am Coll Cardiol* 2013;62:1081–91.
29. Tanaka A, Shimada K, Sano T, et al. Multiple plaque rupture and C-reactive protein in acute myocardial infarction. *J Am Coll Cardiol* 2005;45: 1594–9.
30. Arbustini E, Grasso M, Diegoli M, et al. Coronary atherosclerotic plaques with and without thrombus in ischemic heart syndromes: a morphologic, immunohistochemical, and biochemical study. *Am J Cardiol* 1991;68:36B–50B.
31. Hudson J, McCaughey WT. Mural thrombosis and atherogenesis in coronary arteries and aorta. An investigation using antifibrin and antiplatelet sera. *Atherosclerosis* 1974;19:543–53.
32. Shimamura K, Ino Y, Kubo T, et al. Difference of ruptured plaque morphology between asymptomatic coronary artery disease and non-ST elevation acute coronary syndrome patients: an optical coherence tomography study. *Atherosclerosis* 2014;235:532–7.
33. Tanaka A, Imanishi T, Kitabata H, et al. Distribution and frequency of thin-capped fibroatheromas and ruptured plaques in the entire culprit coronary artery in patients with acute coronary syndrome as determined by optical coherence tomography. *Am J Cardiol* 2008;102:975–9.
34. Takano M, Inami S, Ishibashi F, et al. Angiographic follow-up study of coronary ruptured plaques in nonculprit lesions. *J Am Coll Cardiol* 2005; 45:652–8.
35. Hong MK, Mintz GS, Lee CW, et al. Comparison of coronary plaque rupture between stable angina and acute myocardial infarction: a three-vessel

- intravascular ultrasound study in 235 patients. *Circulation* 2004;110:928–33.
- 36.** Schoenhagen P, Stone GW, Nissen SE, et al. Coronary plaque morphology and frequency of ulceration distant from culprit lesions in patients with unstable and stable presentation. *Arterioscler Thromb Vasc Biol* 2003;23:1895–900.
- 37.** Kotani J, Mintz GS, Castagna MT, et al. Intravascular ultrasound analysis of infarct-related and non-infarct-related arteries in patients who presented with an acute myocardial infarction. *Circulation* 2003;107:2889–93.
- 38.** Cheruvu PK, Finn AV, Gardner C, et al. Frequency and distribution of thin-cap fibroatheroma and ruptured plaques in human coronary arteries: a pathologic study. *J Am Coll Cardiol* 2007;50:940–9.
- 39.** Burke AP, Kolodgie FD, Farb A, et al. Healed plaque ruptures and sudden coronary death: evidence that subclinical rupture has a role in plaque progression. *Circulation* 2001;103:934–40.
- 40.** Mann J, Davies MJ. Mechanisms of progression in native coronary artery disease: role of healed plaque disruption. *Heart* 1999;82:265–8.
- 41.** Rioufol G, Finet G, Ginon I, et al. Multiple atherosclerotic plaque rupture in acute coronary syndrome: a three-vessel intravascular ultrasound study. *Circulation* 2002;106:804–8.
- 42.** Nemetz PN, Roger VL, Ransom JE, et al. Recent trends in the prevalence of coronary disease: a population-based autopsy study of nonnatural deaths. *Arch Intern Med* 2008;168:264–70.
- 43.** Kubo T, Maehara A, Mintz GS, et al. The dynamic nature of coronary artery lesion morphology assessed by serial virtual histology intravascular ultrasound tissue characterization. *J Am Coll Cardiol* 2010;55:1590–7.
- 44.** Motreff P, Rioufol G, Finet G. Seventy-four-month follow-up of coronary vulnerable plaques by serial gray-scale intravascular ultrasound. *Circulation* 2012;126:2878–9.
- 45.** Uemura S, Ishigami K, Soeda T, et al. Thin-cap fibroatheroma and microchannel findings in optical coherence tomography correlate with subsequent progression of coronary atherosomatous plaques. *Eur Heart J* 2012;33:78–85.
- 46.** Tian J, Dauerman H, Toma C, et al. Prevalence and characteristics of TCFA and degree of coronary artery stenosis: an OCT, IVUS, and angiographic study. *J Am Coll Cardiol* 2014;64:672–80.
- 47.** Narula J, Kovacic JC. Putting TCFA in clinical perspective. *J Am Coll Cardiol* 2014;64:681–3.
- 48.** Budoff MJ, Shaw LJ, Liu ST, et al. Long-term prognosis associated with coronary calcification: observations from a registry of 25,253 patients. *J Am Coll Cardiol* 2007;49:1860–70.
- 49.** Budoff MJ, Hokanson JE, Nasir K, et al. Progression of coronary artery calcium predicts all-cause mortality. *J Am Coll Cardio Img* 2010;3:1229–36.
- 50.** Yokoya K, Takatsu H, Suzuki T, et al. Process of progression of coronary artery lesions from mild or moderate stenosis to moderate or severe stenosis: a study based on four serial coronary arteriograms per year. *Circulation* 1999;100:903–9.
- 51.** Yusuf S, Zucker D, Peduzzi P, et al. Effect of coronary artery bypass graft surgery on survival: overview of 10-year results from randomised trials by the Coronary Artery Bypass Graft Surgery Trialists Collaboration. *Lancet* 1994;344:563–70.
- 52.** Cho I, Chang HJ, Sung JM, et al., for the CONFIRM Investigators. Coronary computed tomographic angiography and risk of all-cause mortality and nonfatal myocardial infarction in subjects without chest pain syndrome from the CONFIRM registry (Coronary CT Angiography Evaluation for Clinical Outcomes: An International Multicenter Registry). *Circulation* 2012;126:304–13.
- 53.** Baber U, Mehran R, Sartori S, et al. Detection and impact of subclinical coronary and carotid atherosclerosis on cardiovascular risk prediction and reclassification in asymptomatic US adults: insights from the high risk plaque BioImage study (abstr). *J Am Coll Cardiol* 2014;63:A998.
- 54.** Collins R, Armitage J, Parish S, et al., for the Heart Protection Study Collaborative Group. Effects of cholesterol-lowering with simvastatin on stroke and other major vascular events in 20536 people with cerebrovascular disease or other high-risk conditions. *Lancet* 2004;363:757–67.
- 55.** Cannon CP, Braunwald E, McCabe CH, et al., for the Pravastatin or Atorvastatin Evaluation and Infection Therapy-Thrombolysis in Myocardial Infarction 22 Investigators. Intensive versus moderate lipid lowering with statins after acute coronary syndromes. *N Engl J Med* 2004;350:1495–504.
- 56.** Berger JS, Brown DL, Becker RC. Low-dose aspirin in patients with stable cardiovascular disease: a meta-analysis. *Am J Med* 2008;121:43–9.
- 57.** Stergiopoulos K, Brown DL. Initial coronary stent implantation with medical therapy vs medical therapy alone for stable coronary artery disease: meta-analysis of randomized controlled trials. *Arch Intern Med* 2012;172:312–9.
- 58.** Boden WE. Mounting evidence for lack of PCI benefit in stable ischemic heart disease: what more will it take to turn the tide of treatment? Comment on "Initial coronary stent implantation with medical therapy vs medical therapy alone for stable coronary artery disease." *Arch Intern Med* 2012;172:319–21.
- 59.** Stergiopoulos K, Boden WE, Hartigan P, et al. Percutaneous coronary intervention outcomes in patients with stable obstructive coronary artery disease and myocardial ischemia: a collaborative meta-analysis of contemporary randomized clinical trials. *JAMA Intern Med* 2014;174:232–40.
- 60.** Tonino PA, De Bruyne B, Pijls NH, et al., for the FAME Study Investigators. Fractional flow reserve versus angiography for guiding percutaneous coronary intervention. *N Engl J Med* 2009;360:213–24.
- 61.** Arbab-Zadeh A. Fractional flow reserve guided percutaneous coronary intervention is not a valid concept. *Circulation* 2014;129:1871–8.
- 62.** De Bruyne B, Fearon WF, Pijls NH, et al., for the FAME 2 Trial Investigators. Fractional flow reserve-guided PCI for stable coronary artery disease. *N Engl J Med* 2014;371:1208–17.
- 63.** Niccoli G, Stefanini GG, Capodanno D, et al. Are the culprit lesions severely stenotic? *J Am Coll Cardiol Img* 2013;6:1108–14.
- 64.** Little WC, Constantinescu M, Applegate RJ, et al. Can coronary angiography predict the site of a subsequent myocardial infarction in patients with mild-to-moderate coronary artery disease? *Circulation* 1988;78:1157–66.
- 65.** Fishbein MC, Siegel RJ. How big are coronary atherosclerotic plaques that rupture? *Circulation* 1996;94:2662–6.
- 66.** Ojio S, Takatsu H, Tanaka T, et al. Considerable time from the onset of plaque rupture and/or thrombi until the onset of acute myocardial infarction in humans: coronary angiographic findings within 1 week before the onset of infarction. *Circulation* 2000;102:2063–9.
- 67.** Frobert O, van't Veer M, Aarnoudse W, et al. Acute myocardial infarction and underlying stenosis severity. *Catheter Cardiovasc Interv* 2007;70:958–65.
- 68.** Davies MJ, Thomas A. Thrombosis and acute coronary-artery lesions in sudden cardiac ischemic death. *N Engl J Med* 1984;310:1137–40.
- 69.** Farb A, Burke AP, Tang AL, et al. Coronary plaque erosion without rupture into a lipid core. A frequent cause of coronary thrombosis in sudden coronary death. *Circulation* 1996;93:1354–63.
- 70.** Bittencourt MS, Hulten E, Ghoshhajra B, et al. Prognostic value of nonobstructive and obstructive coronary artery disease detected by coronary computed tomography angiography to identify cardiovascular events. *Circ Cardiovasc Imaging* 2014;7:282–91.
- 71.** Rumberger JA, Simons DB, Fitzpatrick LA, et al. Coronary artery calcium area by electron-beam computed tomography and coronary atherosclerotic plaque area. A histopathologic correlative study. *Circulation* 1995;92:2157–62.
- 72.** Boogers MJ, Broersen A, van Velzen JE, et al. Automated quantification of coronary plaque with computed tomography: comparison with intravascular ultrasound using a dedicated registration algorithm for fusion-based quantification. *Eur Heart J* 2012;33:1007–16.
- 73.** Nicholls SJ, Hsu A, Wolski K, et al. Intravascular ultrasound-derived measures of coronary atherosclerotic plaque burden and clinical outcome. *J Am Coll Cardiol* 2010;55:2399–407.
- 74.** Mancini GB, Hartigan PM, Shaw LJ, et al. Predicting outcome in the COURAGE trial (Clinical Outcomes Utilizing Revascularization and Aggressive Drug Evaluation): coronary anatomy versus ischemia. *J Am Coll Cardiol Intv* 2014;7:195–201.
- 75.** Wald DS, Morris JK, Wald NJ, et al., for the PRAMI Investigators. Randomized trial of preventive angioplasty in myocardial infarction. *N Engl J Med* 2013;369:1115–23.
- 76.** Kelly DJ, McCann GP, Blackman D, et al. Complete Versus Culprit-Lesion Only Primary PCI Trial (CVLPRIT): a multicentre trial testing management strategies when multivessel disease is detected at the time of primary PCI: rationale and design. *Eurointervention* 2013;8:1190–8.
- 77.** Bainey KR, Mehta SR, Lai T, et al. Complete vs culprit-only revascularization for patients with

- multivessel disease undergoing primary percutaneous coronary intervention for ST-segment elevation myocardial infarction: a systematic review and meta-analysis. *Am Heart J* 2014;167:1-14.e2.
- 78.** Complete vs Culprit-Only Revascularization to Treat Multi-Vessel Disease After Primary PCI for STEMI (COMPLETE). Available at: <http://clinicaltrials.gov/ct2/show/NCT01740479>. Accessed December 4, 2014.
- 79.** van Lammeren GW, den Ruijter HM, Vrijenhoek JE, et al. Time-dependent changes in atherosclerotic plaque composition in patients undergoing carotid surgery. *Circulation* 2014;129:2269-76.
- 80.** McDaniel MC, Eshtehardi P, Sawaya FJ, et al. Contemporary clinical applications of coronary intravascular ultrasound. *J Am Coll Cardiol Intv* 2011;4:1155-67.
- 81.** Rodriguez-Granillo GA, Rosales MA, Llaurado C, et al. Guidance of percutaneous coronary interventions by multidetector row computed tomography coronary angiography. *Eurointervention* 2011;6:773-8.
- 82.** Madhavan MV, Tarigopula M, Mintz GS, et al. Coronary artery calcification: pathogenesis and prognostic implications. *J Am Coll Cardiol* 2014;63:1703-14.
- 83.** Ertelet K, Genereux P, Mintz GS, et al. Impact of the severity of coronary artery calcification on clinical events in patients undergoing coronary artery bypass grafting (from the Acute Catheterization and Urgent Intervention Triage Strategy Trial). *Am J Cardiol* 2013;112:1730-7.
- 84.** Hattori K, Ozaki Y, Ismail TF, et al. Impact of statin therapy on plaque characteristics as assessed by serial OCT, grayscale and integrated backscatter-IVUS. *J Am Coll Cardiol Img* 2012;5:169-77.
- 85.** Sadat U, Jaffer FA, van Zandvoort MA, et al. Inflammation and neovascularization intertwined in atherosclerosis: imaging of structural and molecular imaging targets. *Circulation* 2014;130:786-94.
- 86.** Tuzcu EM, Kapadia SR, Tutar E, et al. High prevalence of coronary atherosclerosis in asymptomatic teenagers and young adults: evidence from intravascular ultrasound. *Circulation* 2001;103:2705-10.
- 87.** Damani SB, Topol EJ. Emerging genomic applications in coronary artery disease. *J Am Coll Cardiol Intv* 2011;4:473-82.
- 88.** Taylor M, Farrall M, Holliday EG, et al., for the Australian Stroke Genetics Collaborative, Wellcome Trust Case Control Consortium 2 (WTCCC2), et al., International Stroke Genetics Consortium. Genetic risk factors for ischaemic stroke and its subtypes (the METASTROKE collaboration): a meta-analysis of genome-wide association studies. *Lancet Neurol* 2012;11:951-62.
- 89.** Makinen VP, Civelek M, Meng Q, et al., for the Coronary Artery Disease Genome-Wide Replication and Meta-Analysis (CARDIoGRAM) Consortium, et al. Integrative genomics reveals novel molecular pathways and gene networks for coronary artery disease. *PLoS Genet* 2014;10:e1004502.
- 90.** Montagnana M, Salvagno GL, Lippi G. Circadian variation within hemostasis: an underrecognized link between biology and disease? *Semin Thromb Hemost* 2009;35:23-33.
- 91.** Phang M, Lazarus S, Wood LG, et al. Diet and thrombosis risk: nutrients for prevention of thrombotic disease. *Semin Thromb Hemost* 2011;37:199-208.
- 92.** Servoss SJ, Januzzi JL, Muller JE. Triggers of acute coronary syndromes. *Prog Cardiovasc Dis* 2002;44:369-80.

KEY WORDS atherosclerosis, coronary heart disease, coronary imaging, prevention