



Weighted Gcd-Sum Functions

László Tóth ¹

Department of Mathematics

University of Pécs

Ifjúság u. 6

7624 Pécs

Hungary

and

Institute of Mathematics, Department of Integrative Biology

Universität für Bodenkultur

Gregor Mendel-Straße 33

A-1180 Wien

Austria

ltoth@gamma.ttk.pte.hu

Abstract

We investigate weighted gcd-sum functions, including the alternating gcd-sum function and those having as weights the binomial coefficients and values of the Gamma function. We also consider the alternating lcm-sum function.

1 Introduction

The gcd-sum function, called also Pillai's arithmetical function (OEIS [A018804](#)) is defined by

$$P(n) := \sum_{k=1}^n \gcd(k, n) \quad (n \in \mathbb{N} := \{1, 2, \dots\}). \quad (1)$$

The function P is multiplicative and its arithmetical and analytical properties are determined by the representation

$$P(n) = \sum_{d|n} d \phi(n/d) \quad (n \in \mathbb{N}), \quad (2)$$

¹The author gratefully acknowledges support from the Austrian Science Fund (FWF) under the project Nr. P20847-N18.

where ϕ is Euler's function. See the survey paper [5]. Note that for every prime power p^a ($a \in \mathbb{N}$),

$$P(p^a) = (a + 1)p^a - ap^{a-1}. \quad (3)$$

Now let

$$P_{\text{altern}}(n) := \sum_{k=1}^n (-1)^{k-1} \gcd(k, n) \quad (n \in \mathbb{N}) \quad (4)$$

be the alternating gcd-sum function. As far as we know, the function (4) was not considered before.

Furthermore, let

$$P_{\text{binom}}(n) := \sum_{k=1}^n \binom{n}{k} \gcd(k, n) \quad (n \in \mathbb{N}) \quad (5)$$

(OEIS [A159068](#)), where $\binom{n}{k}$ are the binomial coefficients. Every term of the sum (5) is a multiple of n , since $\gcd(k, n) = kx + ny$ with suitable integers x, y and $k \binom{n}{k} = n \binom{n-1}{k-1}$ ($1 \leq k \leq n$). Note also the symmetry $\binom{n}{k} \gcd(k, n) = \binom{n}{n-k} \gcd(n-k, n)$ ($1 \leq k \leq n-1$).

More generally, consider the weighted gcd-sum functions defined by

$$P_w(n) := \sum_{k=1}^n w(k, n) \gcd(k, n) \quad (n \in \mathbb{N}), \quad (6)$$

where the weights are functions $w : \mathbb{N}^2 \rightarrow \mathbb{C}$.

In this paper we evaluate the alternating gcd-sum function $P_{\text{altern}}(n)$, deduce a formula for the function $P_{\text{binom}}(n)$ and investigate other special cases of (6). We also give a formula for the alternating lcm-sum function defined by

$$L_{\text{altern}}(n) := \sum_{k=1}^n (-1)^{k-1} \text{lcm}[k, n] \quad (n \in \mathbb{N}). \quad (7)$$

Similar results can be derived for the weighted versions of certain analogs and generalizations of the gcd-sum function, see [5], but we confine ourselves to the function (6).

2 General results

We first give the following simple result.

Proposition 1. *i) Let $w : \mathbb{N}^2 \rightarrow \mathbb{C}$ be an arbitrary function. Then*

$$P_w(n) = \sum_{d|n} \phi(d) \sum_{j=1}^{n/d} w(dj, n) \quad (n \in \mathbb{N}). \quad (8)$$

ii) Assume that there is a function $g : (0, 1] \rightarrow \mathbb{C}$ such that $w(k, n) = g(k/n)$ ($1 \leq k \leq n$) and let $G(n) = \sum_{k=1}^n g(k/n)$ ($n \in \mathbb{N}$). Then

$$P_w(n) = \sum_{d|n} \phi(d) G(n/d) \quad (n \in \mathbb{N}). \quad (9)$$

Proof. i) Using Gauss' formula $m = \sum_{d|m} \phi(d)$ for $m = \gcd(k, n)$, grouping the terms of (6) and denoting $k = dj$ we obtain at once

$$P_w(n) := \sum_{k=1}^n w(k, n) \sum_{d|\gcd(k, n)} \phi(d) = \sum_{d|n} \phi(d) \sum_{j=1}^{n/d} w(dj, n).$$

ii) Use (8) and that

$$\sum_{j=1}^{n/d} w(dj, n) = \sum_{j=1}^{n/d} g(dj/n) = \sum_{j=1}^{n/d} g(j/(n/d)) = G(n/d).$$

□

For $w(k, n) = 1$ we reobtain formula (2). In the next section we investigate other special cases, including those already mentioned in the Introduction.

Remark 2. Consider the function

$$R_w(n) := \sum_{\substack{k=1 \\ \gcd(k, n)=1}}^n w(k, n) \quad (n \in \mathbb{N}). \quad (10)$$

Then, similar to the proof of i), now with the Möbius μ function instead of ϕ ,

$$R_w(n) = \sum_{k=1}^n w(k, n) \sum_{d|\gcd(k, n)} \mu(d) = \sum_{d|n} \mu(d) \sum_{j=1}^{n/d} w(dj, n). \quad (11)$$

If condition ii) is satisfied, then we have

$$R_w(n) = \sum_{d|n} \mu(d) G(n/d) \quad (n \in \mathbb{N}). \quad (12)$$

We will also point out some special cases of (11) and (12).

3 Special cases

3.1 Alternating gcd-sum function

Let $w(k, n) = (-1)^{k-1}$ ($k, n \in \mathbb{N}$). Then we have the function $P_{\text{altern}}(n)$ defined by (4).

Proposition 3. *Let $n \in \mathbb{N}$ and write $n = 2^a m$, where $a \in \mathbb{N}_0 := \{0, 1, 2, \dots\}$ and $m \in \mathbb{N}$ is odd. Then*

$$P_{\text{altern}}(n) = \begin{cases} n, & \text{if } n \text{ is odd } (a = 0); \\ -2^{a-1} a P(m) = -\frac{a}{a+2} P(n), & \text{if } n \text{ is even } (a \geq 1). \end{cases} \quad (13)$$

Proof. Use formula (8). Here

$$S_d(n) := \sum_{j=1}^{n/d} w(dj, n) = \sum_{j=1}^{n/d} (-1)^{dj-1} = - \sum_{j=1}^{n/d} (-1)^{dj}.$$

If n is odd, then every divisor d of n is also odd and obtain $S_d(n) = - \sum_{j=1}^{n/d} (-1)^j = 1$, where n/d is odd. Hence, $P_{\text{altern}}(n) = \sum_{d|n} \phi(d) = n$.

Now let n be even and let $d | n$. For d odd, $S_d(n) = - \sum_{j=1}^{n/d} (-1)^j = 0$, since n/d is even. For d even, $S_d(n) = - \sum_{j=1}^{n/d} 1 = -n/d$. We obtain that

$$P_{\text{altern}}(n) = - \sum_{\substack{d|n \\ d \text{ even}}} \phi(d) \frac{n}{d} = - \sum_{d|n} \phi(d) \frac{n}{d} + \sum_{\substack{d|n \\ d \text{ odd}}} \phi(d) \frac{n}{d},$$

where the first sum is $P(n)$ (cf. (2)), and the second one is

$$\sum_{d|m} \phi(d) \frac{2^a m}{d} = 2^a P(m).$$

Using (3), $P(n) = P(2^a)P(m) = 2^{a-1}(a+2)P(m)$ and deduce

$$\begin{aligned} P_{\text{altern}}(n) &= -P(n) + 2^a P(m) = P(m)(2^a - 2^{a-1}(a+2)) \\ &= -2^{a-1}aP(m) = -\frac{a}{a+2}P(n). \end{aligned}$$

□

Remark 4. More generally, consider the polynomial

$$f_n(x) := \sum_{k=1}^n \gcd(k, n)x^{k-1}, \quad (14)$$

i.e., put $w(k, n) = x^{k-1}$ (formally). Then $f_n(1) = P(n)$, $f_n(-1) = P_{\text{altern}}(n)$ and deduce from Proposition 1,

$$f_n(x) := (1-x^n) \sum_{d|n} \frac{\phi(d)x^{d-1}}{1-x^d}. \quad (15)$$

3.2 Logarithms as weights

Let

$$P_{\log}(n) := \sum_{k=1}^n (\log k) \gcd(k, n). \quad (16)$$

Proposition 5. For every $n \in \mathbb{N}$,

$$P_{\log}(n) = P(n) \log n + \sum_{d|n} \log(d!/d^d) \phi(n/d). \quad (17)$$

Proof. Apply formula (8). For $w(k, n) = \log k$,

$$\sum_{j=1}^{n/d} w(dj, n) = \sum_{j=1}^{n/d} \log(dj) = \frac{n}{d} \log d + \log \left(\frac{n}{d} \right)!,$$

hence

$$P_{\log}(n) = \sum_{d|n} \phi(d) \left(\frac{n}{d} \log d + \log \left(\frac{n}{d} \right)! \right),$$

and a short computation leads to (17). \square

Remark 6. Writing the exponential form of (17),

$$\prod_{k=1}^n k^{\gcd(k, n)} = n^{P(n)} \prod_{d|n} \left(\frac{d!}{d^d} \right)^{\phi(n/d)}. \quad (18)$$

Compare this to the known formula

$$\prod_{\substack{k=1 \\ \gcd(k, n)=1}}^n k = n^{\phi(n)} \prod_{d|n} \left(\frac{d!}{d^d} \right)^{\mu(n/d)}, \quad (19)$$

cf. [2, p. 197, Ex. 24] (OEIS [A001783](#)).

3.3 Discrete Fourier transform of the gcd's

Consider $w(k, n) = \exp(2\pi ikr/n)$ ($k, n \in \mathbb{N}$), where $r \in \mathbb{Z}$, and denote

$$P_{\text{DFT}}^{(r)}(n) := \sum_{k=1}^n \exp(2\pi ikr/n) \gcd(k, n), \quad (20)$$

representing the discrete Fourier transform of the function $f(k) = \gcd(k, n)$ ($k \in \mathbb{N}$).

Proposition 7. *For every $n \in \mathbb{N}$, $r \in \mathbb{Z}$,*

$$P_{\text{DFT}}^{(r)}(n) = \sum_{d|\gcd(n, r)} d \phi(n/d). \quad (21)$$

Proof. Here $\exp(2\pi ikr/n) = g(k/n)$ with $g(x) = \exp(2\pi irx)$. Using formula (9) and that

$$\sum_{k=1}^n \exp(2\pi irk/n) = \begin{cases} n, & \text{if } n \mid r; \\ 0, & \text{otherwise;} \end{cases}$$

we obtain

$$P_{\text{DFT}}^{(r)}(n) = \sum_{d|n, n/d|r} \phi(d) \frac{n}{d} = \sum_{d|n, d|r} d \phi(n/d).$$

\square

Remark 8. Formula (21) can be written in the form

$$P_{\text{DFT}}^{(r)}(n) = \sum_{d|n} dc_{n/d}(r), \quad (22)$$

where $c_n(k)$ are the Ramanujan sums. Furthermore, (22) can be extended for r -even functions. See [4], [6, Prop. 2]. Note that in the present treatment we do not need properties of the Ramanujan sums and of r -even functions.

For $r = 0$ (more generally, in case $n \mid r$) we reobtain from (21) formula (2). For $r = 1$ we deduce

$$\sum_{k=1}^n \exp(2\pi ik/n) \gcd(k, n) = \phi(n) \quad (n \in \mathbb{N}), \quad (23)$$

which gives by writing the real and the imaginary parts, respectively,

$$\sum_{k=1}^n \cos(2\pi k/n) \gcd(k, n) = \phi(n) \quad (n \in \mathbb{N}), \quad (24)$$

$$\sum_{k=1}^n \sin(2\pi k/n) \gcd(k, n) = 0 \quad (n \in \mathbb{N}), \quad (25)$$

similar relations being valid for $\gcd(n, r) = 1$.

Formulae (23), (24), (25) were pointed out in [4, Ex. 3].

3.4 Binomial coefficients as weights

Let $w(k, n) = \binom{n}{k}$ ($k, n \in \mathbb{N}$). Then we have the function $P_{\text{binom}}(n)$ defined by (5).

Proposition 9. *For every $n \in \mathbb{N}$,*

$$P_{\text{binom}}(n) = 2^n \sum_{d|n} \frac{\phi(d)}{d} \sum_{\ell=1}^d (-1)^\ell \cos^n(\ell\pi/d) - n. \quad (26)$$

Proof. Let $\varepsilon_r^j = \exp(2\pi ij/r)$ ($1 \leq j \leq r$) denote the r -th roots of unity. Using the known identity

$$\sum_{k=0}^{\lfloor n/r \rfloor} \binom{n}{kr} = \frac{1}{r} \sum_{j=1}^r (1 + \varepsilon_r^j)^n = \frac{2^n}{r} \sum_{j=1}^r \cos^n(j\pi/r) \cos(nj\pi/r) \quad (n, r \in \mathbb{N}), \quad (27)$$

cf. [1, p. 84], and applying (8) we deduce

$$\begin{aligned} P_{\text{binom}}(n) &= \sum_{d|n} \phi(d) \sum_{j=1}^{n/d} \binom{n}{dj} = \sum_{d|n} \phi(d) \left(\frac{2^n}{d} \sum_{\ell=1}^d \cos^n(\ell\pi/d) \cos(n\ell\pi/d) - 1 \right) \\ &= 2^n \sum_{d|n} \frac{\phi(d)}{d} \sum_{\ell=1}^d (-1)^\ell \cos^n(\ell\pi/d) - \sum_{d|n} \phi(d), \end{aligned}$$

giving (26). □

Note that (11) and (27) lead to the following formula for the sequence OEIS [A056188](#):

$$R_{\text{binom}}(n) := \sum_{\substack{k=1 \\ \gcd(k,n)=1}}^n \binom{n}{k} = 2^n \sum_{d|n} \frac{\mu(d)}{d} \sum_{\ell=1}^d (-1)^\ell \cos^n(\ell\pi/d) \quad (n > 1). \quad (28)$$

3.5 Weights concerning the Gamma function

Now let

$$P_{\text{Gamma}}(n) := \sum_{k=1}^n \log \Gamma\left(\frac{k}{n}\right) \gcd(k, n), \quad (29)$$

where Γ is the Gamma function.

Proposition 10. *For every $n \in \mathbb{N}$,*

$$P_{\text{Gamma}}(n) = \frac{\log 2\pi}{2} (P(n) - n) - \frac{1}{2} n \log n + \frac{1}{2} \sum_{d|n} \phi(d) \log d. \quad (30)$$

Proof. This follows by (9) and by

$$\prod_{k=1}^n \Gamma\left(\frac{k}{n}\right) = (2\pi)^{(n-1)/2} n^{-1/2}, \quad (n \in \mathbb{N}),$$

which is a consequence of Gauss' multiplication formula. □

Remark 11. (30) can be written also as

$$P_{\text{Gamma}}(n) = \frac{\log 2\pi}{2} (P(n) - n) - \frac{1}{2} (\phi * \log)(n), \quad (31)$$

where $*$ denotes the Dirichlet convolution. Note that $\phi * \log = \mu * \text{id} * \log = \Lambda * \text{id}$, where $\text{id}(n) = n$ ($n \in \mathbb{N}$) and Λ is the von Mangoldt function.

Writing the exponential form,

$$\prod_{k=1}^n \left(\Gamma\left(\frac{k}{n}\right) \right)^{\gcd(k,n)} = (2\pi)^{(P(n)-n)/2} n^{-n/2} \prod_{d|n} d^{\phi(d)/2}. \quad (32)$$

Compare this to the following formula given in [3]:

$$\prod_{\substack{k=1 \\ \gcd(k,n)=1}}^n \Gamma\left(\frac{k}{n}\right) = \frac{(2\pi)^{\phi(n)/2}}{\exp(\Lambda(n)/2)} = \begin{cases} (2\pi)^{\phi(n)/2} / \sqrt{p}, & n = p^a \text{ (a prime power);} \\ (2\pi)^{\phi(n)/2}, & \text{otherwise.} \end{cases} \quad (33)$$

3.6 Further special cases

It is possible to investigate other special cases, too. As examples we give the next ones with weights regarding, among others, the floor function $\lfloor \cdot \rfloor$, and the saw-tooth function ψ defined as $\psi(x) = x - \lfloor x \rfloor - 1/2$ for $x \in \mathbb{R} \setminus \mathbb{Z}$ and $\psi(x) = 0$ for $x \in \mathbb{Z}$.

Proposition 12. *For every $n \in \mathbb{N}$,*

$$P_{\text{id}}(n) := \sum_{k=1}^n k \gcd(k, n) = \frac{n}{2}(P(n) + n). \quad (34)$$

Proposition 13. *For every $n \in \mathbb{N}$ and $\alpha \in \mathbb{R}$,*

$$P_{\text{floor}}(n) := \sum_{k=1}^n \left\lfloor \alpha + \frac{k}{n} \right\rfloor \gcd(k, n) = \sum_{d|n} \phi(d) \left\lfloor \frac{n\alpha}{d} \right\rfloor. \quad (35)$$

Proposition 14. *For every $n, r \in \mathbb{N}$,*

$$P_{\text{saw-tooth}}^{(r)}(n) := \sum_{k=1}^n \psi(kr/n) \gcd(k, n) = 0. \quad (36)$$

Proposition 15. *For every $n \in \mathbb{N}, n > 1$,*

$$P_{\text{sin}}(n) := \sum_{k=1}^{n-1} (\log \sin(k\pi/n)) \gcd(k, n) = (\phi * \log)(n) - (\log 2)(P(n) - n). \quad (37)$$

Proposition 16. *For every $n \in \mathbb{N}$ and $\alpha \in \mathbb{R}$ with $\alpha + k/n \notin \mathbb{Z}$ ($1 \leq k \leq n$),*

$$P_{\text{cot}}(n) := \sum_{k=1}^n \cot \pi(\alpha + k/n) \gcd(k, n) = n \sum_{d|n} \frac{\phi(d)}{d} \cot(\pi n\alpha/d). \quad (38)$$

These follow from Proposition 1 using the following well-known formulae:

$$\sum_{k=1}^n \left\lfloor \alpha + \frac{k}{n} \right\rfloor = \lfloor n\alpha \rfloor, \quad (n \in \mathbb{N}), \quad (39)$$

$$\sum_{k=1}^n \psi(kr/n) = 0 \quad (n, r \in \mathbb{N}), \quad (40)$$

$$\prod_{k=1}^{n-1} \sin(k\pi/n) = \frac{n}{2^{n-1}} \quad (n \in \mathbb{N}) \quad (41)$$

(for $n = 1$ the empty product is 1),

$$\sum_{k=1}^n \cot \pi(\alpha + k/n) = n \cot \pi n\alpha \quad (n \in \mathbb{N}, \alpha \in \mathbb{R}, \alpha + k/n \notin \mathbb{Z}, 1 \leq k \leq n). \quad (42)$$

4 The alternating lcm-sum function

Some of the previous results have counterparts for the lcm-sum function (OEIS [A051193](#))

$$L(n) := \sum_{k=1}^n \text{lcm}[k, n] = \frac{n}{2} \left(1 + \sum_{d|n} d\phi(d) \right) \quad (n \in \mathbb{N}). \quad (43)$$

We consider here the alternating lcm-sum function defined by (7) and then the analog of (18).

Let $F(n) := \frac{1}{n} \sum_{d|n} d\phi(d)$. Note that $F(n) = \sum_{k=1}^n (\text{gcd}(k, n))^{-1}$ representing the arithmetic mean of the orders of elements in the cyclic group of order n , cf. [5, p. 3]. Furthermore, let $\beta(n) := (\mathbf{1} * \mu \text{id})(n) = \prod_{d|n} (1 - p)$ and let $h(n) := \prod_{k=1}^n k^k$ be the sequence of hyperfactorials (OEIS [A002109](#)).

Proposition 17. *Let $n \in \mathbb{N}$. If n is odd, then*

$$L_{\text{altern}}(n) = \frac{n}{2} \left(1 + \sum_{d|n} d\mu(d)\tau(n/d) \right) = \frac{n}{2} \left(1 + \prod_{p^a||n} (a(1-p) + 1) \right), \quad (44)$$

where τ is the divisor function.

If n is even of the form $n = 2^a m$, where $a \geq 1$ and $m \in \mathbb{N}$ is odd, then

$$L_{\text{altern}}(n) = 2^{a-1} m \left(\frac{2^{2a} - 1}{3} m F(m) - 1 \right) = \frac{n}{2} \left(\frac{2^{2a} - 1}{2^{2a+1} + 1} n F(n) - 1 \right). \quad (45)$$

Proof. Let $\text{id}_{-1}(n) = n^{-1}$ and $\mathbf{1}(n) = 1$ ($n \in \mathbb{N}$). We have

$$\begin{aligned} L_{\text{altern}}(n) &= n \sum_{k=1}^n (-1)^{k-1} k \frac{1}{\text{gcd}(k, n)} = n \sum_{k=1}^n (-1)^{k-1} k \sum_{d|\text{gcd}(k, n)} (\text{id}_{-1} * \mu)(d) \\ &= n \sum_{d|n} \beta(d) \sum_{j=1}^{n/d} (-1)^{dj-1} j. \end{aligned}$$

Now using that $\sum_{k=1}^n (-1)^{k-1} k = (-1)^{n-1} \lfloor (n+1)/2 \rfloor$ ($n \in \mathbb{N}$) the given formulae are obtained along the same lines with the proof of Proposition 3. \square

Proposition 18. *For every $n \in \mathbb{N}$,*

$$\left(\prod_{k=1}^n k^{\text{lcm}[k, n]} \right)^{1/n} = \prod_{d|n} h(n/d)^{\beta(d)} \left(\prod_{d|n} d^{\beta(d)/d} \right)^{n/2} \left(\prod_{d|n} d^{\beta(d)/d^2} \right)^{n^2/2}. \quad (46)$$

Proof. Similar to the proofs of above,

$$\begin{aligned} \sum_{k=1}^n (\log k) \operatorname{lcm}[k, n] &= n \sum_{k=1}^n (k \log k) \frac{1}{\operatorname{gcd}(k, n)} \\ &= n \sum_{k=1}^n (k \log k) \sum_{d|\operatorname{gcd}(k, n)} (\operatorname{id}_{-1} * \mu)(d) = n \sum_{d|n} (\operatorname{id}_{-1} * \mu)(d) \sum_{j=1}^{n/d} jd \log(jd) \\ &= n \sum_{d|n} \beta(d) \log h(n/d) + \frac{n^2}{2} \sum_{d|n} \beta(d) \frac{\log d}{d} + \frac{n^3}{2} \sum_{d|n} \beta(d) \frac{\log d}{d^2}, \end{aligned}$$

equivalent to (46). □

References

- [1] L. Comtet, *Advanced Combinatorics. The Art of Finite and Infinite Expansions*, D. Reidel Publishing Co., 1974.
- [2] I. Niven, H. S. Zuckerman and H. L. Montgomery, *An Introduction to the Theory of Numbers*, 5th ed., John Wiley & Sons, 1991.
- [3] J. Sándor and L. Tóth, A remark on the gamma function, *Elem. Math.* **44** (1989), 73–76.
- [4] W. Schramm, The Fourier transform of functions of the greatest common divisor, *Integers* **8** (2008), #A50.
- [5] L. Tóth, A survey of gcd-sum functions, *J. Integer Sequences* **13** (2010), [Article 109.8.1](#).
- [6] L. Tóth and P. Haukkanen, The discrete Fourier transform of r -even functions, submitted, <http://arxiv.org/abs/1009.5281v1>

2010 *Mathematics Subject Classification*: Primary 11A25; Secondary 05A10, 33B15.

Keywords: gcd-sum function, lcm-sum function, Euler’s function, Möbius function, binomial coefficient, Gamma function.

(Concerned with sequences [A001783](#), [A002109](#), [A018804](#), [A051193](#), [A056188](#), and [A159068](#).)

Received May 11 2011; revised version received July 18 2011. Published in *Journal of Integer Sequences*, September 5 2011.

Return to [Journal of Integer Sequences home page](#).