

# Deep glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet

Mathieu MORLIGHEM<sup>1</sup>, Eric RIGNOT<sup>1,2</sup>, Tobias BINDER<sup>3</sup>, Donald BLANKENSHIP<sup>4</sup>, Reinhard DREWS<sup>3,5</sup>, Graeme EAGLES<sup>3</sup>, Olaf EISEN<sup>3,6</sup>, Fausto FERRACCIOLI<sup>7</sup>, René FORSBERG<sup>8</sup>, Peter FRETWELL<sup>7</sup>, Vikram GOEL<sup>9</sup>, Jamin S. GREENBAUM<sup>4</sup>, Hilmar GUDMUNDSSON<sup>10</sup>, Jingxue GUO<sup>11</sup>, Veit HELM<sup>3</sup>, Coen HOFSTEDÉ<sup>3</sup>, Ian HOWAT<sup>12</sup>, Angelika HUMBERT<sup>3,6</sup>, Wilfried JOKAT<sup>3</sup>, Nanna B. KARLSSON<sup>3,13</sup>, Won Sang LEE<sup>14</sup>, Kenichi MATSUOKA<sup>15</sup>, Romain MILLAN<sup>1</sup>, Jeremie MOUGINOT<sup>1,16</sup>, John PADEN<sup>17</sup>, Frank PATTYN<sup>18</sup>, Jason ROBERTS<sup>19,20,21</sup>, Sebastian ROSIER<sup>10</sup>, Antonia RUPPEL<sup>22</sup>, Helene SEROUSSI<sup>2</sup>, Emma C. SMITH<sup>3</sup>, Daniel STEINHAGE<sup>3</sup>, Bo SUN<sup>11</sup>, Michiel R. VAN DEN BROEKE<sup>23</sup>, Tas D. VAN OMMEN<sup>19,20,21</sup>, Melchior VAN WESSEM<sup>23</sup>, Duncan YOUNG<sup>4</sup>,

<sup>1</sup>*University of California Irvine, USA,*

<sup>2</sup>*Jet Propulsion Laboratory, California Institute of Technology, USA,*

<sup>3</sup>*Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Germany,*

<sup>4</sup>*UT Austin, USA,*

<sup>5</sup>*University of Tübingen, Germany,*

<sup>6</sup>*University of Bremen, Germany,*

<sup>7</sup>*British Antarctic Survey, UK,*

<sup>8</sup>*Technical University of Denmark, Denmark,*

<sup>9</sup>*National Centre for Polar and Ocean Research, India,*

<sup>10</sup>*Northumbria University, UK,*

<sup>11</sup>*Polar Research Institute of China,*

<sup>12</sup>*Ohio State University, USA,*

<sup>13</sup>*Geological Survey of Denmark and Greenland, Denmark,*

<sup>14</sup>*Korea Polar Research Institute, South Korea,*

<sup>15</sup>*Norwegian Polar Institute, Norway,*

<sup>16</sup>*University Grenoble Alpes, France,*

<sup>17</sup>*University of Kansas, USA,*

<sup>18</sup>*Université libre de Bruxelles, Belgium,*

<sup>19</sup>*Australian Antarctic Division, Australia,*

<sup>20</sup>*Antarctic Climate & Ecosystems Cooperative Research Centre, Australia,*

<sup>21</sup>*Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania, Australia,*

<sup>22</sup>*Federal Institute for Geosciences and Natural Resources, Germany,*

<sup>23</sup>*Utrecht University, Netherlands,*

**The Antarctic ice sheet has been losing mass over the past decades through the accelerated flow of its glaciers conditioned by ocean temperature and bed topography. Glaciers retreating along retrograde slopes (i.e., bed elevation drops in the inland direction) are potentially unstable, whereas subglacial ridges slow down the glacial retreat. Despite major advances in mapping subglacial bed topography, significant sectors of Antarctica remain poorly resolved and critical spatial details are missing. Here we present a novel, high-resolution, and physically-based description of Antarctic bed topography using mass conservation. Our results reveal previously unknown basal features with major implications for glacier response**

**to climate change. For instance, glaciers flowing across the Transantarctic Mountains are protected by broad, stabilizing ridges. Conversely, in the marine basin of Wilkes Land, East Antarctica, we find retrograde slopes along Ninnis and Denman glaciers, with stabilizing slopes beneath Moscow University, Totten and Lambert glacier system, despite corrections in bed elevation of up to 1 km for the latter. This transformative description of bed topography redefines the high- and lower-risk sectors for rapid sea level rise from Antarctica; it will also significantly impact model projections of sea level rise from Antarctica in the coming centuries.**

Subglacial bed topography has been most efficiently measured using airborne radio echo sounding<sup>1</sup>. This technique provides bed elevation measurements directly beneath the aircraft path, but despite numerous campaigns, major data gaps remain between flight lines and especially across deep glaciers. As a result, there are vast sectors of Antarctica with no data<sup>2</sup>: 85% of Antarctica's surface area does not have any measurement of bed topography within a 1-km radius, and 50% of the ice sheet is more than 5 km from any measurement. The region inland of Princess Elizabeth Land, North of Dome Argus, has an area more than 90,000 km<sup>2</sup> wide with no measurement. Major data gaps exist east, west, and south of Dome Fuji and west of the Transantarctic Mountains. More importantly, we have no deep sounding near the grounding lines (i.e., at the junction with the ocean) of major glaciers such as Denman Glacier in East Antarctica or the Lambert system.

Bed elevation is difficult to sound for logistical and technical reasons. Radar sounding systems fail to probe deep subglacial troughs because steep valley walls yield side reflections that

mask the bed echoes<sup>3,4</sup> and the rough, broken-up glacier surface generates significant radar clutter. Unfortunately, these areas, while small in total area compared to the rest of the continent, are critical to characterize because they drive most of the ice discharge from Antarctica. The latest Antarctic-wide bed topography dataset<sup>5</sup>, Bedmap2, was a major improvement over previous datasets, but many sectors were undersampled, especially the glacier troughs. A major limitation of prior approaches was the sole reliance on ice thickness data combined with simple interpolation techniques, such as Kriging or thin plate splines. These approaches are highly sensitive to measurement density, resulting in ice thickness errors of several hundreds of meters to 1 km in places with few to no observation as a result of uncontrolled extrapolation. At the grounding line, it is essential to obtain a seamless transition in ice thickness and bed topography because glacier dynamics is particularly sensitive to both properties there. The level of detail required by ice sheet numerical models is typically about one ice thickness, or at least<sup>6</sup> 1 km. It is at that length scale that ridges and sinks in bed topography affect glacier dynamics. The current uncertainty and lack of small-scale detail in existing bed topography profoundly limits our ability to understand current changes in glacier flow and project ice-sheet evolution over the coming decades.

### **Mapping bed topography using mass conservation**

To overcome these difficulties, we use a mass conservation method<sup>7</sup> (MC). A chief advantage of MC is to employ a fundamental physical law to fill data gaps, i.e., the conservation of mass. The output product is fully compatible with numerical models because mass is conserved in the output product<sup>8</sup>. Second, MC employs corrections for surface mass balance and temporal changes

in ice thickness to refine the calculation of ice thickness. The resolution of the data product is no longer defined by the spacing of ice thickness data from radio-echo sounding but by the spatial resolution of the ice surface velocity, which is typically on the order of a few hundred meters for satellite-based datasets. The precision of the product, however, is affected by the spacing between ice thickness measurements, which are used to constrain the calculation, and by uncertainties and errors in the ice velocity and the surface mass balance. This methodology has been successfully applied in Greenland to transform our knowledge of bed topography and in turn our understanding of glacier dynamics, ocean circulation, ocean heat transfer, calving dynamics, and mechanisms of retreat<sup>9,10</sup>. Applying the same methodology to Antarctica presents a number of additional challenges due to the sheer size of the continent and the limited density of ice thickness data compared to Greenland.

In this study, we employ ice thickness data from 19 different research institutes, covering more than 1.5 million line kilometers over the time period 1967 to present. We use gravity-derived inversion for ice-shelf bathymetry from Operation IceBridge and other projects in a few sectors, complemented by seismic data where available. We use ice flow velocity from satellite interferometry<sup>11,12</sup>, surface mass balance from a regional atmospheric climate model<sup>13</sup>, and the surface topography from the Reference Elevation Model of Antarctica<sup>14</sup>. The grid size of the output product is 500 m. The spatial domain is divided into a number of fast flowing areas where we apply MC, and slower moving areas where we use a streamline diffusion method as an alternative to Kriging (see supplement). In total, we revise bed topography over more than 50% of the ice sheet flowing faster than 50 m/a, where MC is most accurate, and cover 71% of the ice discharge

from the continent. The results are accompanied by an error map and a source mask (see supplement), which are needed by modelers and to assist future surveys. The nominal vertical accuracy of MC is 30–60 m but local errors may exceed 100 m in poorly constrained regions. On floating ice, we rely on hydrostatic equilibrium with a firm densification model that is calibrated with all available ice shelf thickness data. This latter approach has the advantage of ensuring continuity in ice thickness across the grounding line.

The new bed compilation is named *BedMachine Antarctica* (Figure 1) because the product is regularly updated with new data. At the large scale, the shape of the bed beneath Antarctica is not fundamentally different from Bedmap2. We calculate a sea level equivalent (SLE) of  $57.9 \pm 0.9$  m for the Antarctic Ice Sheet (Table S3), which is close to the Bedmap2 estimate of 58.3 m. Most differences appear at the smaller scale, yet these local differences have a profound impact on glacier evolution, and in turn on ice sheet mass balance. As an example, we find that local bed slopes are steeper over 62% of the mapped area using MC compared to Bedmap2 (Figure S59). In addition, MC captures high-resolution details where Kriging produced smooth bed topography. The spatial details of the connectivity of individual basins with deep channels and the ocean is revised significantly, which is critical for ice sheet modeling.

### **New details along coastal margins**

In the most rapidly changing sector of Antarctica, the Amundsen Sea Embayment (ASE) (Figures 2a, 2b, and S6-S9), we find that the bed of Thwaites Glacier (65 cm SLE, 118.4 Gt/a discharge<sup>15</sup>)

has a granular texture, with no well-defined troughs, which is indicative of a hard, crystalline bedrock<sup>16</sup>. Asperities and bed ridges in the proximity of the grounding zone were for a large part missing in previous datasets, but are now found to be in excellent agreement with the observed pattern of retreat<sup>17</sup>. We do not find major bumps in bed topography upstream of the current grounding line that could stop the grounding line retreat, except for two prominent ridges about 35 and 50 km upstream (indicated as red lines in fig. 2a). Ice sheet numerical models indicate that once the glacier retreats past the second ridge, the retreat of Thwaites Glacier would become unstoppable<sup>18-20</sup>.

East of Thwaites, the bed topography of Pine Island Glacier at the grounding line (51 cm SLE, 122.6 Gt/a discharge) is 200 m deeper than in Bedmap2 because of erroneous bed picks from the top of bottom crevasses<sup>17</sup>. The older Bedmap2 product, still widely used by the modeling community, yields model simulations with limited grounding-line retreat or even grounding-line advance, both of which contradict observations<sup>21</sup>. Nearby, the bed of Kohler Glacier (Figure 2b) shows a topographically controlled ice flow, typical of selective linear erosion<sup>22,23</sup>, with a significant portion of retrograde slope. The bed of the glaciers between Pope and Smith is more continuous than in Bedmap2 and does not include a ridge across the grounding lines (Figure S9), which was an artifact of the gridding method in Bedmap2. The trough of Smith Glacier is one of the deepest and longest in West Antarctica, reaching 2,500 m below sea level, with retrograde slopes where the grounding line is retreating at record rates<sup>17</sup> of 2 to 2.5 km/a. Along the Shirase coast, West Antarctica (Figure 2c), we find a previously unknown 100-km long, 15-km wide, 1-km deep valley beneath Echelmeyer Ice Stream not resolved in prior maps (Figure S16).

Along the Transantarctic Mountains, we find deeper valleys beneath the outlet glaciers than in Bedmap2 (Figure 2d). Nimrod, Byrd and Mullock glaciers have a smaller ice discharge than Pine Island and Thwaites glaciers but have a sea level potential one order of magnitude greater due to their extensive catchments on the East Antarctic plateau. The glaciers flow along narrow submarine valleys, more than 3,000 m below sea level for Byrd Glacier. These deeply incised troughs have been challenging to resolve for radar sounding for decades, as illustrated in Figure 3b or 3c, which explains errors  $> \pm 1$  km in some places in prior mapping (Figure 3). In all cases, however, we find that the bed elevation rises rapidly above sea level within a few tens of km of the present-day grounding lines. Byrd Glacier has a prominent subglacial ridge across the Transantarctic Mountains that will provide a strong anchor point for its grounding line. David Glacier, further West (Figure 2e and 3d), is currently held by a major ridge above the cauldron area that had not been previously resolved (Figure S21). On the eastern side of David Glacier, we find a 2-km deep, 10-km wide trough that ends with an ice fall into the Drygalski Ice Tongue. The ice thickness of the ice tongue at the grounding line exceeds 2,500 m, which explains its remarkable stability and exceptional (70-km) extension out to sea, but a 10-km wide ridge, 100 m above sea level, a few kilometers upstream of the present-day grounding line will prevent the glacier from rapid retreat into the deep, Wilkes Subglacial basin (red arrow in Figure 2e). Subglacial ridges such as this one were not apparent in previous mappings but are robust features of our inversion which imply that such sectors have a low risk of collapse in decades to come (e.g., Figures 3f, S18-21).

Along George V Land (Figure 2f), the bed of Ninnis Glacier displays strong glacial lin-



eations, tens of kilometers long, likely resulting from bedrock erosion over multiple glacial cycles. The bed is flatter in this region, i.e., the flow of Ninnis is not as strongly topographically controlled as at Byrd Glacier, but is more similar to Thwaites. We find a 10-km wide valley beneath the fast-flowing portion of the glacier that extends 70 km upstream and is thus more prominent and extensive than in Bedmap2 (Figure 2f and S27). This glacier has been relatively stable over the past decades but recently lost a large part of its floating tongue<sup>15</sup>, and its bed topography suggests susceptibility to marine ice sheet instability<sup>24</sup> (MISI) that has not been previously highlighted. Conversely, further west in Wilkes Land (Figure 2g, S32 and S33), we find that Totten Glacier (3.9 m SLE, 65 Gt/a) and Moscow University Ice Shelf flow over a mostly prograde bed for 50 km upstream of the current grounding line at Totten and for 60 km at Moscow. Despite the significant thinning signal observed on Totten Glacier, evidence of a slow grounding line retreat<sup>25</sup>, presence of relatively warm water in front of the glacier<sup>26</sup> and high rates of ice shelf melt, we find that the bed topography is likely to limit any widespread MISI in that sector, until the grounding line retreats past the prograde slope areas.

Further West, Denman Glacier flows through a deep canyon more than  $\sim 3,500$  m below sea level. The full depth of the bed was not resolved even in the most recent radar field campaigns (Figure 3h) due to its deep entrenchment and the presence of a rough and broken-up ice surface<sup>3,5</sup> (Figure S35). BedMachine reveals that the bed beneath this ice stream is the deepest continental point on Earth. Close to the grounding line, the bed slope is gentle and slightly retrograde, which could lead to instability if the grounding line were to retreat inland, making this sector very vulnerable in East Antarctica, with a potential 1.5 m sea level rise.

On Mellor Glacier, upstream of Amery Ice Shelf, we find a 3-km deep bed depression (Figure 2i and 3j) that is inconsistent with prior radar data that indicated a bed only 1000 m below sea level, which also yielded ice fluxes that were much too low to balance upstream accumulation. We conclude that the radar data have been systematically misinterpreted in that region, probably due to side reflections (Figure 3k-1, S37). MC requires ice to be more than 1-km thicker at that location, which is quite plausible because this is a zone of convergence of three glaciers (Lambert, Mellor and Fisher) constrained by mountain ranges. The valleys are mostly prograde and the basin upstream rises rapidly above sea level except along the East Lambert Rift, suggesting that this sector has low potential for MISI in the near future.

Bed topography further west, stretching from Enderby to Queen Maud land, is locally retrograde only for a few tens of kilometers (Fig. S60) and therefore not as vulnerable to MISI as other regions. In the Baudouin sector, West Ragnhild ice stream flows on a prograde submarine valley that extends 80 km further inland<sup>27</sup> than in Bedmap2, but eventually rises above sea level (Figure 2j). Conversely, further west along Coats Land, several major ice streams feeding the Ronne-Filchner ice shelves stand on strongly retrograde bed slopes from 100 km to 600 km further upstream than in Bedmap2: Slessor (2.9 m SLE), Recovery (6.2 SLE), Support Force and Academy (2.5 SLE). Recovery (Figure 2k) is 800 m deeper than previously thought (Figure 3o and S46). This region is a major point of vulnerability in East Antarctica.

At the southeastern tip of the Antarctic Peninsula, we report a well-defined valley that coincides with Evans Ice Stream and four tributaries feeding the main ice stream (Figure 3l). This

sector is an example of selective linear erosion characterized by more rapid basal incision by fast-flowing, warm-based ice, relative to the surrounding slower, cold-based ice. Some tributaries flow in troughs more than 2 km below sea level that drain ice from a predominantly submarine basin.

Among the limitations of our compilation, we note a lack of ocean bathymetry on the continental shelf and beneath ice shelves, which remains a problem over vast portions of the coast of Antarctica. Multibeam echo sounding data, gravity data, seismic data and sea floor depth from robotic devices will be essential to improve bathymetry mapping in this part of Antarctica, which is critical for ice/ocean interactions and for ice sheet mass balance<sup>28</sup>. To improve the mapping of fast flowing regions, we recommend flight tracks perpendicular to the flow direction to maximize constraints on ice flux, especially upstream of Academy and Support force glaciers, along Stancomb-Wills, Gould Coast near the Ross ice shelf, and Wilhem II Coast between Denman and Lambert glaciers.

### **Implications for ice sheet vulnerability**

The new bed topography highlights regions of higher vulnerability in West Antarctica and regions of low risk in the Ross Sea sector, along the Transantarctic Mountains. Glaciers spanning from George V Land to Dibble Glacier in Terre Adélie and Wilkes Land are in contrast located at the mouth of deep submarine basins with retrograde slopes, hence risk zones for MISI. In Wilkes Land, Totten Glacier and Moscow University Ice Shelf would have to retreat about 50 km inland before reaching a zone of retrograde bed, but Denman Glacier stands at the edge of a deep trough

that makes it vulnerable. Further west, the glaciers in Enderby and Queen Maud Land flow over prograde bed slope, except along a narrow coastal margin, and the drainage basins are mostly above sea level, hence more protected from MISI. Conversely, the glaciers feeding the eastern side of the Filchner Ice Shelf have retrograde slopes over vast portions of their basin, hence prone to MISI. It will be essential to refine these results with more precise observations in the future to better inform ice sheet numerical models, but the new product has already brought major changes that call into question prior modeling using older maps. The revised bed topography will enable more robust ice sheet numerical modeling and improved projections of the contribution of Antarctica to sea level rise.

## **Method Summary**

The method of mass conservation, MC<sup>7,29</sup>, yields ice thickness and bed topography compatible with ice sheet numerical models, resolves uncertainties of prior interpretation of radar echoes, and ensures that grounding lines fluxes are compatible with snowfall accumulation and thinning rates in the interior without assuming steady-state. We use radar-derived thickness data from multiple sources, with a vertical precision of  $\sim 30$  m, ice velocity measurements derived from satellite radar data posted at 150 m, with errors of 10 m/a in speed and  $1.5^\circ$  in flow direction<sup>11</sup>, the REMA DEM<sup>14</sup>, gravity-derived bathymetry<sup>28,30,31</sup>, seismic bathymetry<sup>32</sup>, and IBCSO data<sup>33</sup>; and surface mass balance<sup>13</sup> (SMB) averaged for the years 1961-1990 with a 7% accuracy. The algorithm neglects ice motion by internal shear, which is a good approximation<sup>7,29</sup> for fast-flowing glaciers ( $> 50$  m/a). The optimization procedure is not applied in slow-moving sectors, where we use a

streamline diffusion. For floating ice shelves, we rely on hydrostatic equilibrium with a calibrated firn depth correction so the inferred ice thickness is consistent with available ice thickness data. More technical details and error analyses are provided in the supplementary material.

**Correspondence** Correspondence and requests for materials should be addressed to M. Morlighem (email: mathieu.morlighem@uci.edu).

**Acknowledgements** This work was performed at the University of California Irvine under a contract with the National Aeronautics and Space Administration, Cryospheric Sciences Program (#NNX17AI02G, #NNX15AD55G and #NNX14AN03G), MEaSURES-3 Program (#80NSSC18M0083), the NSF-NERC International Thwaites Glacier Collaboration (Award #1739031), and the Australian Research Council's Special Research Initiative for Antarctic Gateway Partnership (Project ID SR140300001). We acknowledge the use of data and/or data products from CReSIS generated with support from the University of Kansas, NSF grant ANT-0424589, and NASA Operation IceBridge grant NNX16AH54G, and data products collected by the ICECAP collaboration under NSF grants ANT-1043761, ANT-1543452, ANT-0733025, ANT-1443690 and ANT-1143843, NASA grants #NNG10HPO6C and #NNX11AD33G, and AAD projects 3013, 4077 and 4346, the Australian Research Council's Special Research Initiative for the Antarctic Gateway Partnership (project ID SR140300001) and the Australian Government's Cooperative Research Centre program through the Antarctic Climate and Ecosystems Cooperative Research Centre, the National Natural Science Foundation of China grant 41876227, with support by the G. Unger Vetlesen Foundation. R. D. was partially supported by the DFG Emmy Noether Grant DR 822/3-1, and W. L. was supported by the Korean Ministry of Oceans and Fisheries (KIMST20190361; PM19020) and KOPRI(PE19110).

**Author Contributions** M.M. developed the algorithm and lead the calculations. H.S. assisted in imple-

menting the algorithm. M.M. and E.R. wrote the first draft of the manuscript. All authors contributed data and to the writing of the paper.

**Data availability** BedMachine Antarctica is publicly available at the NSIDC, Boulder CO, as a MEASURES-3 product (URL coming soon).

**Code availability** The algorithms used to generate the bed topography are included in the open-source Ice Sheet System Model (<https://issm.jpl.nasa.gov>).

1. Evans, S. & Robin, G. d. Q. Glacier depth-sounding from air. *Nature* **210**, 883–885 (1966).
2. Pritchard, H. D. Bedgap: where next for Antarctic subglacial mapping? *Antarct. Sci.* **26**, 742–757 (2014).
3. Holt, J. W., Peters, M. E., Kempf, S. D., Morse, D. L. & Blankenship, D. D. Echo source discrimination in single-pass airborne radar sounding data from the Dry Valleys, Antarctica: Implications for orbital sounding of Mars. *J. Geophys. Res.* **111**, 1–13 (2006).
4. Jezek, K., Wu, X., Paden, J. & Leuschen, C. Radar mapping of Isunnguata Sermia, Greenland. *J. Glaciol.* **59**, 1135–1146 (2013).
5. Fretwell, P. *et al.* Bedmap2: improved ice bed, surface and thickness datasets for Antarctica. *Cryosphere* **7**, 375–393 (2013).
6. Durand, G., Gagliardini, O., Favier, L., Zwinger, T. & le Meur, E. Impact of bedrock description on modeling ice sheet dynamics. *Geophys. Res. Lett.* **38**, L20501, 1–5 (2011).


7. Morlighem, M. *et al.* A mass conservation approach for mapping glacier ice thickness. *Geophys. Res. Lett.* **38**, 1–6 (2011).
8. Seroussi, H. *et al.* Ice flux divergence anomalies on 79north Glacier, Greenland. *Geophys. Res. Lett.* **38**, 1–5 (2011).
9. Morlighem, M., Rignot, E., Mouginot, J., Seroussi, H. & Larour, E. Deeply incised submarine glacial valleys beneath the Greenland Ice Sheet. *Nat. Geosci.* **7**, 418–422 (2014).
10. Morlighem, M. *et al.* BedMachine v3: Complete bed topography and ocean bathymetry mapping of Greenland from multi-beam echo sounding combined with mass conservation. *Geophys. Res. Lett.* **44**, 11,051–11,061 (2017). 2017GL074954.
11. Rignot, E., Velicogna, I., van den Broeke, M. R., Monaghan, A. & Lenaerts, J. Acceleration of the contribution of the Greenland and Antarctic ice sheets to sea level rise. *Geophys. Res. Lett.* **38**, 1–5 (2011).
12. Mouginot, J., Rignot, E., Scheuchl, B. & Millan, R. Comprehensive Annual Ice Sheet Velocity Mapping Using Landsat-8, Sentinel-1, and RADARSAT-2 Data. *Remote Sens.* **9** (2017).
13. van Wessem, J. M. *et al.* Modelling the climate and surface mass balance of polar ice sheets using RACMO2: Part 2: Antarctica (1979-2016). *Cryosphere* **12**, 1479–1498 (2018).
14. Howat, I. M., Porter, C., Smith, B. E., Noh, M.-J. & Morin, P. The Reference Elevation Model of Antarctica. *Cryosphere* **13**, 665–674 (2019).


15. Rignot, E. *et al.* Four decades of Antarctic Ice Sheet mass balance from 1979–2017. *Proc. Natl. Acad. Sci.* **116**, 1095–1103 (2019).  
<https://www.pnas.org/content/early/2019/01/08/1812883116.full.pdf>.
16. Bingham, R. G. *et al.* Diverse landscapes beneath Pine Island Glacier influence ice flow. *Nat. Commun.* **8**, 1618 (2017).
17. Rignot, E., Mouginot, J., Morlighem, M., Seroussi, H. & Scheuchl, B. Widespread, rapid grounding line retreat of Pine Island, Thwaites, Smith and Kohler glaciers, West Antarctica from 1992 to 2011. *Geophys. Res. Lett.* **41**, 3502–3509 (2014).
18. Joughin, I., Smith, B. E. & Medley, B. Marine Ice Sheet Collapse Potentially Underway for the Thwaites Glacier Basin, West Antarctica. *Science* **344**, 735–738 (2014).
19. Seroussi, H. *et al.* Continued retreat of Thwaites Glacier, West Antarctica, controlled by bed topography and ocean circulation. *Geophys. Res. Lett.* **44**, 6191–6199 (2017). 2017GL072910.
20. Yu, H., Rignot, E., Seroussi, H. & Morlighem, M. Retreat of Thwaites Glacier, West Antarctica, over the next 100 years using various ice flow models, ice shelf melt scenarios and basal friction laws. *Cryosphere* **12**, 3861–3876 (2018).
21. Nias, I. J., Cornford, S. L. & Payne, A. J. New mass-conserving bedrock topography for pine island glacier impacts simulated decadal rates of mass loss. *Geophys. Res. Lett.* **45**, 3173–3181 (2018).  
<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2017GL076493>.




22. Sugden, D. E. & John, B. S. *Glaciers and Landscape: A Geomorphological Approach* (Edward Arnold, 1976).
23. Jamieson, S. S. *et al.* The glacial geomorphology of the antarctic ice sheet bed. *Antarct. Sci.* **26**, 724–741 (2014).
24. Weertman, J. Stability of the junction of an ice sheet and an ice shelf. *J. Glaciol.* **13(67)**, 3–11 (1974).
25. Li, X., Rignot, E., Morlighem, M., Mouginot, J. & Scheuchl, B. Grounding line retreat of Totten Glacier, East Antarctica, 1996 to 2013. *Geophys. Res. Lett.* **42**, 8049–8056 (2015).
26. Rintoul, S. R. *et al.* Ocean heat drives rapid basal melt of the Totten Ice Shelf. *Sci. Adv.* **2** (2016).  
<https://advances.sciencemag.org/content/2/12/e1601610.full.pdf>.
27. Eagles, G. *et al.* Erosion at extended continental margins: Insights from new aerogeophysical data in eastern Dronning Maud Land. *Gondwana Res.* **63**, 105 – 116 (2018).
28. Millan, R., Rignot, E., Bernier, V., Morlighem, M. & Dutrieux, P. Bathymetry of the Amundsen Sea Embayment sector of West Antarctica from Operation IceBridge gravity and other data. *Geophys. Res. Lett.* **44**, 1360–1368 (2017).
29. Morlighem, M. *et al.* High-resolution bed topography mapping of Russell Glacier, Greenland, inferred from Operation IceBridge data. *J. Glaciol.* **59**, 1015–1023 (2013).

30. Greenbaum, J. S. *et al.* Ocean access to a cavity beneath Totten Glacier in East Antarctica. *Nat. Geosci.* **8**, 294–298 (2015).
31. Tinto, K. J. *et al.* Ross Ice Shelf response to climate driven by the tectonic imprint on seafloor bathymetry. *Nat. Geosci.* **12**, 441–449 (2019).
32. Rosier, S. H. R. *et al.* A New Bathymetry for the Southeastern Filchner-Ronne Ice Shelf: Implications for Modern Oceanographic Processes and Glacial History. *J. Geophys. Res. - Oceans* **123**, 4610–4623 (2018).  
<https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2018JC013982>.
33. Arndt, J. E. *et al.* The International Bathymetric Chart of the Southern Ocean (IBCSO) Version 1.0: A new bathymetric compilation covering circum-Antarctic waters. *Geophys. Res. Lett.* **40**, 3111–3117 (2013).







**Figure 1** Bed elevation of the Antarctic ice sheet color coded between -3,000 m and 1,500 m above sea level. White line delineates the basins from the ice sheet mass balance inter-comparison exercise (IMBIE).

**Figure 2** Detailed bed topography of Antarctic outlet glaciers. Bed elevation of a) Thwaites, b) Kohler, Smith and Pope glaciers, c) Shirase coast, d) Byrd and Mullock glaciers, e) David, f) Ninnis and Mertz glaciers, g) Moscow University Ice Shelf and Totten glacier, h) Denman, i) Lambert glaciers, j) Roi Baudouin Ice Shelf, k) Recovery and l) Evans Ice Streams, color coded between -3,000 m and 1,500 m above sea level. The black lines show the ice extent and white line the grounding lines.

**Figure 3** Comparison with previous datasets and radar data. Bed elevation of Byrd (a), David (d), Denman (g), Mellor (j) and Recovery (m) glaciers color coded between -3000 m and 1500 m above sea level, with radar profiles shown as white lines where bed reflections were detected. The yellow and red lines (e.g., A1-A2 and A1'-A2' in the first row left and right, respectively, etc.) show the locations where the profiles on the middle and right column panels are extracted. The second column shows the profiles along the red line, which corresponds to a flight line, and the third column shows profiles along the yellow line (along flow). The solid black line shows the surface elevation along the transect, the dashed black line is sea level, the solid blue line is the bed elevation from BedMachine (and associated uncertainty in light blue), and the green line is the bed topography from

Bedmap2 (and associated uncertainty in light green). The black dots in the panels of the second column show the radar derived bed elevation with error bars.