

Fully Homomorphic Encryption with Zero Norm Cipher Text

Masahiro Yagisawa†

†Resident in Yokohama-shi

Sakae-ku, Yokohama-shi, Japan

tfkt8398yagi@outlook.jp

Abstract. Gentry’s bootstrapping technique is the most famous method of obtaining fully homomorphic encryption. In previous work I proposed a fully homomorphic encryption without bootstrapping which has the weak point in the plaintext [1],[15]. I also proposed fully homomorphic encryptions with composite number modulus which avoid the weak point by adopting the plaintext including the random numbers in it [14],[16]. In this paper I propose another fully homomorphic encryption with zero norm cipher text where zero norm medium text is generated and enciphered by using composite number modulus. In the proposed scheme it is proved that if there exists the PPT algorithm that generates the cipher text of the plaintext $-p$ from the cipher text of any plaintext p , there exists the PPT algorithm that factors the given composite number modulus. That is, we include the random parameters in the plaintext so that if the random parameter and the plaintext are separated, then the composite number to be the modulus is factored. Since the scheme is based on computational difficulty to solve the multivariate algebraic equations of high degree while the almost all multivariate cryptosystems [2],[3],[4],[5] proposed until now are based on the quadratic equations avoiding the explosion of the coefficients. Because proposed fully homomorphic encryption scheme is based on multivariate algebraic equations with high degree or too many variables, it is against the Gröbner basis [6] attack, the differential attack, rank attack and so on.

keywords: fully homomorphic encryption, zero norm cipher text, Gröbner basis, octonion, factoring

§1. Introduction

A cryptosystem which supports both addition and multiplication (thereby preserving the ring structure of the plaintexts) is known as fully homomorphic encryption (FHE) and is very powerful. Using such a scheme, any circuit can be homomorphically evaluated, effectively allowing the construction of programs which may be run on encryptions of their inputs to produce an encryption of their output. Since such a

program never decrypts its input, it can be run by an untrusted party without revealing its inputs and internal state. The existence of an efficient and fully homomorphic cryptosystem would have great practical implications in the outsourcing of private computations, for instance, in the context of cloud computing.

With homomorphic encryption, a company could encrypt its entire database of e-mails and upload it to a cloud. Then it could use the cloud-stored data as desired—for example, to calculate the stochastic value of stored data. The results would be downloaded and decrypted without ever exposing the details of a single e-mail.

In 2009 Gentry has created a homomorphic encryption scheme that makes it possible to encrypt the data in such a way that performing a mathematical operation on the encrypted information and then decrypting the result produces the same answer as performing an analogous operation on the unencrypted data[7],[8].

But in Gentry's scheme a task like finding a piece of text in an e-mail requires chaining together thousands of basic operations. His solution was to use a second layer of encryption, essentially to protect intermediate results when the system broke down and needed to be reset.

Some fully homomorphic encryption schemes were proposed until now [9], [10], [11],[12].

In this paper I propose a fully homomorphic encryption scheme on non-associative octonion ring over finite ring with composite number modulus which is based on computational difficulty to solve the multivariate algebraic equations of high degree while the almost all multivariate cryptosystems [2],[3],[4],[5] proposed until now are based on the quadratic equations avoiding the explosion of the coefficients. Our scheme is against the Gröbner basis [6] attack, the differential attack, rank attack and so on.

In this scheme the norm of every medium text is given as the product of two factors made of the parameters in the medium text where the each factor includes the different prime factor of the composite modulus. Then every medium text has zero norm so that every cipher text also has zero norm. In this scheme zero norm medium text is generated and enciphered by using composite number modulus. It is proved in section 4.4 that if there exists the PPT algorithm that generates the ciphertext of the plaintext $-p$ from the ciphertext of any plaintext p , there exists the PPT algorithm that factors the given composite number modulus. In other words we include the random parameter in the plaintext so that if the random parameters and the plaintext are separated, then the composite number to be the modulus is factored.

§2. Preliminaries for octonion operation

In this section we describe the operations on octonion ring and properties of octonion ring. The readers who understand the property of octonion may skip this section.

§2.1 Multiplication and addition on the octonion ring O

Let s and t be 1000-digit primes where s and t are secret. Let $q=st$ be a fixed modulus to be 2000-digit composite number where

$$s < t. \quad (1)$$

Let O be the octonion [13] ring over a ring $\mathbf{Z}/q\mathbf{Z}$.

$$O = \{(a_0, a_1, \dots, a_7) \mid a_j \in \mathbf{Z}/q\mathbf{Z} (j=0,1,\dots,7)\} \quad (2)$$

We define the multiplication and addition of $A, B \in O$ as follows.

$$A = (a_0, a_1, \dots, a_7), a_j \in \mathbf{Z}/q\mathbf{Z} (j=0,1,\dots,7), \quad (3a)$$

$$B = (b_0, b_1, \dots, b_7), b_j \in \mathbf{Z}/q\mathbf{Z} (j=0,1,\dots,7). \quad (3b)$$

$AB \bmod q$

$$\begin{aligned} &= (a_0b_0 - a_1b_1 - a_2b_2 - a_3b_3 - a_4b_4 - a_5b_5 - a_6b_6 - a_7b_7 \bmod q, \\ &\quad a_0b_1 + a_1b_0 + a_2b_4 + a_3b_7 - a_4b_2 + a_5b_6 - a_6b_5 - a_7b_3 \bmod q, \\ &\quad a_0b_2 - a_1b_4 + a_2b_0 + a_3b_5 + a_4b_1 - a_5b_3 + a_6b_7 - a_7b_6 \bmod q, \\ &\quad a_0b_3 - a_1b_7 - a_2b_5 + a_3b_0 + a_4b_6 + a_5b_2 - a_6b_4 + a_7b_1 \bmod q, \\ &\quad a_0b_4 + a_1b_2 - a_2b_1 - a_3b_6 + a_4b_0 + a_5b_7 + a_6b_3 - a_7b_5 \bmod q, \\ &\quad a_0b_5 - a_1b_6 + a_2b_3 - a_3b_2 - a_4b_7 + a_5b_0 + a_6b_1 + a_7b_4 \bmod q, \\ &\quad a_0b_6 + a_1b_5 - a_2b_7 + a_3b_4 - a_4b_3 - a_5b_1 + a_6b_0 + a_7b_2 \bmod q, \\ &\quad a_0b_7 + a_1b_3 + a_2b_6 - a_3b_1 + a_4b_5 - a_5b_4 - a_6b_2 + a_7b_0 \bmod q) \end{aligned} \quad (4)$$

$A+B \bmod q$

$$\begin{aligned} &= (a_0 + b_0 \bmod q, a_1 + b_1 \bmod q, a_2 + b_2 \bmod q, a_3 + b_3 \bmod q, \\ &\quad a_4 + b_4 \bmod q, a_5 + b_5 \bmod q, a_6 + b_6 \bmod q, a_7 + b_7 \bmod q). \end{aligned} \quad (5)$$

Let $|A|$ be the norm of A such that

$$|A|^2 = a_0^2 + a_1^2 + \dots + a_7^2 \bmod q. \quad (6)$$

If $|A|^2$ has the inverse mod q , we can have A^{-1} , the inverse of A by using the algorithm **Octinv**(A) such that

$$A^{-1} = (a_0/|A|^2 \bmod q, -a_1/|A|^2 \bmod q, \dots, -a_7/|A|^2 \bmod q) \leftarrow \mathbf{Octinv}(A). \quad (7)$$

Here details of the algorithm $\mathbf{Octinv}(A)$ are omitted and can be looked up in the **Appendix A**.

§2.2. Property of multiplication over octonion ring O

A, B, C etc. $\in O$ satisfy the following formulae in general where A, B and C have the inverse A^{-1}, B^{-1} and $C^{-1} \bmod q$.

1) Non-commutative

$$AB \neq BA \quad \bmod q.$$

2) Non-associative

$$A(BC) \neq (AB)C \quad \bmod q.$$

3) Alternative

$$(AA)B = A(AB) \quad \bmod q, \quad (8)$$

$$A(BB) = (AB)B \quad \bmod q, \quad (9)$$

$$(AB)A = A(BA) \quad \bmod q. \quad (10)$$

4) Moufang's formulae [13],

$$C(A(CB)) = ((CA)C)B \quad \bmod q, \quad (11)$$

$$A(C(BC)) = ((AC)B)C \quad \bmod q, \quad (12)$$

$$(CA)(BC) = (C(AB))C \quad \bmod q, \quad (13)$$

$$(CA)(BC) = C((AB)C) \quad \bmod q. \quad (14)$$

5) For positive integers n, m , we have

$$(AB)B^n = ((AB)B^{n-1})B = A(B(B^{n-1}B)) = AB^{n+1} \quad \bmod q, \quad (15)$$

$$(AB^n)B = ((AB)B^{n-1})B = A(B(B^{n-1}B)) = AB^{n+1} \quad \bmod q, \quad (16)$$

$$B^n(BA) = B(B^{n-1}(BA)) = ((BB^{n-1})B)A = B^{n+1}A \quad \bmod q, \quad (17)$$

$$B(B^n A) = B(B^{n-1}(BA)) = ((BB^{n-1})B)A = B^{n+1}A \quad \bmod q. \quad (18)$$

From (9) and (16), we have

$$[(AB^n)B]B = [AB^{n+1}]B \quad \bmod q,$$

$$(AB^n)(BB) = [(AB^n)B]B = [AB^{n+1}]B = AB^{n+2} \quad \bmod q,$$

$$(AB^n)B^2 = AB^{n+2} \pmod{q},$$

...

$$(AB^n)B^m = AB^{n+m} \pmod{q}.$$

In the same way we have

$$B^m(B^n A) = B^{n+m}A \pmod{q}. \quad (19)$$

6) Lemma 1

$$A(B((AB)^n)) = (AB)^{n+1} \pmod{q},$$

$$(((AB)^n)A)B = (AB)^{n+1} \pmod{q}.$$

where n is a positive integer and B has the inverse B^{-1} .

(Proof:)

From (12) we have

$$B(A(B((AB)^n))) = ((BA)B)(AB)^n = (B(AB))(AB)^n = B(AB)^{n+1} \pmod{q}.$$

Then

$$B^{-1}(B(A(B(AB)^n))) = B^{-1}(B(AB)^{n+1}) \pmod{q},$$

$$A(B(AB)^n) = (AB)^{n+1} \pmod{q}.$$

In the same way we have

$$(((AB)^n)A)B = (AB)^{n+1} \pmod{q}. \quad \text{q.e.d.}$$

7) Lemma 2

$$A^{-1}(AB) = B \pmod{q},$$

$$(BA)A^{-1} = B \pmod{q}.$$

(Proof:)

Here proof is omitted and can be looked up in the **Appendix B**.

8) Lemma 3

$$A(BA^{-1}) = (AB)A^{-1} \pmod{q}.$$

(Proof:)

From (14) we substitute A^{-1} to C , we have

$$(A^{-1}A)(BA^{-1})=A^{-1}((AB)A^{-1}) \bmod q,$$

$$(BA^{-1})=A^{-1}((AB)A^{-1}) \bmod q.$$

We multiply A from left side,

$$A(BA^{-1})=A(A^{-1}((AB)A^{-1}))=(AB)A^{-1} \bmod q. \quad \text{q.e.d.}$$

We can express $A(BA^{-1})$, $(AB)A^{-1}$ such that

$$ABA^{-1}.$$

9) From (10) and Lemma 2 we have

$$A^{-1}((A(BA^{-1}))A)=A^{-1}(A((BA^{-1})A))=(BA^{-1})A=B \bmod q,$$

$$(A^{-1}((AB)A^{-1}))A=((A^{-1}((AB)A^{-1}))A)=A^{-1}(AB)=B \bmod q.$$

10) **Lemma 4**

$$(BA^{-1})(AB)=B^2 \bmod q.$$

(Proof:)

From (14),

$$(BA^{-1})(AB)=B((A^{-1}A)B)=B^2 \bmod q. \quad \text{q.e.d.}$$

11a) **Lemma 5a**

$$(ABA^{-1})(ABA^{-1})=AB^2A^{-1} \bmod q.$$

(Proof:)

From (14),

$$\begin{aligned} & (ABA^{-1})(ABA^{-1}) \bmod q \\ &= [A^{-1}(A^2(BA^{-1}))][(AB)A^{-1}] = A^{-1} \{ [(A^2(BA^{-1}))(AB)]A^{-1} \} \bmod q \\ &= A^{-1} \{ [(A(A(BA^{-1}))) (AB)]A^{-1} \} \bmod q \\ &= A^{-1} \{ [(A((AB)A^{-1})) (AB)]A^{-1} \} \bmod q \\ &= A^{-1} \{ [(A(AB))A^{-1}] (AB)]A^{-1} \} \bmod q. \end{aligned}$$

We apply (12) to inside of [.],

$$= A^{-1} \{ [(A((AB)(A^{-1}(AB))))]A^{-1} \} \bmod q$$

$$\begin{aligned}
&= A^{-1} \{ [(A((AB)B))] A^{-1} \} \pmod q \\
&= A^{-1} \{ [A(A(BB))] A^{-1} \} \pmod q \\
&= \{ A^{-1} [A(A(BB))] \} A^{-1} \pmod q \\
&= (A(BB)) A^{-1} \pmod q \\
&= AB^2 A^{-1} \pmod q. \qquad \qquad \qquad \text{q.e.d.}
\end{aligned}$$

11b) Lemma 5b

$$\begin{aligned}
&[A_1(\dots(A_r B A_r^{-1})\dots)A_1^{-1}] [A_1(\dots(A_r B A_r^{-1})\dots)A_1^{-1}] \\
&= A_1(\dots(A_r B^2 A_r^{-1})\dots)A_1^{-1} \pmod q.
\end{aligned}$$

where

$$A_i \in O \text{ has the inverse } A_i^{-1} \pmod q \text{ (} i=1, \dots, r \text{)}.$$

(Proof:)

As we use Lemma 5a repeatedly we have

$$\begin{aligned}
&\{ A_1([A_2(\dots(A_r B A_r^{-1})\dots)A_2^{-1}]) A_1^{-1} \} \{ A_1([A_2(\dots(A_r B A_r^{-1})\dots)A_2^{-1}]) A_1^{-1} \} \pmod q \\
&= A_1([A_2(\dots(A_r B A_r^{-1})\dots)A_2^{-1}] [A_2(\dots(A_r B A_r^{-1})\dots)A_2^{-1}]) A_1^{-1} \pmod q \\
&= A_1(A_2([A_3(\dots(A_r B A_r^{-1})\dots)A_3^{-1}][A_3(\dots(A_r B A_r^{-1})\dots)A_3^{-1}] A_2^{-1}) A_1^{-1} \pmod q \\
&\qquad \qquad \qquad \dots \qquad \qquad \dots \\
&= A_1(A_2(\dots([A_r B A_r^{-1}] [A_r B A_r^{-1}])\dots)A_2^{-1}) A_1^{-1} \pmod q \\
&= A_1(A_2(\dots(A_r B^2 A_r^{-1})\dots)A_2^{-1}) A_1^{-1} \pmod q \\
&\qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \text{q.e.d.}
\end{aligned}$$

11c) Lemma 5c

$$\begin{aligned}
&A_1^{-1} (A_1 B A_1^{-1}) A_1 \\
&= B \pmod q.
\end{aligned}$$

where

$$A_1 \in O \text{ has the inverse } A_1^{-1} \pmod q.$$

(Proof:)

$$A_1^{-1} (A_1 B A_1^{-1}) A_1 = A_1^{-1} [((A_1 B) A_1^{-1}) A_1] \pmod q,$$

From Lemma 2 we have

$$= A_1^{-1} (A_1 B) = B \pmod{q}. \quad \text{q.e.d.}$$

11d) **Lemma 5d**

$$\begin{aligned} & A_r^{-1} (\dots (A_1^{-1} [A_1 (\dots (A_r B A_r^{-1}) \dots) A_1^{-1}] A_1) \dots) A_r \\ & = B \pmod{q}. \end{aligned}$$

where

$$A_i \in O \text{ has the inverse } A_i^{-1} \pmod{q} \ (i=1, \dots, r).$$

(*Proof:*)

As we use Lemma 5c repeatedly we have

$$\begin{aligned} & A_r^{-1} (\dots (A_1^{-1} [A_1 (\dots (A_r B A_r^{-1}) \dots) A_1^{-1}] A_1) \dots) A_r \\ = & A_r^{-1} (\dots (A_2^{-1} [A_2 (\dots (A_r B A_r^{-1}) \dots) A_2^{-1}] A_2) \dots) A_r \pmod{q} \\ & \dots \quad \dots \\ & = A_r^{-1} [A_r B A_r^{-1}] A_r \pmod{q} \\ & = B \pmod{q} \quad \text{q.e.d.} \end{aligned}$$

12) **Lemma 6**

$$(A B^m A^{-1})(A B^n A^{-1}) = A B^{m+n} A^{-1} \pmod{q}.$$

(*Proof:*)

From (13),

$$\begin{aligned} & [A^{-1} (A^2 (B^m A^{-1}))][A (B^n) A^{-1}] = \{A^{-1} [(A^2 (B^m A^{-1})) (A B^n)]\} A^{-1} \pmod{q} \\ & = A^{-1} \{ [(A (A (B^m A^{-1}))) (A B^n)] A^{-1} \} \pmod{q} \\ & = A^{-1} \{ [(A ((A B^m) A^{-1})) (A B^n)] A^{-1} \} \pmod{q} \\ & = A^{-1} \{ [((A (A B^m)) A^{-1}) (A B^n)] A^{-1} \} \pmod{q} \\ & = A^{-1} \{ [((A^2 B^m) A^{-1}) (A B^n)] A^{-1} \} \pmod{q}. \end{aligned}$$

We apply (12) to inside of { . },

$$= A^{-1} \{ (A^2 B^m) [A^{-1} ((A B^n) A^{-1})] \} \pmod{q}$$

$$\begin{aligned}
&= A^{-1} \{ (A^2 B^m)[A^{-1}(A(B^n A^{-1}))]\} \pmod q \\
&= A^{-1} \{ (A^2 B^m)(B^n A^{-1})\} \pmod q \\
&= A^{-1} \{ (A^{-1}(A^3 B^m))(B^n A^{-1})\} \pmod q.
\end{aligned}$$

We apply (12) to inside of { . },

$$\begin{aligned}
&= A^{-1} \{ [(A^{-1}(A^3 B^m))B^n]A^{-1}\} \pmod q \\
&= A^{-1} \{ ((A^2 B^m)B^n)A^{-1}\} \pmod q \\
&= A^{-1} \{ (A^2 B^{m+n})A^{-1}\} \pmod q \\
&= \{ A^{-1} (A^2 B^{m+n}) \} A^{-1} \pmod q \\
&= (A B^{m+n}) A^{-1} \pmod q \\
&= A B^{m+n} A^{-1} \pmod q. \qquad \qquad \qquad \text{q.e.d}
\end{aligned}$$

13) $A \in O$ satisfies the following theorem.

[Theorem 1]

$$A^2 = w\mathbf{1} + vA \pmod q,$$

where

$$\exists w, v \in \mathbf{Z}/q\mathbf{Z},$$

$$\mathbf{1} = (1, 0, 0, 0, 0, 0, 0) \in O,$$

$$A = (a_0, a_1, \dots, a_7) \in O.$$

(Proof:)

$$\begin{aligned}
&A^2 \pmod q \\
&= (a_0 a_0 - a_1 a_1 - a_2 a_2 - a_3 a_3 - a_4 a_4 - a_5 a_5 - a_6 a_6 - a_7 a_7) \pmod q, \\
&\quad a_0 a_1 + a_1 a_0 + a_2 a_4 + a_3 a_7 - a_4 a_2 + a_5 a_6 - a_6 a_5 - a_7 a_3 \pmod q, \\
&\quad a_0 a_2 - a_1 a_4 + a_2 a_0 + a_3 a_5 + a_4 a_1 - a_5 a_3 + a_6 a_7 - a_7 a_6 \pmod q, \\
&\quad a_0 a_3 - a_1 a_7 - a_2 a_5 + a_3 a_0 + a_4 a_6 + a_5 a_2 - a_6 a_4 + a_7 a_1 \pmod q, \\
&\quad a_0 a_4 + a_1 a_2 - a_2 a_1 - a_3 a_6 + a_4 a_0 + a_5 a_7 + a_6 a_3 - a_7 a_5 \pmod q, \\
&\quad a_0 a_5 - a_1 a_6 + a_2 a_3 - a_3 a_2 - a_4 a_7 + a_5 a_0 + a_6 a_1 + a_7 a_4 \pmod q,
\end{aligned}$$

$$\begin{aligned}
& a_0a_6+a_1a_5-a_2a_7+a_3a_4-a_4a_3-a_5a_1+a_6a_0+a_7a_2 \pmod q, \\
& a_0a_7+a_1a_3+a_2a_6-a_3a_1+a_4a_5-a_5a_4-a_6a_2+a_7a_0 \pmod q) \\
& =(2a_0^2-L \pmod q, 2a_0a_1 \pmod q, 2a_0a_2 \pmod q, 2a_0a_3 \pmod q, \\
& 2a_0a_4 \pmod q, 2a_0a_5 \pmod q, 2a_0a_6 \pmod q, 2a_0a_7 \pmod q)
\end{aligned}$$

where

$$L = a_0^2 + a_1^2 + a_2^2 + a_3^2 + a_4^2 + a_5^2 + a_6^2 + a_7^2 \pmod q.$$

Now we try to obtain $u, v \in Fq$ that satisfy $A^2 = w\mathbf{1} + vA \pmod q$.

$$\begin{aligned}
w\mathbf{1} + vA &= w(1, 0, 0, 0, 0, 0, 0, 0) + v(a_0, a_1, \dots, a_7) \pmod q, \\
A^2 &= (2a_0^2 - L \pmod q, 2a_0a_1 \pmod q, 2a_0a_2 \pmod q, 2a_0a_3 \pmod q, \\
& 2a_0a_4 \pmod q, 2a_0a_5 \pmod q, 2a_0a_6 \pmod q, 2a_0a_7 \pmod q).
\end{aligned}$$

Then we have

$$A^2 = w\mathbf{1} + vA = -L\mathbf{1} + 2a_0A \pmod q, \quad (20)$$

$$w = -L \pmod q,$$

$$v = 2a_0 \pmod q. \quad \text{q.e.d.}$$

14) [Theorem 2]

$$A^h = w_h\mathbf{1} + v_hA \pmod q$$

where h is an integer and $w_h, v_h \in \mathbf{Z}/q\mathbf{Z}$.

(Proof:)

From Theorem 1

$$A^2 = w_2\mathbf{1} + v_2A = -L\mathbf{1} + 2a_0A \pmod q. \quad (21)$$

If we can express A^h such that

$$A^h = w_h\mathbf{1} + v_hA \pmod q \in \mathcal{O}, \quad w_h, v_h \in \mathbf{Z}/q\mathbf{Z},$$

Then

$$\begin{aligned}
A^{h+1} &= (w_h\mathbf{1} + v_hA)A \pmod q \\
&= w_hA + v_h(-L\mathbf{1} + 2a_0A) \pmod q \\
&= -Lv_h\mathbf{1} + (w_h + 2a_0v_h)A \pmod q.
\end{aligned}$$

We have

$$\begin{aligned}w_{h+1} &= -L v_h \bmod q \in \mathbf{Z}/q\mathbf{Z}, \\v_{h+1} &= w_h + 2a_0 v_h \bmod q \in \mathbf{Z}/q\mathbf{Z}. \quad \text{q.e.d.}\end{aligned}$$

15) [**Theorem 3**]

$D \in O$ does not exist that satisfies the following equation.

$$B(AX) = DX \bmod q,$$

where $B, A, D \in O$, and X is a variable.

(*Proof:*)

When $X=1$, we have

$$BA = D \bmod q.$$

Then

$$B(AX) = (BA)X \bmod q.$$

We can select $C \in O$ that satisfies

$$B(AC) \neq (BA)C \bmod q. \quad (22)$$

We substitute $C \in O$ to X to obtain

$$B(AC) = (BA)C \bmod q. \quad (23)$$

(23) is contradictory to (22). q.e.d.

16) [**Theorem 4**]

$D \in O$ does not exist that satisfies the following equation.

$$C(B(AX)) = DX \bmod q \quad (24)$$

where $C, B, A, D \in O$, C has inverse $C^{-1} \bmod q$ and X is a variable.

B, A, C are non-associative, that is,

$$B(AC) \neq (BA)C \bmod q. \quad (25)$$

(*Proof:*)

If D exists, we have at $X=1$

$$C(BA)=D \text{ mod } q.$$

Then

$$C(B(AX))=(C(BA))X \text{ mod } q.$$

We substitute C to X to obtain

$$C(B(AC))=(C(BA))C \text{ mod } q.$$

From (10)

$$C(B(AC))=(C(BA))C=C((BA)C) \text{ mod } q$$

Multiplying C^{-1} from left side ,

$$B(AC)=(BA)C \text{ mod } q \quad (26)$$

(26) is contradictory to (25). q.e.d.

17) [Theorem 5]

D and $E \in O$ do not exist that satisfy the following equation.

$$C(B(AX))= E (DX) \text{ mod } q$$

where C, B, A, D and $E \in O$ have inverse and X is a variable.

A, B, C are non-associative, that is,

$$C(BA) \neq (CB)A \text{ mod } q. \quad (27)$$

(Proof:)

If D and E exist, we have at $X=\mathbf{1}$

$$C(BA)=ED \text{ mod } q \quad (28)$$

We have at $X=(ED)^{-1}=D^{-1}E^{-1} \text{ mod } q.$

$$C(B(A(D^{-1}E^{-1})))= E (D(D^{-1}E^{-1})) \text{ mod } q=\mathbf{1},$$

$$(C(B(A(D^{-1}E^{-1}))))^{-1} \text{ mod } q=\mathbf{1},$$

$$((ED)A^{-1})B^{-1}C^{-1} \text{ mod } q=\mathbf{1},$$

$$ED =(CB)A \text{ mod } q. \quad (29)$$

From (28) and (29) we have

$$C(BA) = (CB)A \pmod{q}. \quad (30)$$

(30) is contradictory to (27). q.e.d.

18) [Theorem 6]

$D \in O$ does not exist that satisfies the following equation.

$$A(B(A^{-1}X)) = DX \pmod{q}$$

where $B, A, D \in O$, A has inverse $A^{-1} \pmod{q}$ and X is a variable.

(Proof:)

If D exists, we have at $X=1$

$$A(BA^{-1}) = D \pmod{q}.$$

Then

$$A(B(A^{-1}X)) = (A(BA^{-1}))X \pmod{q}. \quad (31)$$

We can select $C \in O$ such that

$$(BA^{-1})(CA^2) \neq (BA^{-1})CA^2 \pmod{q}. \quad (32)$$

That is, (BA^{-1}) , C and A^2 are non-associative.

Substituting $X=CA$ in (31), we have

$$A(B(A^{-1}(CA))) = (A(BA^{-1}))(CA) \pmod{q}.$$

From Lemma 3

$$A(B((A^{-1}C)A)) = (A(BA^{-1}))(CA) \pmod{q}.$$

From (14)

$$A(B((A^{-1}C)A)) = A([(BA^{-1})C]A) \pmod{q}.$$

Multiply A^{-1} from left side we have

$$B((A^{-1}C)A) = ((BA^{-1})C)A \pmod{q}.$$

From Lemma 3

$$B(A^{-1}(CA)) = ((BA^{-1})C)A \pmod{q}.$$

Transforming CA to $((CA^2)A^{-1})$, we have

$$B(A^{-1}((CA^2)A^{-1}))=((BA^{-1})C)A \bmod q.$$

From (12) we have

$$((BA^{-1})(CA^2))A^{-1}=((BA^{-1})C)A \bmod q.$$

Multiply A from right side we have

$$((BA^{-1})(CA^2))=((BA^{-1})C)A^2 \bmod q. \quad (33)$$

(33) is contradictory to (32).

q.e.d.

§3. Concept of proposed fully homomorphic encryption scheme

Homomorphic encryption is a form of encryption which allows specific types of computations to be carried out on ciphertext and obtain an encrypted result which decrypted matches the result of operations performed on the plaintext. For instance, one person could add two encrypted numbers and then another person could decrypt the result, without either of them being able to find the value of the individual numbers.

§3.1 Definition of homomorphic encryption

A homomorphic encryption scheme $\mathbf{HE} := (\mathbf{KeyGen}; \mathbf{Enc}; \mathbf{Dec}; \mathbf{Eval})$ is a quadruple of PPT (Probabilistic polynomial time) algorithms.

In this paper the medium text space M_e of the encryption schemes will be octonion ring, and the functions to be evaluated will be represented as arithmetic circuits over this ring, composed of addition and multiplication gates. The syntax of these algorithms is given as follows.

-Key-Generation. The algorithm \mathbf{KeyGen} , on input the security parameter 1^λ , outputs $(\mathbf{sk}) \leftarrow \mathbf{KeyGen}(1^\lambda)$, where \mathbf{sk} is a secret encryption/decryption key.

-Encryption. The algorithm \mathbf{Enc} , on input system parameter $q=st$ where s and t are primes, secret keys (\mathbf{sk}) and a plaintext $p \in \mathbb{Z}_t\mathbb{Z}$, outputs a ciphertext $C \leftarrow \mathbf{Enc}(\mathbf{sk}; p)$.

-Decryption. The algorithm \mathbf{Dec} , on input system parameter q , secret key (\mathbf{sk}) and a ciphertext C , outputs a plaintext $p^* \leftarrow \mathbf{Dec}(\mathbf{sk}; C)$.

-Homomorphic-Evaluation. The algorithm \mathbf{Eval} , on input system parameter q , an arithmetic circuit ckt , and a tuple of n ciphertexts (C_1, \dots, C_n) , outputs a ciphertext $C' \leftarrow \mathbf{Eval}(\text{ckt}; C_1, \dots, C_n)$.

§3.2 Definition of fully homomorphic encryption

A scheme HE is fully homomorphic if it is both compact and homomorphic with respect to a class of circuits. More formally:

Definition (Fully homomorphic encryption). A homomorphic encryption scheme $FHE := (\mathbf{KeyGen}; \mathbf{Enc}; \mathbf{Dec}; \mathbf{Eval})$ is fully homomorphic if it satisfies the following properties:

1. Homomorphism: Let $CR = \{CR_\lambda\}_{\lambda \in \mathbb{N}}$ be the set of all polynomial sized arithmetic circuits. On input $\mathbf{sk} \leftarrow \mathbf{KeyGen}(1^\lambda), \forall \text{ckt} \in CR_\lambda, \forall (p_1, \dots, p_n) \in (\mathbb{Z}/t\mathbb{Z})^n$ where $n = n(\lambda)$ and $\forall (C_1, \dots, C_n)$ where $C_i \leftarrow \mathbf{Enc}(\mathbf{sk}; p_i)$, it holds that:

$$\Pr[\mathbf{Dec}(\mathbf{sk}; \mathbf{Eval}(\text{ckt}; C_1, \dots, C_n)) \neq \text{ckt}(p_1, \dots, p_n)] = \text{negl}(\lambda).$$

2. Compactness: There exists a polynomial $\mu = \mu(\lambda)$ such that the output length of \mathbf{Eval} is at most μ bits long regardless of the input circuit ckt and the number of its inputs.

§3.3 Proposed fully homomorphic enciphering/deciphering functions

We propose a fully homomorphic encryption (FHE) scheme based on the enciphering/deciphering functions on octonion ring over $\mathbb{Z}/q\mathbb{Z}$.

First we define the secret parameters $B \in O$ and $G \in O$ as follows. Let s and t be 1000-digit secret primes, and $q=st$ be published as a system parameter

where

$$s < t.$$

Let k and h be the integers that satisfy the following equation from chinese remainder theorem,

$$ks+ht = 1 \pmod{q},$$

$$k, h \in \mathbb{Z}/q\mathbb{Z}.$$

We select the element $B = (b_0, b_1, \dots, b_7) \in O$ and $G = (g_0, -b_1, \dots, -b_7) \in O$ such that

$$L_B := |B|^2 = b_0^2 + b_1^2 + \dots + b_7^2 \pmod{q} = 0,$$

$$L_G := |G|^2 = g_0^2 + b_1^2 + \dots + b_7^2 \pmod{q}$$

$$\begin{aligned}
&= g_0^2 - b_0^2 \neq 0 \pmod{q}. \\
&\text{GCD}(b_i, q) = 1 \quad (i=0,1), \\
&B+G = (b_0 + g_0)\mathbf{1} \pmod{q}. \tag{34}
\end{aligned}$$

From theorem1 we have

$$\begin{aligned}
B^2 &= 2b_0B \pmod{q}, \\
G^2 &= -(g_0^2 - b_0^2)\mathbf{1} + 2g_0G \pmod{q}.
\end{aligned}$$

To multiply (34) by B from right side, we have

$$\begin{aligned}
(B+G)B &= (b_0 + g_0)B \pmod{q}, \\
B^2 + GB &= (b_0 + g_0)B \pmod{q}, \\
2b_0B + GB &= (b_0 + g_0)B \pmod{q}, \\
GB &= (-b_0 + g_0)B \pmod{q}.
\end{aligned}$$

To multiply (34) by B from left side, we have

$$\begin{aligned}
B(B+G) &= (b_0 + g_0)B \pmod{q}, \\
B^2 + BG &= (b_0 + g_0)B \pmod{q}, \\
2b_0B + BG &= (b_0 + g_0)B \pmod{q}, \\
BG &= (-b_0 + g_0)B \pmod{q} = GB \pmod{q}.
\end{aligned}$$

Let $d \in \mathbf{Z}/s\mathbf{Z}, u, v$ and $w \in \mathbf{Z}/q\mathbf{Z}$ be the random numbers. Let $p \in \mathbf{Z}/t\mathbf{Z}$ be a plaintext and $f, y, z \in \mathbf{Z}/q\mathbf{Z}$ are defined such that

$$f = y + z \pmod{q},$$

where

$$y = psk \pmod{q} = u + 2b_0v + (g_0 - b_0)w \pmod{q} \in \mathbf{Z}/q\mathbf{Z},$$

$$z = dth \pmod{q} = u + (g_0 + b_0)w \pmod{q} \in \mathbf{Z}/q\mathbf{Z}.$$

Then f is given such that

$$f = 2u + 2b_0v + 2g_0w \pmod{q} \in \mathbf{Z}/q\mathbf{Z}.$$

We define the medium text M by

$$M = (m_0, \dots, m_7) := R_1(\dots(R_r([u\mathbf{1} + vB + wG]))R_r^{-1})\dots R_1^{-1} \pmod{q} \in O,$$

where

$R_i \in O$ such that R_i^{-1} exists ($i=1, \dots, r$) and

$$R_i B \neq B R_i \pmod{q} (i=1, \dots, r),$$

$$R_i G \neq G R_i \pmod{q} (i=1, \dots, r).$$

As $M = u\mathbf{1} + vB + wG = (u + b_0v + g_0w, vb_1 - wb_1, \dots, vb_7 - wb_7) \pmod{q}$, we can calculate $|M|^2 \pmod{q}$ as follows.

$$\begin{aligned} |M|^2 \pmod{q} &= (u + b_0v + g_0w)^2 + (vb_1 - wb_1)^2 + \dots + (vb_7 - wb_7)^2 \pmod{q} \\ &= (u + b_0v + g_0w)^2 + (v-w)^2(b_1^2 + \dots + b_7^2) \pmod{q} \\ &= (u + b_0v + g_0w)^2 - ((v-w)b_0)^2 \pmod{q} \\ &= (u + b_0v + g_0w + (v-w)b_0)(u + b_0v + g_0w - (v-w)b_0) \pmod{q} \\ &= (u + 2b_0v + (g_0 - b_0)w)(u + (g_0 + b_0)w) \pmod{q} \\ &= yz \pmod{q} \\ &= (psk)(dth) \pmod{q} \\ &= pdkhq \pmod{q} \\ &= 0 \pmod{q}. \end{aligned}$$

As $u\mathbf{1} + vB + wG = (u + b_0v + g_0w, vb_1 - wb_1, \dots, vb_7 - wb_7)$, we have

$$2[u\mathbf{1} + vB + wG]_0 = 2u + 2b_0v + 2g_0w \pmod{q} \in \mathbf{Z}/q\mathbf{Z} = f$$

and

$$f \pmod{t} = psk + dth \pmod{t} = p$$

where $[M]_i$ denotes the i -th element of $M \in O$.

Here we show a simple numerical example to help with the understanding of the reader.

$$\text{Let } s=7, t=11, q=st=77, k=8, h=2, (8*7+2*11=1 \pmod{77}).$$

$$\text{Let } B=(39,19,45,15,4,0,1,3), G=(40,58,32,62,73,0,76,74).$$

That is, $b_0=1/2 = 39 \pmod{77}$, $g_0=3/2 = 40 \pmod{77}$. We select p_1, d_1 and w_1 as follows.

$$p_1=4, d_1=3, w_1=19,$$

$$y_1=4*(8*7) \bmod 77=70=u_1+v_1+w_1 \bmod q,$$

$$z_1=3*(2*11) \bmod 77=66 =u_1+2w_1 \bmod q,$$

$$f_1= y_1+ z_1 \bmod 77=59.$$

As $w_1=19$, we have $u_1=28, v_1=23$.

Then we have,

$$M_1= (u_1+v_1/2+3w_1/2, v_1b_1-w_1b_1, \dots, v_1b_7-w_1b_7)$$

$$= (19, 4b_1, \dots, 4b_7)$$

$$= (68, 76, 26, 60, 16, 0, 4, 12),$$

$$f_1=2*[M_1]_0 \bmod 77=2*68 \bmod 77=59,$$

$$p_1=f_1 \bmod 11=4,$$

$$|M_1|^2 \bmod 77=68^2+4^2(b_1^2+\dots+b_7^2) \bmod 77$$

$$=4-16*(1/2)^2$$

$$=4-4 =0 \bmod 77.$$

Next we select p_2, d_2, w_2 as follows.

$$p_2=8, d_2=5, w_2=43,$$

$$y_2=8*(8*7) =63 \bmod 77=u_2+v_2+w_2 \bmod 77,$$

$$z_2=5*(2*11) =33 \bmod 77=u_2+2w_2 \bmod 77,$$

$$f_2= y_2+ z_2 \bmod 77=19.$$

As $w_2=43$, we have $u_2=24, v_2=73$.

Then we have,

$$M_2=(u_2+v_2/2+3w_2/2, v_2b_1-w_2b_1, \dots, v_2b_7-w_2b_7)$$

$$=(48, 30b_1, \dots, 30b_7)$$

$$=(48, 31, 41, 65, 43, 0, 30, 13),$$

$$f_2=2*[M_2]_0 \bmod 77=2*48 \bmod 77=19.$$

$$p_2=f_2 \bmod 11=8,$$

$$\begin{aligned}
|M_2|^2 \bmod 77 &= 48^2 + 30^2(b_1^2 + \dots + b_7^2) \bmod 77 \\
&= 71 - 53 \cdot (1/2)^2 \\
&= 71 - 71 = 0 \bmod 77.
\end{aligned}$$

$$M_1 + M_2 \bmod 77 = (39, 30, 67, 48, 59, 0, 34, 25),$$

$$|M_1 + M_2|^2 \bmod 77 = 0,$$

$$p_1 + p_2 = 4 + 8 = 1 \bmod 11,$$

$$p_{1+2} = 2 * [M_1 + M_2]_0 = 2 * 39 \bmod 11 = 1 \bmod 11.$$

$$M_1 M_2 \bmod 77 = (60, 58, 32, 62, 73, 0, 76, 74),$$

$$|M_1 M_2|^2 \bmod 77 = 0,$$

$$p_1 p_2 = 4 * 8 = 10 \bmod 11,$$

$$p_{12} = 2 * [M_1 M_2]_0 = 2 * 60 = 10 \bmod 11. \quad \square$$

Here we simplify the expression of medium text M such that

$$M := \mathbf{R}([u\mathbf{1} + vB + wG])\mathbf{R}^{-1} \in O.$$

We show relation between M and p .

$$(m'_0, m'_1, \dots, m'_7) := \mathbf{R}_r^{-1}(\dots(\mathbf{R}_1^{-1}(M \mathbf{R}_1)\dots)\mathbf{R}_r) = u\mathbf{1} + vB + wG \in O,$$

$$m'_0 + m'_1 (b_0/b_1) = u + b_0v + g_0w + (vb_1 - wb_1) (b_0/b_1) \bmod q$$

$$= u + 2b_0v + (g_0 - b_0)w \bmod q$$

$$= y = psk \bmod q.$$

Then we have

$$y \bmod t = p.$$

$$m'_0 - m'_1 (b_0/b_1) = u + b_0v + g_0w - (vb_1 - wb_1) (b_0/b_1) \bmod q$$

$$= u + (g_0 + b_0)w \bmod q$$

$$= z = dth \bmod q.$$

$$z \bmod s = d.$$

f is given such that

$$f=y+z=2u+2b_0v+2g_0w \pmod q.$$

Let

$$M_1:=\mathbf{R}(u_1\mathbf{1}+v_1B+w_1G)\mathbf{R}^{-1}\in O,$$

$$M_2:=\mathbf{R}(u_2\mathbf{1}+v_2B+w_2G)\mathbf{R}^{-1}\in O.$$

We have

$$\begin{aligned} M_1M_2 &= \mathbf{R}[(u_1u_2-(g_0^2-b_0^2)w_1w_2)\mathbf{1} \\ &\quad +(u_1v_2+v_1u_2+2b_0v_1v_2+(-b_0+g_0)(v_1w_2+w_1v_2))B \\ &\quad +(u_1w_2+w_1u_2+2g_0w_1w_2)G]\mathbf{R}^{-1}=M_2M_1 \pmod q. \end{aligned}$$

We show the above equation by using Lemma 5b as follows.

$$[\mathbf{R}B\mathbf{R}^{-1}][\mathbf{R}B\mathbf{R}^{-1}]=\mathbf{R}B^2\mathbf{R}^{-1} \pmod{ (=2b_0\mathbf{R}B\mathbf{R}^{-1} \pmod q)},$$

$$[\mathbf{R}G\mathbf{R}^{-1}][\mathbf{R}G\mathbf{R}^{-1}]=\mathbf{R}G^2\mathbf{R}^{-1} \pmod{ (= \mathbf{R}[2g_0G-(g_0^2-b_0^2)\mathbf{1}]\mathbf{R}^{-1} \pmod q)}.$$

From $B+G=(b_0+g_0)\mathbf{1} \pmod q$,

$$[\mathbf{R}B\mathbf{R}^{-1}+\mathbf{R}G\mathbf{R}^{-1}]=[\mathbf{R}(B+G)\mathbf{R}^{-1}]=(b_0+g_0)\mathbf{1} \pmod q. \quad (35)$$

We multiply $[\mathbf{R}B\mathbf{R}^{-1}]$ from right side, we have

$$[\mathbf{R}B\mathbf{R}^{-1}+\mathbf{R}G\mathbf{R}^{-1}][\mathbf{R}B\mathbf{R}^{-1}]=(b_0+g_0)\mathbf{1}[\mathbf{R}B\mathbf{R}^{-1}] \pmod q,$$

$$2b_0[\mathbf{R}B\mathbf{R}^{-1}]+\mathbf{R}G\mathbf{R}^{-1}[\mathbf{R}B\mathbf{R}^{-1}]=(b_0+g_0)[\mathbf{R}B\mathbf{R}^{-1}] \pmod q,$$

$$[\mathbf{R}G\mathbf{R}^{-1}][\mathbf{R}B\mathbf{R}^{-1}]=(-b_0+g_0)[\mathbf{R}B\mathbf{R}^{-1}] \pmod q.$$

From $(-b_0+g_0)B=GB \pmod q$,

$$[\mathbf{R}G\mathbf{R}^{-1}][\mathbf{R}B\mathbf{R}^{-1}]=(-b_0+g_0)[\mathbf{R}(B)\mathbf{R}^{-1}]=[\mathbf{R}(GB)\mathbf{R}^{-1}] \pmod q.$$

We multiply $[\mathbf{R}B\mathbf{R}^{-1}]$ from left side of (35), we have

$$[\mathbf{R}B\mathbf{R}^{-1}][\mathbf{R}B\mathbf{R}^{-1}+\mathbf{R}G\mathbf{R}^{-1}]=(b_0+g_0)[\mathbf{R}B\mathbf{R}^{-1}]\mathbf{1} \pmod q,$$

$$2b_0[\mathbf{R}B\mathbf{R}^{-1}]+\mathbf{R}B\mathbf{R}^{-1}[\mathbf{R}G\mathbf{R}^{-1}]=(b_0+g_0)[\mathbf{R}B\mathbf{R}^{-1}] \pmod q,$$

$$[\mathbf{R}B\mathbf{R}^{-1}][\mathbf{R}G\mathbf{R}^{-1}]=(-b_0+g_0)[\mathbf{R}B\mathbf{R}^{-1}] \pmod q.$$

From $(-b_0+g_0)B=BG \pmod q$

$$[\mathbf{RBR}^{-1}][\mathbf{RGR}^{-1}] = (-b_0+g_0)[\mathbf{R(B)R}^{-1}]=[\mathbf{R(BG)R}^{-1}] \pmod q (= [\mathbf{R(GB)R}^{-1}]).$$

Then we have

$$\begin{aligned} M_1M_2 &= [\mathbf{R}(u_1\mathbf{1}+v_1B+w_1G)\mathbf{R}^{-1}][\mathbf{R}(u_2\mathbf{1}+v_2B+w_2G)\mathbf{R}^{-1}] \\ &= [u_1\mathbf{R1R}^{-1}+v_1\mathbf{RBR}^{-1}+w_1\mathbf{RGR}^{-1}][u_2\mathbf{R1R}^{-1}+v_2\mathbf{RBR}^{-1}+w_2\mathbf{RGR}^{-1}] \\ &= [u_1\mathbf{R1R}^{-1}][u_2\mathbf{R1R}^{-1}]+[u_1\mathbf{R1R}^{-1}][v_2\mathbf{RBR}^{-1}]+[u_1\mathbf{R1R}^{-1}][w_2\mathbf{RGR}^{-1}] \\ &\quad +[v_1\mathbf{RBR}^{-1}][u_2\mathbf{R1R}^{-1}]+[v_1\mathbf{RBR}^{-1}][v_2\mathbf{RBR}^{-1}]+[v_1\mathbf{RBR}^{-1}][w_2\mathbf{RGR}^{-1}] \\ &\quad +[w_1\mathbf{RGR}^{-1}][u_2\mathbf{R1R}^{-1}]+[w_1\mathbf{RGR}^{-1}][v_2\mathbf{RBR}^{-1}]+[w_1\mathbf{RGR}^{-1}][w_2\mathbf{RGR}^{-1}] \\ &= [\mathbf{R}(u_1u_2)\mathbf{1R}^{-1}]+[\mathbf{R}(u_1v_2)\mathbf{BR}^{-1}]+[\mathbf{R}(u_1w_2)\mathbf{GR}^{-1}] \\ &\quad +[\mathbf{R}(v_1u_2)\mathbf{BR}^{-1}]+[\mathbf{R}(v_1v_2)\mathbf{BBR}^{-1}]+[\mathbf{R}(v_1w_2)\mathbf{BGR}^{-1}] \\ &\quad +[\mathbf{R}(w_1u_2)\mathbf{GR}^{-1}]+[\mathbf{R}(w_1v_2)\mathbf{GBR}^{-1}]+[\mathbf{R}(w_1w_2)\mathbf{GGR}^{-1}] \\ &= \mathbf{R}[(u_1u_2)\mathbf{1}+(u_1v_2)B+(u_1w_2)G+(v_1u_2)B+2b_0(v_1v_2)B+(-b_0+g_0)(v_1w_2)B \\ &\quad +(w_1u_2)G+(-b_0+g_0)(w_1v_2)B+(w_1w_2)[-(g_0^2-b_0^2)\mathbf{1}+2g_0G]\mathbf{R}^{-1}] \\ &= \mathbf{R}[(u_1u_2-(g_0^2-b_0^2)w_1w_2)\mathbf{1}+(u_1v_2+v_1u_2+2b_0v_1v_2+(-b_0+g_0)(v_1w_2+w_1v_2))B \\ &\quad +(u_1w_2+w_1u_2+2g_0w_1w_2)G]\mathbf{R}^{-1}. \quad \square \end{aligned}$$

$$\begin{aligned} \text{Let } (m^*_0, m^*_1, \dots, m^*_7) &:= R_r^{-1}(\dots(R_1^{-1}([M_1M_2]R_1)\dots)R_r \pmod q \\ &= (u_1\mathbf{1}+v_1B+w_1G)(u_2\mathbf{1}+v_2B+w_2G) \\ &= (u_1u_2-(g_0^2-b_0^2)w_1w_2)\mathbf{1} \\ &\quad +(u_1v_2+v_1u_2+2b_0v_1v_2+(-b_0+g_0)(v_1w_2+w_1v_2))B \\ &\quad +(u_1w_2+w_1u_2+2g_0w_1w_2)G \pmod q. \end{aligned} \tag{36}$$

We can show that we obtain p_1p_2 , the multiple of p_1 and p_2 from $(m^*_0, m^*_1, \dots, m^*_7)$ as follows.

$$y_1 = p_1sk \pmod q = u_1+2b_0v_1+(g_0-b_0)w_1 \pmod q \in \mathbf{Z}/q\mathbf{Z},$$

$$z_2 = d_2th \pmod q = u_2+(g_0+b_0)w_2 \pmod q \in \mathbf{Z}/q\mathbf{Z}.$$

Then we have

$$y_1z_2 = (p_1sk)(d_2th) \pmod q = 0 = (u_1+2b_0v_1+(g_0-b_0)w_1)(u_2+(g_0+b_0)w_2) \pmod q.$$

In the same manner we have

$$y_2 z_1 = (p_2 s k)(d_1 t h) \pmod q = 0 = (u_2 + 2b_0 v_2 + (g_0 - b_0)w_2)(u_1 + (g_0 + b_0)w_1) \pmod q.$$

From (36)

$$\begin{aligned} 2m^*_0 &= 2(u_1 u_2 - (g_0^2 - b_0^2)w_1 w_2) + 2(u_1 v_2 + v_1 u_2 + 2b_0 v_1 v_2 + (-b_0 + g_0)(v_1 w_2 + w_1 v_2)) b_0 \\ &\quad + 2(u_1 w_2 + w_1 u_2 + 2g_0 w_1 w_2) g_0 \pmod q. \end{aligned}$$

Here we calculate $2m^*_0 = 2m^*_0 + y_1 z_2 + y_2 z_1 \pmod q$ by adding $y_1 z_2 + y_2 z_1$ as follows.

$$\begin{aligned} 2m^*_0 &= 2m^*_0 + y_1 z_2 + y_2 z_1 \pmod q \\ &= 2(u_1 u_2 - (g_0^2 - b_0^2)w_1 w_2) + 2(u_1 v_2 + v_1 u_2 + 2b_0 v_1 v_2 + (-b_0 + g_0)(v_1 w_2 + w_1 v_2)) b_0 \\ &\quad + 2(u_1 w_2 + w_1 u_2 + 2g_0 w_1 w_2) g_0 \\ &\quad + (u_1 + 2b_0 v_1 + (g_0 - b_0)w_1)(u_2 + (g_0 + b_0)w_2) \\ &\quad + (u_2 + 2b_0 v_2 + (g_0 - b_0)w_2)(u_1 + (g_0 + b_0)w_1) \pmod q \\ &= 4u_1 u_2 + 4b_0 (u_1 v_2 + v_1 u_2) + (4b_0^2) v_1 v_2 \\ &\quad + (2b_0 (-b_0 + g_0) + 2b_0 (g_0 + b_0))(v_1 w_2 + w_1 v_2) \\ &\quad + (2g_0 + 2g_0)(u_1 w_2 + w_1 u_2) + (4g_0^2 - 2(g_0^2 - b_0^2) + 2(g_0^2 - b_0^2))w_1 w_2 \pmod q \\ &= 4u_1 u_2 + 4b_0 (u_1 v_2 + v_1 u_2) + (4b_0^2) v_1 v_2 + 4b_0 g_0 (v_1 w_2 + w_1 v_2) \\ &\quad + 4g_0 (u_1 w_2 + w_1 u_2) + 4g_0^2 w_1 w_2 \pmod q \\ &= (2u_1 + 2b_0 v_1 + 2g_0 w_1) (2u_2 + 2b_0 v_2 + 2g_0 w_2) \\ &= f_1 f_2 \pmod q \in \mathbf{Z}/q\mathbf{Z}. \end{aligned}$$

Then we have

$$2m^*_0 \pmod t = f_1 f_2 \pmod t = p_1 p_2 \pmod t.$$

And we have $y_1 y_2$ and $z_1 z_2$ from $m^*_0 + m^*_1 (b_0/b_1)$ and $m^*_0 - m^*_1 (b_0/b_1)$ respectively as follows.

$$\begin{aligned} &m^*_0 + m^*_1 (b_0/b_1) \pmod q \\ &= (u_1 u_2 - (g_0^2 - b_0^2)w_1 w_2) + (u_1 v_2 + v_1 u_2 + 2b_0 v_1 v_2 + (-b_0 + g_0)(v_1 w_2 + w_1 v_2)) b_0 \\ &\quad + (u_1 w_2 + w_1 u_2 + 2g_0 w_1 w_2) g_0 \\ &\quad + [(u_1 v_2 + v_1 u_2 + 2b_0 v_1 v_2 + (-b_0 + g_0)(v_1 w_2 + w_1 v_2))b_1 - (u_1 w_2 + w_1 u_2 + 2g_0 w_1 w_2)b_1](b_0/b_1) \\ &\pmod q \end{aligned}$$

$$\begin{aligned}
&= u_1u_2+2b_0(u_1v_2+v_1u_2)+4b_0v_1v_2+b_0(-b_0+g_0)v_1w_2+w_1v_2) \\
&\quad +(-b_0+g_0)(u_1w_2+w_1u_2)+(-b_0+g_0)^2w_1w_2 \\
&= (u_1+2b_0v_1+(g_0-b_0)w_1)(u_2+2b_0v_2+(g_0-b_0)w_2) \bmod q \\
&= y_1y_2 \in \mathbf{Z}/q\mathbf{Z}. \tag{37a}
\end{aligned}$$

We notice that

$$y_1y_2 \bmod t = (y_1 \bmod t)(y_2 \bmod t) \bmod t = p_1p_2 \bmod t.$$

$$\begin{aligned}
&m^*_0 - m^*_1(b_0/b_1) \bmod q \\
&= (u_1u_2 - (g_0^2 - b_0^2)w_1w_2) + (u_1v_2 + v_1u_2 + 2b_0v_1v_2 + (-b_0 + g_0)(v_1w_2 + w_1v_2))b_0 \\
&\quad + (u_1w_2 + w_1u_2 + 2g_0w_1w_2)g_0 \\
&\quad - [(u_1v_2 + v_1u_2 + 2b_0v_1v_2 + (-b_0 + g_0)(v_1w_2 + w_1v_2))b_1 - (u_1w_2 + w_1u_2 + 2g_0w_1w_2)b_1](b_0/b_1) \\
&\bmod q \\
&= u_1u_2 + (b_0 + g_0)(u_1w_2 + w_1u_2) + (b_0 + g_0)^2w_1w_2 \\
&= (u_1 + (g_0 + b_0)w_1)(u_2 + (g_0 + b_0)w_2) \bmod q \\
&= z_1z_2 \bmod q \in \mathbf{Z}/q\mathbf{Z}. \tag{37b}
\end{aligned}$$

From $y_1 = p_1sk \bmod q$, $y_2 = p_2sk \bmod q$, $z_1 = d_1th \bmod q$ and $z_2 = d_2th \bmod q$, we also have

$$\begin{aligned}
f_{12} &:= y_1y_2 + z_1z_2 \bmod q \\
&= (y_1 + z_1)(y_2 + z_2) \bmod q \\
&= f_1f_2 \bmod q \in \mathbf{Z}/q\mathbf{Z}. \tag{37c}
\end{aligned}$$

Here we define the some parameters for describing FHE. Let s and t ($s < t$) be secret 1000-digit primes. Let $q = st$ be a 2000-digit composite number to be published as a system parameter. Let $M = (m_0, m_1, \dots, m_7) := \mathbf{R}(u\mathbf{1} + v\mathbf{B} + w\mathbf{G})\mathbf{R}^{-1} \in O$ be the medium text where

$$\begin{aligned}
f &= y + z \bmod q \in \mathbf{Z}/q\mathbf{Z}, \\
y &= psk \bmod q = u + 2b_0v + (g_0 - b_0)w \bmod q \in \mathbf{Z}/q\mathbf{Z}, \\
z &= dth \bmod q = u + (g_0 + b_0)w \bmod q \in \mathbf{Z}/q\mathbf{Z}.
\end{aligned}$$

Let $p \in \mathbf{Z}/t\mathbf{Z}$ be a plaintext, $d \in \mathbf{Z}/s\mathbf{Z}$ and $w \in \mathbf{Z}/q\mathbf{Z}$ be secret random numbers, $u, v \in \mathbf{Z}/q\mathbf{Z}$ be secret numbers such that

$$\begin{aligned} u &= dth - (g_0 + b_0)w \pmod{q} \in \mathbf{Z}/q\mathbf{Z}, \\ v &= (psk - u - (g_0 - b_0)w)(2b_0)^{-1} \pmod{q} \in \mathbf{Z}/q\mathbf{Z}, \end{aligned}$$

where

$$sk + ht = 1 \pmod{q}.$$

Let $X = (x_0, \dots, x_7) \in O[X]$ be a variable. Let $E(p, X)$ and $D(X)$ be an enciphering and a deciphering function. Let $C(X) = E(p, X) \in O[X]$ be the ciphertext. $A_i, Z_i \in O$ are selected randomly such that A_i^{-1} and Z_i^{-1} exist ($i=1, \dots, k$) which are the secret keys.

Enciphering function $E(p, X) = C(X)$ is defined as follows.

$$E(p, X) = C(X) :=$$

$$\begin{aligned} & A_1((\dots((A_k((M[(A_k^{-1}((\dots((A_1^{-1}X)Z_1))\dots))Z_k])Z_k^{-1})).\dots))Z_1^{-1}) \pmod{q} \in O[X] \\ & = (e_{00}x_0 + e_{01}x_1 + \dots + e_{07}x_7, \\ & \quad e_{10}x_0 + e_{11}x_1 + \dots + e_{17}x_7, \\ & \quad \dots \quad \dots \\ & \quad e_{70}x_0 + e_{71}x_1 + \dots + e_{77}x_7), \tag{38} \\ & = \{e_{ij}\} (i, j=0, \dots, 7) \end{aligned}$$

with $e_{ij} \in \mathbf{Z}/q\mathbf{Z}$ ($i, j=0, \dots, 7$) which is published in cloud centre, where

$$(m'_0, m'_1, \dots, m'_7) := R_r^{-1}(\dots(R_1^{-1}(MR_1)\dots)R_r = (u\mathbf{1} + vB + wG) \pmod{q} \in O.$$

Here we notice how to construct enciphering function. We show a part of process for constructing enciphering function $E(p, X)$ as follows.

$$\begin{aligned} & A_1^{-1}X \\ & (A_1^{-1}X)Z_1 \\ & A_2^{-1}((A_1^{-1}X)Z_1) \\ & (A_2^{-1}((A_1^{-1}X)Z_1))Z_2 \\ & \dots \\ & (A_k^{-1}((\dots((A_1^{-1}X)Z_1))\dots))Z_k \\ & M[(A_k^{-1}((\dots((A_1^{-1}X)Z_1))\dots))Z_k] \\ & (M[(A_k^{-1}((\dots((A_1^{-1}X)Z_1))\dots))Z_k])Z_k^{-1} \\ & A_k(M[(A_k^{-1}((\dots((A_1^{-1}X)Z_1))\dots))Z_k])Z_k^{-1} \end{aligned}$$

$$A_1(\dots((A_k((M[(A_k^{-1}(\dots((A_1^{-1}X)Z_1))\dots))Z_k])Z_k^{-1}))\dots))Z_1^{-1}$$

Let D be the deciphering function defined as follows .

$$\begin{aligned} D_1(X) &:= A_k^{-1}(\dots((A_1^{-1}X)Z_1))\dots)Z_k, \\ D_2(X) &:= A_1(\dots((A_k(X Z_k^{-1}))\dots))Z_1^{-1}, \\ D(X) &:= D_1(C(D_2(X)) \bmod q) = MX. \end{aligned} \quad (39)$$

$$\begin{aligned} D(\mathbf{1}) &= M = (m_0, m_1, \dots, m_7) = R_1(\dots(R_r(u\mathbf{1} + vB + wG)R_r^{-1})\dots)R_1^{-1} \\ &= \mathbf{R}[u\mathbf{1} + vB + wG]\mathbf{R}^{-1} = \mathbf{R}[u\mathbf{1} + v(b_0, b_1, \dots, b_7) + w(g_0, -b_1, \dots, -b_7)]\mathbf{R}^{-1} \end{aligned}$$

where

$$\begin{aligned} \text{GCD}(b_i, q) &= 1 \quad (i=0,1), \\ g_0^2 - b_0^2 &\neq 0 \bmod q. \end{aligned}$$

The norm of the ciphertext $C(X)$ is given such that

$$\begin{aligned} |C(X)|^2 &= |A_1(\dots((A_k((M[(A_k^{-1}(\dots((A_1^{-1}X)Z_1))\dots))Z_k])Z_k^{-1}))\dots))Z_1^{-1}|^2 \bmod q \\ &= (|A_1| \dots |A_k| |M| |A_k|^{-1} \dots |A_1|^{-1} |X| |Z_k| \dots |Z_k| |Z_k|^{-1} \dots |Z_1|^{-1})^2 \bmod q \\ &= |M|^2 |X|^2 \bmod q = 0 \end{aligned}$$

because

$$|M|^2 = yz = (psk)(dth) = pdkhq \bmod q = 0 \bmod q.$$

Then we can obtain the plaintext p as follows.

Let

$$(m'_0, m'_1, \dots, m'_7) := R_r^{-1}(\dots(R_1^{-1}(M R_1)\dots)R_r = u\mathbf{1} + vB + wG \in O.$$

From $(m'_0, m'_1, \dots, m'_7)$, we obtain y, z, f and the plaintext p .

$$\begin{aligned} m'_0 + m'_1 (b_0/2b_1) &= u + b_0v + g_0w + (vb_1 - wb_1) (b_0/2b_1) \bmod q \\ &= u + 2b_0v + (g_0 - b_0)w \bmod q \\ &= y \bmod q \in \mathbf{Z}/q\mathbf{Z}. \end{aligned} \quad (40a)$$

$$\begin{aligned} m'_0 - m'_1 (b_0/2b_1) &= u + b_0v + g_0w - (vb_1 - wb_1) (b_0/2b_1) \\ &= u + (g_0 + b_0)w \bmod q \end{aligned}$$

$$=z \bmod q \in \mathbf{Z}/q\mathbf{Z}. \quad (40b)$$

$$f=y+z \bmod q \in \mathbf{Z}/q\mathbf{Z}. \quad (40c)$$

Here we have

$$f=y+z=2m'_0 \bmod q=2u+2b_0v+2g_0w \bmod q \in \mathbf{Z}/q\mathbf{Z},$$

$$2m'_0 \bmod t=f \bmod t=y \bmod t=psk \bmod t=p.$$

§3.4 Assumption

Here we describe the assumption on which the proposed scheme bases.

§3.4.1 Factoring assumption $\text{Fact}(q)$

Let q be as a large composite number where $q = st$ with $q = q(\lambda)$, where λ is a security parameter, and s and t ($s < t$) are prime numbers.

In the $\text{Fact}(q)$ assumption, the PPT(Probabilistic polynomial time) algorithm AL is given q and the goal is to find primes s and t .

For a parameter $q = q(\lambda)$ defined in terms of the security parameter λ and for any PPT algorithm AL, we have

$$\Pr[q=st \text{ with } q = q(\lambda) : (s, t) \leftarrow \text{AL}(1^\lambda, q)] = \text{negl}(\lambda). \quad (41)$$

§3.4.2 Elements on octonion ring assumption $\text{EOR}(k, r, n; q)$

Let q be a 2000-digit composite number. Let k, r and n be integer parameters. Let $\mathbf{A} := (A_1, \dots, A_k) \in O^k$, $\mathbf{Z} := (Z_1, \dots, Z_k) \in O^k$, $\mathbf{R} := (R_1, \dots, R_r) \in O^r$. Let $C_i(X) := E(p_i, X) = (A_1((\dots((A_k(M_i[(A_k^{-1}((\dots((A_1^{-1}X)Z_1))\dots))Z_k]))Z_k^{-1})))\dots))Z_1^{-1} \bmod q \in O[X]$ where medium text $M_i = (m_{i0}, \dots, m_{i7}) := R_1(\dots(R_r(u_i \mathbf{1} + v_i B + w_i G)R_r^{-1})\dots)R_1^{-1} \bmod q \in O$, plaintext $p_i = y_i + z_i = 2u_i + 2b_0v_i + 2g_0w_i \bmod q$, $y_i = p_i sk = u_i + 2b_0v_i + (g_0 - b_0)w_i \bmod q$, $z_i = d_i th = u_i + (g_0 + b_0)w_i \bmod q$ ($i=1, \dots, n$), $sk + th = 1 \bmod q$, and X is a variable.

In the $\text{EOR}(k, r, n; q)$ assumption, the PPT adversary A_d is given $C_i(X)$ ($i=1, \dots, n$) randomly and his goal is to find a set of elements $\mathbf{A} = (A_1, \dots, A_k) \in O^k$, $\mathbf{Z} = (Z_1, \dots, Z_k) \in O^k$, $\mathbf{R} = (R_1, \dots, R_r) \in O^r$, with the order of the elements $A_1, \dots, A_k, Z_1, \dots, Z_k, R_1, \dots, R_r$ and plaintexts p_i ($i=1, \dots, n$). For parameters $k = k(\lambda)$, $r = r(\lambda)$ and $n = n(\lambda)$ defined in terms of the security parameter λ and for any PPT adversary A_d we have

$$\Pr[(A_1((\dots((A_k(M_i[(A_k^{-1}((\dots((A_1^{-1}X)Z_1))\dots))Z_k]))Z_k^{-1})))\dots))Z_1^{-1} \bmod q = C_i(X) \quad (i=1, \dots, n) : \mathbf{A}, \mathbf{Z}, \mathbf{R}, p_i(i=1, \dots, n) \leftarrow A_d(1^\lambda, C_i(X) (i=1, \dots, n))] = \text{negl}(\lambda). \quad (42)$$

To solve directly $\mathbf{EOR}(k,r,n;q)$ assumption is known to be the problem for solving the multivariate algebraic equations of high degree which is known to be NP-hard.

§3.5 Syntax of proposed algorithms

The syntax of proposed scheme is given as follows.

-Key-Generation. The algorithm **KeyGen**, on input the security parameter 1^λ and system parameter q , outputs $\mathbf{sk}=(A,Z,R,B,G,s,t)\leftarrow\mathbf{KeyGen}(1^\lambda)$, where \mathbf{sk} is a secret encryption /decryption key and $q=st$.

-Encryption. The algorithm **Enc**, on input system parameter q and secret keys $\mathbf{sk}=(A,Z,R,B,G,s,t)$ and a plaintext $p\in\mathbf{Z}/q\mathbf{Z}$, outputs a ciphertext $C(X;\mathbf{sk},p)\leftarrow\mathbf{Enc}(\mathbf{sk};p)$.

-Decryption. The algorithm **Dec**, on input system parameter q , secret keys \mathbf{sk} and a ciphertext $C(X;\mathbf{sk},p)$, outputs plaintext $\mathbf{Dec}(\mathbf{sk}; C(X;\mathbf{sk},p))$ where $C(X;\mathbf{sk},p)\leftarrow\mathbf{Enc}(\mathbf{sk}; p)$.

-Homomorphic-Evaluation. The algorithm **Eval**, on input system parameter q , an arithmetic circuit ckt, and a tuple of n ciphertexts (C_1,\dots, C_n) , outputs an evaluated ciphertext $C'\leftarrow\mathbf{Eval}(\text{ckt}; C_1,\dots, C_n)$ where $C_i=C(X;\mathbf{sk},p_i)$ ($i=1,\dots,n$).

[Theorem 7]

For any $p, p' \in \mathbf{Z}/t\mathbf{Z}$,

if $E(p, X) = E(p', X) \pmod q$, then $p = p' \pmod t$

where

$$E(p, X) = A_1((\dots((A_k((M[(A_k^{-1}((\dots((A_1^{-1}X)Z_1))\dots))Z_k])Z_k^{-1}))\dots))Z_1^{-1}) \pmod q,$$

$$M := R_1(\dots(R_r(u\mathbf{1} + vB + wG)R_r^{-1})\dots)R_1^{-1} \pmod q,$$

$$f = y + z \pmod q,$$

$$y = pks \pmod q = u + 2b_0v + (g_0 - b_0)w \pmod q,$$

$$z = dht \pmod q = u + (g_0 + b_0)w \pmod q,$$

$$E(p', X) = A_1((\dots((A_k((M'[(A_k^{-1}((\dots((A_1^{-1}X)Z_1))\dots))Z_k])Z_k^{-1}))\dots))Z_1^{-1}) \pmod q,$$

$$M' := R_1(\dots(R_r(u'\mathbf{1} + v'B + w'G)R_r^{-1})\dots)R_1^{-1} \pmod q,$$

$$f' = y' + z' \pmod{q},$$

$$y' = p'ks \pmod{q} = u' + 2b_0v' + (g_0 - b_0)w' \pmod{q},$$

$$z' = d'ht \pmod{q} = u' + (g_0 + b_0)w' \pmod{q}.$$

That is, if $p \neq p' \pmod{t}$, then $E(p, X) \neq E(p', X) \pmod{q}$.

(Proof)

If $E(p, X) = E(p', X) \pmod{q}$, then

$$\begin{aligned} D_1(E(p, (D_2(X))) &= D_1(E(p', (D_2(X))) \pmod{q} \\ MX &= M'X \pmod{q} \end{aligned}$$

We substitute $\mathbf{1}$ to X in above expression, we obtain

$$M = M' \pmod{q}.$$

$$\begin{aligned} R_1(\dots(R_r(u\mathbf{1} + vB + wG)R_r^{-1})\dots)R_1^{-1} \\ = R_1(\dots(R_r(u'\mathbf{1} + v'B + w'G)R_r^{-1})\dots)R_1^{-1} \pmod{q} \\ u\mathbf{1} + vB + wG = u'\mathbf{1} + v'B + w'G \pmod{q}. \end{aligned}$$

Then we have from $[u\mathbf{1} + vB + wG]_i$ and $[u'\mathbf{1} + v'B + w'G]_i$ ($i=0,1$)

$$u + v b_0 + w g_0 = u' + v' b_0 + w' g_0 \pmod{q},$$

and

$$(v - w)b_1 = (v' - w')b_1 \pmod{q}.$$

$$v - w = v' - w' \pmod{q}, \text{ (from } \text{GCD}(b_1, q) = 1)$$

$$u + v b_0 + w g_0 + (v - w) b_0 = u' + v' b_0 + w' g_0 + (v' - w') b_0 \pmod{q},$$

$$u + 2b_0v + (g_0 - b_0)w = u' + 2b_0v' + (g_0 - b_0)w' \pmod{q},$$

$$y = y' \pmod{q},$$

$$y \pmod{t} = p = y' \pmod{t} = p' \qquad \text{q.e.d.}$$

Next it is shown that the encrypting function $E(p, X)$ has the property of fully homomorphism.

We simply express the encrypting function such that

$$\begin{aligned} A_1(\dots((A_k((M[(A_k^{-1}((\dots((A_1^{-1}X)Z_1))\dots))Z_k])Z_k^{-1}))\dots))Z_1^{-1}) \pmod{q} \\ = A((M[(A^{-1}X)Z])Z^{-1}) \pmod{q}. \end{aligned}$$

§3.6 Addition scheme on ciphertexts

Here we consider the additive operation on the ciphertexts.

Let $M_1 := R[u_1\mathbf{1} + v_1B + w_1G]R^{-1}$, $M_2 := R[u_2\mathbf{1} + v_2B + w_2G]R^{-1} \in O$ be medium texts.

Let $f_1 = 2u_1 + 2b_0v_1 + 2g_0w_1 \pmod q \in \mathbf{Z}/q\mathbf{Z}$, $f_2 = 2u_2 + 2b_0v_2 + 2g_0w_2 \pmod q \in \mathbf{Z}/q\mathbf{Z}$.

Let $p_1 = f_1 \pmod t$, $p_2 = f_2 \pmod t$ be plaintexts to be encrypted.

Let $C_1(X) = E(p_1, X)$ and $C_2(X) = E(p_2, X)$ be the ciphertexts corresponding to the plaintexts p_1 and p_2 .

$$\begin{aligned}
C_1(X) + C_2(X) \pmod q &= E(p_1, X) + E(p_2, X) \pmod q \\
&= A((M_1[(A^{-1}X)\mathbf{Z}])\mathbf{Z}^{-1}) + A((M_2[(A^{-1}X)\mathbf{Z}])\mathbf{Z}^{-1}) \pmod q \\
&= A((M_1 + M_2)[(A^{-1}X)\mathbf{Z}])\mathbf{Z}^{-1} \pmod q \\
&= A((R(u_1\mathbf{1} + v_1B + w_1G + u_2\mathbf{1} + v_2B + w_2G)R^{-1})[(A^{-1}X)\mathbf{Z}])\mathbf{Z}^{-1} \pmod q \\
&= A((R((u_1 + u_2)\mathbf{1} + (v_1 + v_2)B + (w_1 + w_2)G)R^{-1})[(A^{-1}X)\mathbf{Z}])\mathbf{Z}^{-1} \pmod q. \\
f_{1+2} &= 2(u_1 + u_2) + 2b_0(v_1 + v_2) + 2g_0(w_1 + w_2) = f_1 + f_2 \pmod q \in \mathbf{Z}/q\mathbf{Z},
\end{aligned}$$

Then we have

$$E(p_1, X) + E(p_2, X) \pmod q = E(p_1 + p_2, X) \pmod q.$$

§3.7 Multiplication scheme on ciphertexts

Here we consider the multiplicative operation on the ciphertexts. Let $C_1(X) = E(p_1, X)$ and $C_2(X) = E(p_2, X)$ be the ciphertexts corresponding to the plaintexts p_1 and p_2 .

$$C_1(C_2(X)) \pmod q = E(p_1, E(p_2, X)) \pmod q$$

$$\begin{aligned}
&= A_1(\dots((A_k((M_1[(A_k^{-1}(\dots((A_1^{-1}\{A_1(\dots((A_k((M_2[(A_k^{-1}(\dots((A_1^{-1}X)Z_1))\dots))Z_k])Z_k^{-1}))\dots))Z_1^{-1})\dots))Z_1^{-1})\dots))Z_k^{-1})\dots))Z_1^{-1}) \pmod q \\
&= A_1(\dots((A_k((M_1[M_2[(A_k^{-1}(\dots((A_1^{-1}X)Z_1))\dots))Z_k])Z_k^{-1}))\dots))Z_1^{-1}) \pmod q \\
&= A_1(\dots((A_k(M_1(M_2[(A_k^{-1}(\dots((A_1^{-1}X)Z_1))\dots))Z_k])Z_k^{-1}))\dots))Z_1^{-1}) \pmod q. \\
&= A((M_1(M_2[(A^{-1}X)\mathbf{Z}]))\mathbf{Z}^{-1}) \pmod q. \tag{43}
\end{aligned}$$

Here we show the operation on B and G beforehand.

$$\begin{aligned}
&A((RBR^{-1})([RGR^{-1}][(A^{-1}X)\mathbf{Z}]))\mathbf{Z}^{-1} \pmod q \\
&= A((RBR^{-1})([R((b_0 + g_0)\mathbf{1} - B)R^{-1}][(A^{-1}X)\mathbf{Z}]))\mathbf{Z}^{-1} \pmod q \\
&= A((RBR^{-1})([R((b_0 + g_0)\mathbf{1})R^{-1}][(A^{-1}X)\mathbf{Z}]))\mathbf{Z}^{-1} - A((RBR^{-1})([RBR^{-1}][(A^{-1}X)\mathbf{Z}]))\mathbf{Z}^{-1} \pmod q
\end{aligned}$$

$$\begin{aligned}
&=(b_0+g_0) A(([RBR^{-1}]([A^{-1}XZ]))Z^{-1})-A(([RB^2R^{-1}][A^{-1}XZ]))Z^{-1}) \bmod q \\
&=(b_0+g_0) A(([RBR^{-1}]([A^{-1}XZ]))Z^{-1})- 2b_0 A(([RBR^{-1}][A^{-1}XZ]))Z^{-1}) \bmod q \\
&=(-b_0+g_0) A(([RBR^{-1}]([A^{-1}XZ]))Z^{-1}) \bmod q \\
&=A(([R(-b_0+g_0) BR^{-1}]([A^{-1}XZ]))Z^{-1}) \bmod q.
\end{aligned}$$

As $BG=(-b_0+g_0) B \bmod q$,

$$A(([RBR^{-1}][R GR^{-1}][A^{-1}XZ]))Z^{-1})=A(([R(BG)R^{-1}]([A^{-1}XZ]))Z^{-1}) \bmod q.$$

In the same manner we have

$$A(([RGR^{-1}][R BR^{-1}][A^{-1}XZ]))Z^{-1})=A(([R(GB)R^{-1}]([A^{-1}XZ]))Z^{-1}) \bmod q.$$

Substituting $R(u_1\mathbf{1}+v_1B+w_1G)R^{-1}$, $R(u_2\mathbf{1}+v_2B+w_2G)R^{-1}$ to M_1 , M_2 in (43) respectively, we have

$$\begin{aligned}
&C_1(C_2(X)) \bmod q =E(p_1, E(p_2, X)) \bmod q=A((M_1(M_2[A^{-1}XZ]))Z^{-1}) \bmod q. \\
&=A(([R(u_1\mathbf{1}+v_1B+w_1G)R^{-1}][R(u_2\mathbf{1}+v_2B+w_2G)R^{-1}][A^{-1}XZ]))Z^{-1}) \bmod q, \\
&=A(([R(u_1\mathbf{1})R^{-1}][R(u_2\mathbf{1}+v_2B+w_2G)R^{-1}][A^{-1}XZ]))Z^{-1}) \bmod q. \\
&+ A(([R(v_1B)R^{-1}][R(u_2\mathbf{1}+v_2B+w_2G)R^{-1}][A^{-1}XZ]))Z^{-1}) \bmod q \\
&+ A(([R(w_1G)R^{-1}][R(u_2\mathbf{1}+v_2B+w_2G)R^{-1}][A^{-1}XZ]))Z^{-1}) \bmod q. \\
&=A(([R(u_1\mathbf{1})R^{-1}][R(u_2\mathbf{1})R^{-1}][A^{-1}XZ]))Z^{-1}) \bmod q. \\
&+ A(([R(u_1\mathbf{1})R^{-1}][R(v_2B)R^{-1}][A^{-1}XZ]))Z^{-1}) \bmod q \\
&+ A(([R(u_1\mathbf{1})R^{-1}][R(w_2G)R^{-1}][A^{-1}XZ]))Z^{-1}) \bmod q \\
&+ A(([R(v_1B)R^{-1}][R(u_2\mathbf{1})R^{-1}][A^{-1}XZ]))Z^{-1}) \bmod q \\
&+ A(([R(v_1B)R^{-1}][R(v_2B)R^{-1}][A^{-1}XZ]))Z^{-1}) \bmod q \\
&+ A(([R(v_1B)R^{-1}][R(w_2G)R^{-1}][A^{-1}XZ]))Z^{-1}) \bmod q \\
&+ A(([R(w_1G)R^{-1}][R(u_2\mathbf{1})R^{-1}][A^{-1}XZ]))Z^{-1}) \bmod q \\
&+ A(([R(w_1G)R^{-1}][R(v_2B)R^{-1}][A^{-1}XZ]))Z^{-1}) \bmod q \\
&+ A(([R(w_1G)R^{-1}][R(w_2G)R^{-1}][A^{-1}XZ]))Z^{-1}) \bmod q \\
&=A(([R(u_1u_2\mathbf{1}+u_1v_2B+u_1w_2G+v_1u_2B+v_1v_2BB+v_1w_2BG \\
&+w_1u_2G+w_1v_2GB+w_1w_2GG)R^{-1}][A^{-1}XZ]))Z^{-1}) \bmod q
\end{aligned}$$

$$\begin{aligned}
&= A(([\mathbf{R}(u_1\mathbf{1}+v_1B+w_1G)\mathbf{R}^{-1})(\mathbf{R}(u_2\mathbf{1}+v_2B+w_2G)\mathbf{R}^{-1})][(\mathbf{A}^{-1}X\mathbf{Z})])\mathbf{Z}^{-1}) \bmod q \\
&= A((M_1M_2)[(\mathbf{A}^{-1}X\mathbf{Z})])\mathbf{Z}^{-1}) \bmod q.
\end{aligned}$$

Then we have

$$\begin{aligned}
&C_1(C_2(X)) \bmod q = E(p_1, E(p_2, X)) \bmod q \\
&= A(([\mathbf{R}(u_1u_2\mathbf{1}+u_1v_2B+u_1w_2G+v_1u_2B+v_1v_2BB+v_1w_2BG \\
&\quad +w_1u_2G+w_1v_2GB+w_1w_2GG)\mathbf{R}^{-1}][(\mathbf{A}^{-1}X\mathbf{Z})])\mathbf{Z}^{-1}) \bmod q \\
&= A(([\mathbf{R}(u_1u_2\mathbf{1}-(g_0^2-b_0^2)w_1w_2\mathbf{1}+(u_1v_2+v_1u_2+2b_0v_1v_2)B+(v_1w_2+w_1v_2)(-b_0+g_0)B \\
&\quad +(u_1w_2+w_1u_2+2g_0w_1w_2)G)\mathbf{R}^{-1}][(\mathbf{A}^{-1}X\mathbf{Z})])\mathbf{Z}^{-1}) \bmod q \\
&= A(([\mathbf{R}((u_1u_2-(g_0^2-b_0^2)w_1w_2)\mathbf{1}+(u_1v_2+v_1u_2+2b_0v_1v_2+(v_1w_2+w_1v_2)(-b_0+g_0))B \\
&\quad +(u_1w_2+w_1u_2+2g_0w_1w_2)G)\mathbf{R}^{-1}][(\mathbf{A}^{-1}X\mathbf{Z})])\mathbf{Z}^{-1}) \bmod q \\
&\text{Let } (m_0^*, m_1^*, \dots, m_7^*) := R_r^{-1}(\dots(R_1^{-1}([M_1M_2]R_1)\dots)R_r \bmod q \\
&= (u_1u_2-(g_0^2-b_0^2)w_1w_2)\mathbf{1}+(u_1v_2+v_1u_2+2b_0v_1v_2+(v_1w_2+w_1v_2)(-b_0+g_0))B \\
&\quad +(u_1w_2+w_1u_2+2g_0w_1w_2)G \bmod q.
\end{aligned}$$

We have the plaintext p_{12} of $C_1(C_2(X))$ as follows.

As $y_1z_2 + y_2z_1 \bmod q = 0$,

$$\begin{aligned}
&f_{12} := 2m_0^* \bmod q, \\
&= 2m_0^* + y_1z_2 + y_2z_1 \bmod q \\
&= 2(u_1u_2-(g_0^2-b_0^2)w_1w_2) + 2(u_1v_2+v_1u_2+2b_0v_1v_2+(-b_0+g_0)(v_1w_2+w_1v_2))b_0 \\
&\quad + 2(u_1w_2+w_1u_2+2g_0w_1w_2)g_0 \\
&\quad + (u_1+2b_0v_1+(g_0-b_0)w_1)(u_2+(g_0+b_0)w_2) + (u_2+2b_0v_2+(g_0-b_0)w_2)(u_1+(g_0+b_0)w_1) \bmod q \\
&= 4u_1u_2 + 4b_0(u_1v_2+v_1u_2) + (4b_0^2)v_1v_2 + (2b_0(-b_0+g_0)+2b_0(g_0+b_0))(v_1w_2+w_1v_2) \\
&\quad + (2g_0+2g_0)(u_1w_2+w_1u_2) + (4g_0^2-2(g_0^2-b_0^2)+2(g_0^2-b_0^2))w_1w_2 \bmod q \\
&= 4u_1u_2 + 4b_0(u_1v_2+v_1u_2) + (4b_0^2)v_1v_2 + 4b_0g_0(v_1w_2+w_1v_2) + 4g_0(u_1w_2+w_1u_2) \\
&\quad + 4g_0^2w_1w_2 \bmod q \\
&= (2u_1+2b_0v_1+2g_0w_1)(2u_2+2b_0v_2+2g_0w_2) \\
&= f_1f_2 \bmod q \in \mathbf{Z}/q\mathbf{Z}.
\end{aligned}$$

$$p_{12} := f_{12} \bmod t = (f_1 \bmod t)(f_2 \bmod t) = p_1 p_2 \bmod t.$$

Then we have

$$C_1(C_2(X)) \bmod q = E(p_1, E(p_2, X)) \bmod q = E(p_1 p_2, X) \bmod q. \quad (44)$$

It has been shown that this scheme has the multiplicative homomorphism.

§3.8 Fully homomorphism of proposed fully homomorphic encryption

(Fully homomorphic encryption). Proposed fully homomorphic encryption $= (\mathbf{KeyGen}; \mathbf{Enc}; \mathbf{Dec}; \mathbf{Eval})$ is fully homomorphic because it satisfies the following properties:

1. Homomorphism: Let $CR = \{CR_\lambda\}_{\lambda \in \mathbb{N}}$ be the set of all polynomial sized arithmetic circuits. On input $\mathbf{sk} \leftarrow \mathbf{KeyGen}(1^\lambda)$, $\forall \text{ckt} \in CR_\lambda$, $\forall (p_1, \dots, p_n) \in (\mathbb{Z}/t\mathbb{Z})^n$ where $n = n(\lambda)$, $\forall (C_1, \dots, C_n)$ where $C_i \leftarrow (E(p_i, X))$, $(i=1, \dots, n)$, we have $D(\mathbf{sk}; \mathbf{Eval}(\text{ckt}; C_1, \dots, C_n)) = \text{ckt}(p_1, \dots, p_n)$.

Then it holds that:

$$\Pr[D(\mathbf{sk}; \mathbf{Eval}(\text{ckt}; C_1, \dots, C_n)) \neq \text{ckt}(p_1, \dots, p_n)] = \text{negl}(\lambda).$$

2. Compactness: As the output length of **Eval** is at most $k \log_2 q = k\lambda$ where k is a positive integer, there exists a polynomial $\mu = \mu(\lambda)$ such that the output length of **Eval** is at most μ bits long regardless of the input circuit ckt and the number of its inputs.

§4. Analysis of proposed scheme

Here we analyze the proposed fully homomorphic encryption scheme.

§4.1 Computing parameter “y” from coefficients of ciphertext $E(p, X)$ to be published

Ciphertext $E(p, X)$ is published by cloud data centre as follows.

$$\begin{aligned} E(p, X) &= A((M[(A^{-1}X)Z])Z^{-1}) \bmod q \in O[X] \\ &= (e_{00}x_0 + e_{01}x_1 + \dots + e_{07}x_7, \\ &\quad e_{10}x_0 + e_{11}x_1 + \dots + e_{17}x_7, \\ &\quad \dots \quad \dots \\ &\quad e_{70}x_0 + e_{71}x_1 + \dots + e_{77}x_7) \bmod q, \end{aligned}$$

$$= \{e_{jt}\}(j,t=0,\dots,7)$$

with $e_{jt} \in \mathbf{Z}/q\mathbf{Z}$ ($j,t=0,\dots,7$) which is published in cloud data centre, where

$$M = \mathbf{R}(u\mathbf{1} + v\mathbf{B} + w\mathbf{G})\mathbf{R}^{-1} \bmod q,$$

$$f = y + z \bmod q,$$

$$y = pks \bmod q = u + 2b_0v + (g_0 - b_0)w \bmod q,$$

$$z = dth \bmod q = u + (g_0 + b_0)w \bmod q.$$

The user who knows the secret key $\mathbf{ks} = (\mathbf{A}, \mathbf{Z}, \mathbf{R}, B, G, s, t)$ can calculate the plaintext $p \in \mathbf{Z}/t\mathbf{Z}$ from the cipher text $E(p, X) \in O[X]$ as follows.

$$M' = (m'_0, m'_1, \dots, m'_7) = R_r^{-1}(\dots(R_1^{-1}(M R_1)\dots)R_r) = (u\mathbf{1} + v\mathbf{B} + w\mathbf{G}) \bmod q$$

$$= u\mathbf{1} + v(b_0, b_1, \dots, b_7) + w(g_0, -b_1, \dots, -b_7) \bmod q.$$

$$m'_0 + m'_1(b_0/b_1) = u + 2b_0v + (g_0 - b_0)w \bmod q = y = pks \bmod q,$$

$$m'_0 - m'_1(b_0/b_1) = u + (g_0 + b_0)w \bmod q = z = dth \bmod q.$$

$$2m'_0 = 2u + 2vb_0 + 2wg_0 \bmod q = y + z = f \in \mathbf{Z}/q\mathbf{Z}.$$

$A_i, Z_i, R_j \in O$ are the secret keys to be selected such that A_i^{-1}, Z_i^{-1} and R_j^{-1} exist ($i=1, \dots, k; j=1, \dots, r$).

[Theorem 8]

If there exists the PPT algorithm AL1 for obtaining the parameter “y” from coefficients of any $E(p, X)$, $e_{jt} \in \mathbf{Z}/q\mathbf{Z}$ ($j,t=0,\dots,7$), there exists the PPT algorithm that factors modulus q .

(Proof:)

As $y = pks \bmod q$, we have

$$\text{GCD}(y, q) = \text{GCD}(pks \bmod q, st) = s. \quad \text{q.e.d.}$$

§4.2 Computing plaintext p and A_i, Z_i ($i=1, \dots, k$) from coefficients of ciphertext $E(p, X)$ to be published

Ciphertext $E(p_d, X)$ ($d=1, 2, 3$) is published by cloud data centre as follows.

$$E(p_d, X) = \mathbf{A}((M_d[(\mathbf{A}^{-1}X)\mathbf{Z}])\mathbf{Z}^{-1}) \bmod q$$

$$\begin{aligned}
&= \mathbf{A}((\mathbf{R}[u_d \mathbf{1} + v_d \mathbf{B} + w_d \mathbf{G}]\mathbf{R}^{-1})[(\mathbf{A}^{-1} \mathbf{X})\mathbf{Z}])\mathbf{Z}^{-1} \bmod q \in \mathcal{O}[\mathbf{X}], \\
&= (e_{d00}x_0 + e_{d01}x_1 + \dots + e_{d07}x_7, \\
&\quad e_{d10}x_0 + e_{d11}x_1 + \dots + e_{d17}x_7, \\
&\quad \dots \quad \dots \\
&\quad e_{d70}x_0 + e_{d71}x_1 + \dots + e_{d77}x_7) \bmod q, \\
&= \{e_{djk}\}(j, r=0, \dots, 7; d=1, 2, 3)
\end{aligned}$$

with $e_{djt} \in \mathbf{Z}/q\mathbf{Z}$ ($j, t=0, \dots, 7; d=1, 2, 3$) which is published, where $B, G, A_i, Z_i, R_j \in \mathcal{O}$ to be selected such that A_i^{-1}, Z_i^{-1} and R_j^{-1} exist ($i=1, \dots, k; j=1, \dots, r$) are the secret keys.

We try to find plaintext p from coefficients of $E(p_d, X)$, $e_{djt} \in \mathbf{Z}/q\mathbf{Z}$ ($j, t=0, \dots, 7; d=1, 2, \dots$).

In case that $k=8, r=8$ and $d=3$ the number of unknown variables ($p_d, u_d, v_d, w_d, b_0, g_0, b_1, \dots, b_7, A_i, Z_i, R_j$ ($i, j=1, \dots, 8; d=1, 2, 3$)) is $213 (=4*3+9+3*8*8)$, the number of equations is $192 (=64*3)$ such that

$$\begin{aligned}
&F_{100}(M, A_i, Z_i, R_i) = e_{100} \bmod q, \\
&F_{101}(M, A_i, Z_i, R_i) = e_{101} \bmod q, \\
&\quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \\
&F_{107}(M, A_i, Z_i, R_i) = e_{107} \bmod q, \\
&\quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \\
&F_{377}(M, A_i, Z_i, R_i) = e_{377} \bmod q,
\end{aligned} \tag{45}$$

where F_{100}, \dots, F_{377} are the $50 (=8*2*3+2)^{\text{th}}$ algebraic multivariate equations.

Then the complexity G_{reb} required for solving above simultaneous equations by using Gröbner basis is given [6] such as

$$G_{reb} > G_{reb}' = (191 + d_{reg} C_{dreg})^w = (4895 C_{191})^w = 2^{2769} \gg 2^{80},$$

where G_{reb}' is the complexity required for solving 192 simultaneous algebraic equations with 191 variables by using Gröbner basis, where $w=2.39$, and

$$d_{reg} = 4704 (=192*(50-1)/2 - 0\sqrt{(192*(50^2-1)/6)}).$$

The complexity G_{reb} required for solving above simultaneous equations by using

Gröbner basis is enough large for secure.

§4.3 Computing plaintext p and d_{ijk} ($i,j,k=0,\dots,7$)

We try to computing plaintext p_r ($r=0,\dots,7$) and d_{ijk} ($i,j,k=0,\dots,7$) from coefficients of ciphertext $E(p_r, X)$ to be published.

At first let $Enc(Y, X) \in O[X, Y]$ be the enciphering function such as

$$\begin{aligned}
 Enc(Y, X) &:= A((Y[(A^{-1}X)Z])Z^{-1}) \bmod q \in O[X, Y], \\
 &= (d_{000}x_0y_0 + d_{001}x_0y_1 + \dots + d_{077}x_7y_7, \\
 &\quad d_{100}x_0y_0 + d_{101}x_0y_1 + \dots + d_{177}x_7y_7, \\
 &\quad \dots \quad \dots \\
 &\quad d_{700}x_0y_0 + d_{701}x_0y_1 + \dots + d_{777}x_7y_7) \bmod q, \\
 &= \{d_{ijk}\} (i, j, k = 0, \dots, 7)
 \end{aligned} \tag{46}$$

with $d_{ijk} \in \mathbf{Z}/q\mathbf{Z}$ ($i, j, k = 0, \dots, 7$).

Next we substitute M_r to Y where

$$M_r = (m_{r0}, m_{r1}, \dots, m_{r7}) = \mathbf{R} [p_r \mathbf{1} + u_r \mathbf{B} + v_r \mathbf{G}] \mathbf{R}^{-1} \bmod q \in O. \tag{47}$$

We have

$$\begin{aligned}
 E(p_r, X) &= A((M_r[(A^{-1}X)Z])Z^{-1}) \bmod q \in O[X], \\
 &= (d_{000}x_0m_{r0} + d_{001}x_0m_{r1} + \dots + d_{077}x_7m_{r7}, \\
 &\quad d_{100}x_0m_{r0} + d_{101}x_0m_{r1} + \dots + d_{177}x_7m_{r7}, \\
 &\quad \dots \quad \dots \\
 &\quad d_{700}x_0m_{r0} + d_{701}x_0m_{r1} + \dots + d_{777}x_7m_{r7}) \bmod q.
 \end{aligned} \tag{48}$$

Then we obtain 64 equations from (38) and (48) as follows.

$$\left. \begin{aligned}
 d_{000}m_{i0} + d_{001}m_{i1} + \dots + d_{007}m_{i7} &= e_{00} \bmod q \\
 d_{010}m_{i0} + d_{011}m_{i1} + \dots + d_{017}m_{i7} &= e_{01} \bmod q \\
 &\dots \quad \dots \\
 d_{070}m_{i0} + d_{071}m_{i1} + \dots + d_{077}m_{i7} &= e_{07} \bmod q
 \end{aligned} \right\} \tag{49a}$$

$$\left. \begin{aligned}
 d_{100}m_{i0} + d_{101}m_{i1} + \dots + d_{107}m_{i7} &= e_{10} \bmod q
 \end{aligned} \right\}$$

$$d_{110}m_{i0}+d_{111}m_{i1}+ \dots +d_{117}m_{i7}=e_{11} \bmod q \quad (49b)$$

.....

$$d_{170}m_{i0}+d_{171}m_{i1}+ \dots +d_{177}m_{i7}=e_{17} \bmod q$$

.....

.....

$$d_{700}m_{i0}+d_{701}m_{i1}+ \dots +d_{707}m_{i7}=e_{70} \bmod q$$

$$d_{710}m_{i0}+d_{711}m_{i1}+ \dots +d_{717}m_{i7}=e_{71} \bmod q$$

.....

$$d_{770}m_{i0}+d_{771}m_{i1}+ \dots +d_{777}m_{i7}=e_{77} \bmod q$$

}
}
}

(49c)

For M_0, \dots, M_7 we obtain the same equations, the number of which is 512.

We also obtain the 8 equations such as

$$|E(p_i, \mathbf{1})|^2 = |M_i|^2 = m_{i0}^2 + m_{i1}^2 + \dots + m_{i7}^2 = 0 \bmod q, (i=0, \dots, 7). \quad (50)$$

The number of unknown variables M_i and d_{ijk} ($i, j, k=0, \dots, 7$) is $576 (=512+64)$. The number of equations is $520 (=512+8)$. Then the complexity G_{reb} required for solving above simultaneous quadratic algebraic equations by using Gröbner basis is given such as

$$G_{reb} > G_{reb}' = (520 + d_{reg} C_{dreg})^w = (763 C_{243})^w = 2^{1634} \gg 2^{80},$$

where G_{reb}' is the complexity required for solving 520 simultaneous quadratic algebraic equations with 520 variables by using Gröbner basis,

where $w=2.39$, and

$$d_{reg} = 243 (=520*(2-1)/2 - 1\sqrt{(520*(4-1)/6}).$$

It is thought to be difficult computationally to solve the above simultaneous algebraic equations by using Gröbner basis.

§4.4 Attack by using the ciphertexts of p and $-p$

I show that we can not easily distinguish the ciphertexts of p and $-p$. We try to attack by using “ p and $-p$ attack”.

Let $M := \mathbf{R}(p\mathbf{1} + u\mathbf{B} + v\mathbf{G})\mathbf{R}^{-1} \in \mathcal{O}$ be a medium text, $p \in \mathbf{Z}/t\mathbf{Z}$ be a plaintext, $u_+, v_+, w_+, d_+ \in \mathbf{Z}/q\mathbf{Z}$ be parameters, and $E(p, X)$ be the ciphertext of p .

By using simple style expression of $E(p, X)$ we have

$$C_+(X) := E(p, X) = A((M[(A^{-1}X)Z])Z^{-1}) \bmod q \in O[X],$$

$$f_+ := y_+ + z_+ \bmod q,$$

where

$$y_+ := psk \bmod q = u_+ + 2b_0v_+ + (g_0 - b_0)w_+ \bmod q \in \mathbf{Z}/q\mathbf{Z},$$

$$z_+ := d_+th \bmod q = u_+ + (g_0 + b_0)w_+ \bmod q \in \mathbf{Z}/q\mathbf{Z}.$$

Let $E(-p, X)$ be the ciphertext of $-p \bmod q$.

$$C_-(X) := E(-p, X) = A(((M_-.[(A^{-1}X)Z])Z^{-1}) \bmod q \in O[X],$$

$$M_- := \mathbf{R}(u_- \mathbf{1} + v_- B + w_- G) \mathbf{R}^{-1} \bmod q \in O,$$

where

$$f_- := y_- + z_- \bmod q,$$

$$y_- := -psk \bmod q = u_- + 2b_0v_- + (g_0 - b_0)w_- \bmod q \in \mathbf{Z}/q\mathbf{Z},$$

$$z_- := d_-th \bmod q = u_- + (g_0 + b_0)w_- \bmod q \in \mathbf{Z}/q\mathbf{Z}.$$

$$u_-, v_-, w_-, d_- \in \mathbf{Z}/q\mathbf{Z}.$$

Then we have

$$y_+ + y_- \bmod q = psk - psk \bmod q = 0 \bmod q,$$

$$y_+ + y_- \bmod q = u_+ + u_- + 2b_0(v_+ + v_-) + (g_0 - b_0)(w_+ + w_-) \bmod q = 0 \bmod q,$$

$$u_+ + u_- = -2b_0(v_+ + v_-) - (g_0 - b_0)(w_+ + w_-) \bmod q.$$

[Theorem 9]

If there exists the PPT algorithm AL2 for generating the ciphertext $C_-(X) = E(-p, X)$ from any ciphertext $C_+(X) = E(p, X)$, there exists the PPT algorithm that factors modulus q .

(Proof:)

By substituting $\mathbf{1}$ to X of $C_+(X) + C_-(X)$, we have

$$C_+(\mathbf{1}) + C_-(\mathbf{1}) = A(([M_+ + M_-] [(A^{-1}\mathbf{1})Z])Z^{-1}) \bmod q$$

$$= A([\mathbf{R}(u_+ \mathbf{1} + v_+ B + w_+ G + u_- \mathbf{1} + v_- B + w_- G) \mathbf{R}^{-1}] [(A^{-1}\mathbf{1})Z])Z^{-1}) \bmod q$$

$$= A([\mathbf{R}((u_+ + u_-)\mathbf{1} + (v_+ + v_-)B + (w_+ + w_-)G) \mathbf{R}^{-1}] [(A^{-1}\mathbf{1})Z])Z^{-1}) \bmod q$$

$$= A([\mathbf{R}((v_+ + v_-)(B - 2b_0\mathbf{1}) + (w_+ + w_-)(G - (g_0 - b_0)\mathbf{1})) \mathbf{R}^{-1}] [(A^{-1}\mathbf{1})Z])Z^{-1}) \bmod q$$

$$= (v_+ + v_- - w_+ - w_-)A([RHR^{-1}] [(A^{-1}\mathbf{1})\mathbf{Z}])\mathbf{Z}^{-1}) \bmod q$$

where

$$H = (-b_0, b_1, \dots, b_7) \in O,$$

$$|H|^2 = b_0^2 + b_1^2 + \dots + b_7^2 = 0 \bmod q.$$

$$\text{As } 2b_0(v_+ + v_- - w_+ - w_-) = y_+ + y_- - z_+ - z_- \bmod q$$

$$= psk - psk - d_+ht - d_-ht \bmod q$$

$$= -(d_+ + d_-)ht \bmod q,$$

We have

$$C^* := C_+(\mathbf{1}) + C_-(\mathbf{1}) = -(d_+ + d_-)ht(2b_0)^{-1}A([RHR^{-1}] [(A^{-1}\mathbf{1})\mathbf{Z}])\mathbf{Z}^{-1}) \bmod q.$$

Let $C^* = (c_0, \dots, c_7) \in O$. We have with overwhelming probability

$$\text{GCD}(c_i, q) = t, (i \in \{0, \dots, 7\}).$$

That is, we have factored composite number q . q.e.d.

Then it is said that the complexity required for attack by using “ p and $-p$ attack” is as large as the complexity required for factoring the composite number q . We can not easily distinguish the ciphertexts of p and $-p$.

§5. The size of the modulus q and the complexity for enciphering/

deciphering

We consider the size of the system parameter q . We select the size of q such that $O(q)$, the size of the composite number is as large as 2^{2000} . Then we need to select modulus $O(q) = 2^{2000}$ where $O(s) = 2^{1000}$ and $O(t) = 2^{1000}$.

- 1) In case of $k=8$, $O(q) = 2^{2000}$, the size of $e_{ij} \in \mathbf{Z}/q\mathbf{Z} (i, j = 0, \dots, 7)$ which are the coefficients of elements in $E(p, X) = A((M[(A^{-1}X)\mathbf{Z}])\mathbf{Z}^{-1}) \bmod q \in O[X]$ is $(64)(\log_2 q)$ bits = 128 kbits, and the size of system parameters q is 2 kbits.
- 2) In case of $k=8$, $O(q) = 2^{2000}$, the complexity G_{enc} to obtain $Enc(Y, X)$ required one time at the start of the system is $(15 \cdot 64 \cdot 8 + 15 \cdot 512 \cdot 8 + 16 \cdot 8 \cdot 8)(\log_2 q)^2 + K_{AZR} = 2^{39}$ bit-operations,

where

$K_{AZR}=24*16*(\log_2q)^2+24*(\log_2q)^3=2^{37.5}$ bit-operations is the complexity required for A_i^{-1} , Z_i^{-1} and R_i^{-1} ($i=0,\dots,7$).

- 3) In case of $k=8$, $r=8$, $O(q)=2^{2000}$, the complexity G_E to obtain $E(p,X)$ from $Enc(Y,X)$ required at every enciphering is $(64*2*8+O(10))(\log_2q)^2 = 2^{32}$ bit-operations.
- 4) The complexity G_D required for deciphering by using A_i^{-1} and Z_i^{-1} ($i=0,\dots,7$) is given as follows.

Let $C:=A_1((\dots((A_k((M[(A_k^{-1}((\dots((A_1^{-1}\mathbf{1})Z_1))\dots))Z_k])Z_k^{-1})))\dots))Z_1^{-1}) \bmod q$.

We have

$$(A_k((\dots((A_1^{-1}C)Z_1))Z_2))\dots)Z_k = M[(A_k^{-1}((\dots((A_1^{-1}\mathbf{1})Z_1))\dots))Z_k] \bmod q,$$

$$M=[(A_k((\dots((A_1^{-1}C)Z_1))Z_2))\dots)Z_k][(A_k^{-1}((\dots((A_1^{-1}\mathbf{1})Z_1))\dots))Z_k]^{-1} \bmod q.$$

$$=R_1(\dots(R_r(u\mathbf{1}+vB+wG)R_r^{-1})\dots)R_1^{-1}$$

$$(m_0', m_1', \dots, m_7') := (u\mathbf{1}+vB+wG) = R_r^{-1}(\dots(R_1^{-1}M R_1)\dots)R_r$$

$$2m_0' \bmod t=p.$$

Then the complexity G_D required for every deciphering is

$$(16*64+15*64+64)(\log_2s)^2+(1+1)*(\log_2s)^3+(16+2)*(\log_2s)^2 = 2^{32} \text{ bit-operations.}$$

- 5) The complexity required for addition operation on ciphertexts, $E(p_1, X)+E(p_2, X)$ has no multiplication.
- 6) The complexity G_M required for multiplication operation on ciphertexts, $E(p_1, E(p_2, X))$ is $(8*8*8)(\log_2q)^2 = 2^{31}$ bit-operations.

On the other hand the complexity required for enciphering and deciphering in RSA scheme is $O(2(\log n)^3) = 2^{34}$ bit-operations each where $O(n)=2^{2000}$.

Though our scheme requires memory space larger than RAS scheme, the complexity required to encipher and decipher is smaller than RSA scheme.

§6. Conclusion

We proposed the fully homomorphism encryption scheme based on the octonion ring over finite ring. It was shown that our scheme is immune from the Gröbner basis attacks by calculating the complexity to obtain the Gröbner basis for the multivariate

algebraic equations.

The proposed scheme does not require a “bootstrapping” process. We proved that if there exists the PPT algorithm that generates ciphertext of the plaintext $-p$ from the ciphertext of any plaintext p , there exists the PPT algorithm that factors the given composite number modulus q .

§7. Acknowledgments

In this paper we have proposed the scheme which we improve the encryption scheme described in chapter 4 of my work “Fully Homomorphic Encryption without bootstrapping” published in March, 2015 which was published by LAP LAMBERT Academic Publishing, Saarbrücken/Germany [1].

§8. BIBLIOGRAPHY

- [1] Masahiro, Y. (2015). Fully Homomorphic Encryption without bootstrapping. Saarbrücken/Germany: LAP LAMBERT Academic Publishing.
- [2] Shigeo Tsujii , Kohtaro Tadaki , Masahito Gotaishi ,Ryo Fujita ,and Masao Kasahara , "Proposal Integrated MPKC:PPS—STS Enhanced Perturbed Piece in Hand Method---," IEICE Tech. Rep.ISEC2009-27,SITE2009-19,ICSS2009-41(2009-07),July 2009.
- [3] S. Tsujii, K. Tadaki, and R. Fujita, "Piece in hand concept for enhancing the security of multivariate type public key cryptosystems: Public key without containing all the information of secret key," Cryptology ePrint Archive, Report 2004/366, 2004.
- [4] C.Wolf, and B. Preneel, "Taxonomy of public key schemes based on the problem of multivariate quadratic equations," Cryptology ePrint Archive, Report 2005/077, 2005, <http://eprint.iacr.org/>.
- [5] Shigeo Tsujii , Kohtaro Tadaki , Masahito Gotaishi ,Ryo Fujita ,and Masao Kasahara , "Proposal Integrated MPKC:PPS—STS Enhanced Perturbed Piece in Hand Method---," IEICE Tech. Rep.ISEC2009-27, SITE2009-19, ICSS2009-41(2009-07), July 2009.
- [6] M. Bardet, J. C. Faugere, and B. Salvy, "On the complexity of Gröbner basis computation of semi-regular overdetermined algebraic equations," Proceeding of the International Conference on Polynomial System Solving(ICPSS2004), pp.71-75, November 2004.
- [7] Craig Gentry. Fully Homomorphic Encryption Using Ideal Lattices.In the 41st ACM Symposium on Theory of Computing (STOC), 2009.
- [8] Craig Gentry, A Fully Homomorphic Encryption Scheme, 2009. Available at <http://crypto.stanford.edu/craig/craig-thesis.pdf> .
- [9] Marten van Dijk; Craig Gentry, Shai Halevi, and Vinod Vaikuntanathan (2009-12-11). "[Fully Homomorphic Encryption over the Integers](#)" (PDF). International Association for Cryptologic Research. Retrieved 2010-03-18.
- [10] Damien Stehle; Ron Steinfeld (2010-05-19). "Faster Fully Homomorphic Encryption" (PDF). International Association for Cryptologic Research. Retrieved 2010-09-15.
- [11] JS Coron, A Mandal, D Naccache, M Tibouchi," Fully homomorphic encryption over the integers with shorter public keys", Advances in Cryptology–CRYPTO 2011, 487-504.
- [12] Nuida and Kurosawa,"(Batch) Fully Homomorphic Encryption over Integers for Non-Binary Message Spaces", Cryptology ePrint Archive, Report 2014/777, 2014. <http://eprint.iacr.org/>.

- [13] John H. Conway, Derek A. Smith co-authored, translated by Syuuji Yamada, "On Quaternions and Octonions " Baifuukan Publication Center, Tokyo, .2006.
- [14] Masahiro Yagisawa," Fully Homomorphic Encryption with composite number modulus", Cryptology ePrint Archive, Report 2015/1040, 2015.
<http://eprint.iacr.org/>.
- [15] Masahiro Yagisawa," Fully Homomorphic Encryption Without bootstrapping", Cryptology ePrint Archive, Report 2015/474, 2015. <http://eprint.iacr.org/>.
- [16] Masahiro Yagisawa," Improved Fully Homomorphic Encryption with composite number modulus", Cryptology ePrint Archive, Report 2016/050, 2016.

Appendix A:**Octinv(A)** -----

```

S ← a02+a12+...+a72 mod q.
% S-1 mod q
q[1] ← q div S ;% integer part of q/S
r[1] ← q mod S ;% residue
k ← 1
q[0] ← q
r[0] ← S
while r[k] ≠ 0
    begin
        k ← k + 1
        q[k] ← r[k-2] div r[k-1]
        r[k] ← r[k-2] mod [rk-1]
    end
Q [k-1] ← (-1)*q[k-1]
L[ k-1] ← 1
i ← k-1
while i > 1
    begin
        Q[ i-1] ← (-1)*Q[ i ]*q[i-1] + L[ i ]
        L[ i-1 ] ← Q[ i ]
        i ← i-1
    end

invS ← Q[1] mod q
invA[0] ← a0*invS mod q
For i=1,...,7,
    invA[i] ← (-1)*ai*invS mod q
Return A-1 = (invA[0], invA[1],..., invA[7])

```

Appendix B:**Lemma 2**

$$A^{-1}(AB) = B$$

$$(BA)A^{-1} = B$$

(Proof:)

$$A^{-1} = (a_0/|A|^2 \bmod q, -a_1/|A|^2 \bmod q, \dots, -a_7/|A|^2 \bmod q).$$

$$AB \bmod q$$

$$\begin{aligned} &= (a_0b_0 - a_1b_1 - a_2b_2 - a_3b_3 - a_4b_4 - a_5b_5 - a_6b_6 - a_7b_7 \bmod q, \\ &\quad a_0b_1 + a_1b_0 + a_2b_4 + a_3b_7 - a_4b_2 + a_5b_6 - a_6b_5 - a_7b_3 \bmod q, \\ &\quad a_0b_2 - a_1b_4 + a_2b_0 + a_3b_5 + a_4b_1 - a_5b_3 + a_6b_7 - a_7b_6 \bmod q, \\ &\quad a_0b_3 - a_1b_7 - a_2b_5 + a_3b_0 + a_4b_6 + a_5b_2 - a_6b_4 + a_7b_1 \bmod q, \\ &\quad a_0b_4 + a_1b_2 - a_2b_1 - a_3b_6 + a_4b_0 + a_5b_7 + a_6b_3 - a_7b_5 \bmod q, \\ &\quad a_0b_5 - a_1b_6 + a_2b_3 - a_3b_2 - a_4b_7 + a_5b_0 + a_6b_1 + a_7b_4 \bmod q, \\ &\quad a_0b_6 + a_1b_5 - a_2b_7 + a_3b_4 - a_4b_3 - a_5b_1 + a_6b_0 + a_7b_2 \bmod q, \\ &\quad a_0b_7 + a_1b_3 + a_2b_6 - a_3b_1 + a_4b_5 - a_5b_4 - a_6b_2 + a_7b_0 \bmod q). \end{aligned}$$

$$[A^{-1}(AB)]_0$$

$$\begin{aligned} &= \{ a_0(a_0b_0 - a_1b_1 - a_2b_2 - a_3b_3 - a_4b_4 - a_5b_5 - a_6b_6 - a_7b_7) \\ &\quad + a_1(a_0b_1 + a_1b_0 + a_2b_4 + a_3b_7 - a_4b_2 + a_5b_6 - a_6b_5 - a_7b_3) \\ &\quad + a_2(a_0b_2 - a_1b_4 + a_2b_0 + a_3b_5 + a_4b_1 - a_5b_3 + a_6b_7 - a_7b_6) \\ &\quad + a_3(a_0b_3 - a_1b_7 - a_2b_5 + a_3b_0 + a_4b_6 + a_5b_2 - a_6b_4 + a_7b_1) \\ &\quad + a_4(a_0b_4 + a_1b_2 - a_2b_1 - a_3b_6 + a_4b_0 + a_5b_7 + a_6b_3 - a_7b_5) \\ &\quad + a_5(a_0b_5 - a_1b_6 + a_2b_3 - a_3b_2 - a_4b_7 + a_5b_0 + a_6b_1 + a_7b_4) \\ &\quad + a_6(a_0b_6 + a_1b_5 - a_2b_7 + a_3b_4 - a_4b_3 - a_5b_1 + a_6b_0 + a_7b_2) \\ &\quad + a_7(a_0b_7 + a_1b_3 + a_2b_6 - a_3b_1 + a_4b_5 - a_5b_4 - a_6b_2 + a_7b_0) \} / |A|^2 \bmod q \end{aligned}$$

$$= \{ (a_0^2 + a_1^2 + \dots + a_7^2) b_0 \} / |A|^2 = b_0 \bmod q$$

where $[M]_i$ denotes the i -th element of $M \in O$.

$$[A^{-1}(AB)]_1$$

$$\begin{aligned} &= \{ a_0(a_0b_1 + a_1b_0 + a_2b_4 + a_3b_7 - a_4b_2 + a_5b_6 - a_6b_5 - a_7b_3) \\ &\quad - a_1(a_0b_0 - a_1b_1 - a_2b_2 - a_3b_3 - a_4b_4 - a_5b_5 - a_6b_6 - a_7b_7) \\ &\quad - a_2(a_0b_4 + a_1b_2 - a_2b_1 - a_3b_6 + a_4b_0 + a_5b_7 + a_6b_3 - a_7b_5) \end{aligned}$$

$$\begin{aligned}
& -a_3(a_0b_7+a_1b_3+a_2b_6-a_3b_1+a_4b_5-a_5b_4-a_6b_2+a_7b_0) \\
& +a_4(a_0b_2-a_1b_4+a_2b_0+a_3b_5+a_4b_1-a_5b_3+a_6b_7-a_7b_6) \\
& - a_5(a_0b_6+a_1b_5-a_2b_7+a_3b_4-a_4b_3-a_5b_1+a_6b_0+a_7b_2) \\
& +a_6(a_0b_5-a_1b_6+a_2b_3-a_3b_2-a_4b_7+a_5b_0+a_6b_1+a_7b_4) \\
& +a_7(a_0b_3-a_1b_7-a_2b_5+a_3b_0+a_4b_6+a_5b_2-a_6b_4+a_7b_1) \} /|A|^2 \bmod q \\
& =\{(a_0^2+a_1^2+\dots+a_7^2) b_1\} /|A|^2=b_1 \bmod q.
\end{aligned}$$

Similarly we have

$$[A^{-1}(AB)]_i=b_i \bmod q \quad (i=2,3,\dots,7).$$

Then

$$A^{-1}(AB)=B \bmod q. \quad \text{q.e.d.}$$