



Title	Crop classification from Sentinel-2 derived vegetation indices using ensemble learning
Author(s)	Sonobe, Rei; Yamaya, Yuki; Tani, Hiroshi; Wang, Xiufeng; Kobayashi, Nobuyuki; Mochizuki, Kan-ichiro
Citation	The Journal of Applied Remote Sensing, 12(2), 026019 https://doi.org/10.1117/1.JRS.12.026019
Issue Date	2018-05-18
Doc URL	http://hdl.handle.net/2115/71029
Rights	© 2018 Society of Photo-Optical Instrumentation Engineers(SPIE). One print or electronic copy may be made for personal use only. Systematic reproduction and distribution, duplication of any material in this paper for a fee or for commercial purposes, or modification of the content of the paper are prohibited., Rei Sonobe, Yuki Yamaya, Hiroshi Tani, Xiufeng Wang, Nobuyuki Kobayashi, Kan-ichiro Mochizuki, " Crop classification from Sentinel-2 derived vegetation indices using ensemble learning, " The Journal of Applied Remote Sensing, 12(2), 026019, (2018); https://doi.org/10.1117/1.JRS.12.026019 .
Type	article (author version)
File Information	Manuscript_JARS-180133.pdf



[Instructions for use](#)

Crop classification from Sentinel-2 derived vegetation indices using ensemble learning

Rei Sonobe,^{a,*} Yuki Yamaya,^b Hiroshi Tani,^c Xiufeng Wang,^c Nobuyuki Kobayashi,^d Kan-ichiro Mochizuki^e

^aShizuoka University, Faculty of Agriculture, Shizuoka, Japan, 422-8529

^bHokkaido University, Graduate School of Agriculture, Sapporo, Japan, 060-5859

^cHokkaido University, Research Faculty of Agriculture, Sapporo, Japan, 060-5859

^dSmart Link Hokkaido, Iwamizawa, Japan, 068-0034

^ePASCO Corporation, Tokyo, Japan, 153-0043

Abstract. The identification and mapping of crops are important for estimating potential harvest as well as for agricultural field management. Optical remote sensing is one of the most attractive options because it offers vegetation indices and some data have been distributed free of charge. Especially, Sentinel-2A, which is equipped with a multispectral sensor (MSI) with blue, green, red and near-infrared-1 bands at 10 m; red edge 1 to 3, near-infrared-2 and shortwave infrared 1 and 2 at 20 m; and 3 atmospheric bands (Band 1, Band 9 and Band 10) at 60 m, offers some vegetation indices calculated to assess vegetation status. However, sufficient consideration has not been given to the potential of vegetation indices calculated from MSI data. Thus, 82 published indices were calculated and their importance were evaluated for classifying crop types. In this study, the two most common classification algorithms, random forests (RF) and support vector machine (SVM), were applied to conduct cropland classification from MSI data. Additionally, super learning was applied for more improvement, achieving overall accuracies of 90.2–92.2%. Of the two algorithms applied (RF and SVM), the accuracy of SVM was superior and 89.3–92.0% of overall accuracies were confirmed. Furthermore, stacking contributed to higher overall accuracies (90.2–92.2%) and significant differences were confirmed with the results of SVM and RF. Our results showed that vegetation indices had the greatest contributions in identifying specific crop types.

Keywords: crop, random forests, Sentinel-2, stacking, support vector machine, vegetation index.

*First Author, E-mail: reysnb@gmail.com

1 Introduction

From a land-planning perspective, cropland diversity is vital and crop cover maps provide information for estimating potential harvest and agricultural field management. To document field properties such as cultivated crops and locations, some local governments in Japan have been using manual methods¹. However, more efficient techniques are required to reduce the high expense of these methods. Thus, satellite data-based cropland mapping has gained attention. Some spectral indices, which are combinations of spectral measurements at different wavelengths, have been used to evaluate phenology or quantify biophysical parameters²⁻⁵. As a

result, they have also made crop maps more accurate in previous studies⁶ and the abilities of optical remote sensing data have been improved for monitoring agricultural fields. The opportunities to obtain optical remote sensing data have improved due to the Sentinel-2A satellite launch on June 23, 2015. Now, it is collecting multispectral data including 13 bands covering the visible, SWIR wavelength regions. Sentinel-2B, which possesses the same specifications, was launched on March 7, 2017 and creates greater opportunities for monitoring agricultural fields. Furthermore, various spectral indices can be extracted including indices based on shortwave infrared bands (SWIR), which are influenced by plant constituents such as pigments, leaf water contents and biochemicals^{7,8}. Furthermore, vegetation indices derived from reflectance data acquired from optical sensors have been widely used to assess variations in the physiological states and biophysical properties of vegetation⁹⁻¹¹. Specifically, the Normalized Difference Vegetation Index (NDVI)¹², Soil-Adjusted Vegetation Index (SAVI)¹³ and Enhanced Vegetation Index (EVI)¹⁴ have been used for monitoring vegetation systems or ecological responses to environmental change¹⁵. MSI data have been used for identifying crop types¹⁶⁻¹⁸, plastic-covered greenhouses¹⁹, water bodies²⁰ and some previous studies showed the potential of VIs calculated from MSI data. However, it is possible to calculate a vast number of VIs from MSI data and most of them have been ignored in the previous studies. In this study 82 published indices and original reflectance data sources were evaluated to classify six crop types including beans, beetroot, grass, maize, potato and winter wheat, which are dominant crops on the western Tokachi plain, Hokkaido, Japan.

In addition to qualities of remote sensing data, classification algorithms are important to improve classification accuracies of crop maps. Recently, random forests (RF) is a widely used machine learning algorithm consisting of an ensemble of decision trees and it has been an

extremely successful machine learning algorithm for classification and regression method²¹. It has been applied for generating land cover maps^{22, 23} and reached around 65% (tree species identification)¹⁷, 76% (crop types identification)¹⁷ and 90% (greenhouse detection)¹⁹ using MSI data in the previous studies. .

Some studies showed that support vector machine (SVM) performed better than RF for this purpose and it has been widely applied for crop for crop classification^{22, 24-26}. Its robustness to outliers has been demonstrated and SVM is an excellent classifier when the number of input features is large²⁷.

The super learner (SL) methodology²⁵, also called stacking, is an ensemble learning method in which the user-supplied library of algorithms is combined through a convex weighted combination, with the optimal weights to make the cross-validated empirical risk smaller. Therefore, SL could be expected to classify crop types more accurately than the single use of RF or SVM, both considered in this study. Next, an ensemble approach based on SL was applied for improving classification accuracies.

Within this framework, the main objectives of the present study were to evaluate the potential of Sentinel-2 data for crop type classification and the potential of ensemble learning based on RF and SVM.

2. Materials and Methods

2.1. Study area

The study area was located in the western part of Tokachi plain, Hokkaido, Japan (Fig.1, 142°42'51" to 143°08'47" E, 42°43'20" to 43°07'24" N). Main cultivated crops types are beans,

beetroots, grasses, maize, potatoes and winter wheat. The average monthly temperatures were 8.3–21.8°C and monthly precipitation was 12.0–94.5 mm from May to October.

Field location and attribute data, such as crop types, were based on manual surveys and provided by Tokachi Nosai (Obihiro, Hokkaido) as a polygon shape file. A total of 12639 fields (2265 beans fields, 1548 beetroot fields, 2110 grasslands (timothy and orchard grass), 1000 maize fields, 2452 potato fields and 3264 winter wheat fields) were observed. The fields ranged from 0.05 ha to 18.21 ha with an averaged value of 2.54 ha. Grasslands were located on the outskirts of the built-up area.

<Fig. 1 Study area and the distribution of croplands (background map shows Sentinel-2A data obtained on August 11, 2016, R: Band 4, G: Band 3, B: Band 2).>

2.2. Remote sensing data

The data acquired from Sentinel-2 Multispectral Imager (MSI) contained blue, green, red and near-infrared-1 bands at 10 m; red edge 1 to 3, near-infrared-2 and SWIR 1 and 2 at 20 m; and 3 atmospheric bands (Band 1, Band 9 and Band 10) at 60 m. In this study, the three atmospheric bands were removed because they were dedicated to atmospheric corrections and cloud screening²⁸.

Although Sentinel-2A imagery was gathered seven times from May to September 2016 for the whole site, all images were covered with clouds except for one acquired on 11 August. The Level 1C data acquired on August 11, 2016 were downloaded from EarthExplorer (<https://earthexplorer.usgs.gov/>). All bands were converted to 10 m resolution with a cubic convolution resampling method and average reflectance values of each band were calculated for

each field using the field polygons to compensate for spatial variability and to avoid problems related to uncertainty in georeferencing.

Some vegetation indices such as NDVI have been used for improving classification accuracies in previous studies^{16, 22, 29, 30}. Eighty-two published vegetation indices for evaluating various vegetation properties were calculated in this study (Table 1).

Table 1 Vegetation indices calculated from Sentinel-2 MSI data.

Abbreviation	Index	Formula
AFRI1.6 ³¹	Aerosol free vegetation index 1.6	$\frac{Band8a - 0.66 * Band11}{Band8a + 0.66 * Band11}$
AFRI2.1 ³¹	Aerosol free vegetation index 2.1	$\frac{Band8a - 0.5 * Band12}{Band8a + 0.5 * Band12}$
ARI ³²	Anthocyanin reflectance index	$\frac{Band3}{1} - \frac{Band5}{1}$
ARVI ³³	Atmospherically resistant vegetation index	$\frac{(Band8 - (Band4 - \gamma(Band2 - Band4)))}{(Band8 + (Band4 - \gamma(Band2 - Band4)))}$
ARVI2 ³³	Atmospherically resistant vegetation index 2	$-0.18 + 1.17 * \left(\frac{Band8 - Band4}{Band8 + Band4}\right)$
ATSAVI ³⁴	Adjusted transformed soil-adjusted vegetation index	$\frac{a * (Band8 - a * Band4 - b)}{Band8 + Band4 - ab + X(1 + a^2)}$ a = 1.22, b = 0.03, X = 0.08
AVI ³⁵	Ashburn vegetation index	2*Band8a - Band4
BNDVI ³⁶	Blue-normalized difference vegetation index	$(Band8 - Band2)/(Band8 + Band2)$
BRI ³⁷	Browning reflectance index	$\frac{1/Band3 - 1/Band5}{Band6}$
BWDRVI ³⁸	Blue-wide dynamic range vegetation index	$\frac{0.1 * Band7 - Band2}{0.1 * Band7 + Band2}$
CARI ³⁹	Chlorophyll absorption ratio index	$\frac{Band5 * \sqrt{(a * Band4 + Band4 + b)^2}}{Band4} * (a^2 + 1)^{0.5}$ a = (Band5 - Band3)/150 b = Band3 * 550 * a
CCCI ⁴⁰	Canopy chlorophyll content index	$\frac{(Band8 - Band5)}{(Band8 + Band5)} * \frac{(Band8 - Band4)}{(Band8 + Band4)}$
CRI550 ⁴¹	Carotenoid reflectance index 550	$\frac{Band2}{1} - \frac{Band3}{1}$
CRI700 ⁴¹	Carotenoid reflectance index 700	$\frac{Band2}{1} - \frac{Band3}{1}$
CVI ⁴²	Chlorophyll vegetation index	$\frac{Band2 * Band5}{Band8 * Band4} * \frac{1}{(Band3)^2}$

The γ is a weighting function that depends on aerosol type. In this study, a value of 1 for γ .

Datt1 ⁴³	Vegetation index proposed by Datt 1	$\frac{Band8 - Band5}{Band8 - Band4}$
Datt2 ⁴⁴	Vegetation index proposed by Datt 2	$\frac{Band4}{Band3 * Band5}$
Datt3 ⁴⁴	Vegetation index proposed by Datt 3	$\frac{Band8a}{Band3 * Band5}$
DVI ⁴⁵	Differenced vegetation index	$2.4 * Band8 - Band4$
EPIcar ⁴⁴	Eucalyptus pigment index for carotenoid	$0.0049 * \left(\frac{Band4}{Band3 * Band5} \right)^{0.7488}$
EPICChla ⁴⁴	Eucalyptus pigment index for chlorophyll a	$0.0161 * \left(\frac{Band4}{Band3 * Band5} \right)^{0.7784}$
EPICChlab ⁴⁴	Eucalyptus pigment index for chlorophyll a+b	$0.0236 * \left(\frac{Band4}{Band3 * Band5} \right)^{0.7954}$
EPICChlb ⁴⁴	Eucalyptus pigment index for chlorophyll b	$0.0337 * \left(\frac{Band4}{Band3} \right)^{1.8695}$
EVI ¹⁴	Enhanced vegetation index	$2.5 * \frac{Band8 - Band4}{Band8 + 6 * Band4 - 7.5 * Band2 + 1}$
EVI2 ⁴⁶	Enhanced vegetation index 2	$2.4 * \frac{Band8 + Band4 + 1}{Band8 - Band4}$
EVI2.2 ⁴⁷	Enhanced vegetation index 2.2	$2.5 * \frac{Band8 + 2.4 * Band4 + 1}{Band8 - Band4}$
GARI ⁴⁸	Green atmospherically resistant vegetation index	$\frac{Band8 - (Band3 - (Band2 - Band4))}{Band8 - (Band3 + (Band2 - Band4))}$
GBNDVI ⁴⁹	Green-Blue normalized difference vegetation index	$\frac{Band8 - (Band3 + Band2)}{Band8 + (Band3 + Band2)}$
GDVI ⁵⁰	Green difference vegetation index	$Band8 - Band3$
GEMI ⁵¹	Global environment monitoring index	$n * \frac{(1 - 0.25 * n) - Band4 - 0.125}{1 - Band4}$
GLI ⁵²	Green leaf index	$n = \frac{2 * Band5^2 - Band4^2 + 1.5 * Band5 + 0.5 * Band4}{Band5 + Band4 + 0.5}$ $\frac{2 * Band3 - Band5 - Band2}{2 * Band3 + Band5 + Band2}$
GNDVI ⁴⁸	Green normalized difference vegetation index	$\frac{Band8 - Band3}{Band8 + Band3}$
GNDVI2 ⁴⁸	Green normalized difference vegetation index 2	$\frac{Band7 - Band3}{Band7 + Band3}$
GOSAVI ⁵³	Green optimized soil adjusted vegetation index	$\frac{Band8 - Band3}{Band8 + Band3 + 0.16}$
GRNDVI ⁵⁴	Green-Red normalized difference vegetation index	$\frac{Band8 - (Band3 + Band5)}{Band8 + (Band3 + Band5)}$
GVMi ⁵⁵	Global vegetation moisture index	$\frac{(Band8 + 0.1) - (Band12 + 0.02)}{(Band8 + 0.1) + (Band12 + 0.02)}$
Hue ⁵⁶	Hue	$\text{atan} \left(\frac{2 * Band5 - Band3 - Band2}{30.5} * (Band3 - Band2) \right)$
IPVI ⁵⁷	Infrared percentage vegetation index	$\frac{Band8}{Band8 + Band5} \left(\frac{Band5 - Band3}{Band5 + Band5} + 1 \right)$
LCI ⁴³	Leaf chlorophyll index	$\frac{Band8 - Band5}{Band8 + Band4}$

Maccioni ⁵⁸	Vegetation index proposed by Maccioni	$\frac{Band7 - Band5}{Band7 - Band4}$
MCARI ⁵⁹	Modified chlorophyll absorption in reflectance index	$((Band5 - Band4) - 0.2 * (Band5 - Band3)) * \frac{Band5}{Band4}$
MCARI/MTVI2 ⁶⁰	MCARI/MTVI2	$MCARI/MTVI2$
MCARI/OSAVI ⁶¹	MCARI/OSAVI	$MCARI/OSAVI$
MCARI1 ⁶¹	Modified chlorophyll absorption in reflectance index 1	$1.2 * (2.5 * (Band8 - Band4) - 1.3 * (Band8 - Band3))$
MCARI2 ⁶¹	Modified chlorophyll absorption in reflectance index 2	$1.5 * \frac{2.5 * (Band8 - Band4) - 1.3 * (Band8 - Band3)}{\sqrt{(2 * Band8 + 1)^2 - (6 * Band8 - 5 * \sqrt{Band4})}} - 0.5$
MGVI ⁶²	Green vegetation index proposed by Misra	$-0.386 * Band3 - 0.530 * Band4 + 0.535 * Band6 + 0.532 * Band8$
mNDVI ⁶³	Modified normalized difference vegetation index	$\frac{Band8 - Band4}{Band8 + Band4 - 2 * Band2}$
MNSI ⁶²	Non such index proposed by Misra	$0.404 * Band3 + 0.039 * Band4 - 0.505 * Band6 + 0.762 * Band8$
MSAVI ⁶⁴	Modified soil adjusted vegetation index	$\frac{2 * Band8 + 1 - \sqrt{(2 * Band8 + 1)^2 - 8 * (Band8 - Band5)}}{2}$
MSAVI2 ⁶⁴	Modified soil adjusted vegetation index 2	$\frac{2 * Band8 + 1 - \sqrt{(2 * Band8 + 1)^2 - 8 * (Band8 - Band4)}}{2}$
MSBI ⁶²	Soil brightness index proposed by Misra	$0.406 * Band3 + 0.600 * Band4 + 0.645 * Band6 + 0.243 * Band8$
MSR670 ⁶⁵	Modified simple ratio 670/800	$\frac{\frac{Band8}{Band4} - 1}{\sqrt{\frac{Band8}{Band4} + 1}}$
MSRNir/Red ⁶⁶	Modified simple ratio Nir/Red	$\frac{\frac{Band8}{Band5} - 1}{\sqrt{\frac{Band8}{Band5} + 1}}$
MTVI2 ⁶¹	Modified triangular vegetation index 2	$1.5 * \frac{1.2 * (Band8 - Band3) - 2.5 * (Band4 - Band3)}{\sqrt{(2 * Band8 + 1)^2 - (6 * Band8 - 5 * \sqrt{Band4})}} - 0.5$
NBR ⁶⁷	Normalized difference Nir/Swir normalized burn ratio	$\frac{Band8 - Band12}{Band8 + Band12}$
ND774/677 ⁶⁸	Normalized difference 774/677	$\frac{Band7 - Band4}{Band7 + Band4}$
NDII ⁶⁹	Normalized difference infrared index	$\frac{Band8 - Band11}{Band8 + Band11}$
NDRE ⁷⁰	Normalized difference Red-edge	$\frac{Band7 - Band5}{Band7 + Band5}$
NDSI ⁷¹	Normalized difference salinity index	$\frac{Band11 - Band12}{Band11 + Band12}$
NDVI ¹²	Normalized difference vegetation index	$\frac{Band8 - Band4}{Band8 + Band4}$
NDVI2 ⁵⁰	Normalized difference vegetation index 2	$\frac{Band12 - Band8}{Band12 + Band8}$
NGRDI ⁶⁸	Normalized green red difference index	$\frac{Band3 - Band5}{Band3 + Band5}$
OSAVI ^{53, 72}	Optimized soil adjusted vegetation index	$1.16 * \frac{Band3 + Band5}{Band8 - Band4}$

PNDVI ⁵⁴	Pan normalized difference vegetation index	$\frac{Band8 - (Band3 + Band5 + Band2)}{Band8 + (Band3 + Band5 + Band2)}$
PVR ⁷³	Photosynthetic vigour ratio	$\frac{Band3 - Band4}{Band3 + Band4}$
RBNDVI ⁵⁴	Red-Blue normalized difference vegetation index	$\frac{Band8 - (Band4 + Band2)}{Band8 + (Band4 + Band2)}$
RDVI ⁷⁴	Renormalized difference vegetation index	$\frac{Band8 - Band4}{\sqrt{Band8 + Band4}}$
REIP ⁷⁵	Red-edge inflexion point	$700 + 40 * \left(\frac{\left(\frac{Band4 + Band7}{2} \right) - Band5}{Band6 - Band5} \right)$
Rre ⁷⁶	Reflectance at the inflexion point	$\frac{Band4 + Band7}{2}$
SAVI ¹³	Soil adjusted vegetation index	$1.5 * \frac{Band8 - Band4}{Band8 + Band4 + 0.5}$
SBL ⁴⁵	Soil background line	$Band8 - 2.4 * Band4$
SIPI ⁷⁷	Structure intensive pigment index	$\frac{Band8 - Band2}{Band8 - Band4}$
SIWSI ⁷⁸	Shortwave infrared water stress index	$\frac{Band8a - Band11}{Band8a + Band11}$
SLAVI ⁷⁹	Specific leaf area vegetation index	$\frac{Band8}{Band4 + Band12}$
TCARI ⁵⁹	Transformed chlorophyll absorption Ratio	$3 * \left((Band5 - Band4) - 0.2 * (Band5 - Band3) \left(\frac{Band5}{Band4} \right) \right)$
TCARI/OSAVI ⁷²	TCARI/OSAVI	$TCARI/OSAVI$
TCI ^{42, 80}	Triangular chlorophyll index	$1.2 * (Band5 - Band3) - 1.5 * (Band4 - Band3) * \sqrt{\frac{Band5}{Band4}}$
TVI ⁸¹	Transformed vegetation index	$\sqrt{NDVI + 0.5}$
VARI700 ⁸²	Visible atmospherically resistant index 700	$\frac{Band5 - 1.7 * Band4 + 0.7 * Band2}{Band5 + 2.3 * Band4 - 1.3 * Band2}$
VARIgreen ⁸²	Visible atmospherically resistant index green	$\frac{Band3 + Band4 - Band2}{Band5 - Band4}$
VI700 ⁸³	Vegetation index 700	$\frac{Band5 + Band4}{0.1 * Band8 - Band4}$
WDRVI ⁸⁴	Wide dynamic range vegetation index	$\frac{0.1 * Band8 - Band4}{0.1 * Band8 + Band4}$

2.3. Classification algorithm

All samples were divided into the following three groups using a stratified random sampling approach: training data (50%) for developing classification models, validation data (25%) for hyperparameter tuning and test data (25%) for evaluation of classification accuracies⁸⁵ and table 2 shows the numbers of fields of each crop type.

Table 2 Crop type and number of fields.

Crop type	Training data	Validation data	Test data
Beans	1132	566	567
Beetroot	774	387	387
Grassland	1055	527	528
Maize	500	250	250
Potato	1226	613	613
Wheat	1632	816	816

SVM partitions data using maximum separation margins⁸⁶ and the ‘kernel trick’ has frequently been applied instead of attempting to fit a non-linear model in previous studies²⁹. In this study, the Gaussian Radial Basis Function (RBF) kernel, which has mostly been used for classification purposes²⁹, was used as a kernel and two parameters were tuned to control the flexibility of the classifier, the regularization parameter C and the kernel bandwidth γ . If the C value is too large, there is a high penalty for no separable points and we may store many support vectors and overfit. If it is too small, there may be under-fitting. It controls the trade-off between errors of the SVM on training data and margin maximization ($C = \infty$ leads to hard margin SVM). The γ value defines how far the influence of a single training example reaches, with low values meaning ‘far’ and high values meaning ‘close.’

RF is an ensemble learning technique composed of multiple decision trees based on random bootstrapped samples of the training data⁸⁷. The output is determined by a majority vote of the results of decision trees. There are two user-defined hyperparameters including the number of trees (n_{tree}) and the number of variables used to split the nodes (m_{try}). If n_{tree} is made larger, the generalization error always converges, and over-training will not be a problem. On the other hand, a reduction in m_{try} makes each individual decision tree weaker.

The best combinations of these hyperparameters were determined using the Gaussian process, Bayesian optimization⁸⁸, which has been widely applied for hyperparameter tuning of machine learning algorithms¹.

Ensemble machine learning methods have been used to obtain better predictive performance than from single learning algorithms and the SL methodology has been proposed⁸⁹. In this method, given algorithms are combined through a convex weighted combination to minimize cross-validated errors. First, classification models based on RF or SVM were trained as the base algorithms using the training data. Next, a ten-fold cross-validation was performed on each and the cross-validated predicted results were obtained. N is the number of rows in the training data, cross-validated predicted results were combined and an N by two matrix was obtained as the “level-one” data and meta-learning model was generated. To predict the test data, the predictions from the base learners were feed into the meta-learning model to generate the ensemble prediction.

The data-based sensitivity analysis (DSA)⁹⁰, which performs a pure black box use of the fitted models by querying the fitted models with sensitivity samples and recording their responses, was applied for assessing the sensitivity of the classification models.

2.4. Accuracy assessment

Classification accuracies were evaluated based on the simple measures of quantity disagreement (QD) and allocation disagreement (AD)⁹¹. They provide an effective summary of confusion matrices⁹².

The proportion of fields that are classified as crop i and their actual classes are crop j (P_{ij}) is expressed in the following equation (1):

$$P_{ij} = W_i \frac{n_{ij}}{n_{i+}} \quad (1)$$

where W_i are the fields classified as crop i , n_{ij} is the number of fields classified as crop i and their actual classes are crop j . n_{i+} is the row totals of the confusion matrix. In this case, AD and QD are calculated using the following equations (2–5):

$$AD_i = 2 \min(p_{i+}, p_{+i}) - 2p_{ii} \quad (2)$$

$$AD = \frac{1}{2} \sum_{i=1}^{N_c} AD_i \quad (3)$$

$$QD_i = |p_{i+} - p_{+i}| \quad (4)$$

$$QD = \frac{1}{2} \sum_{i=1}^{N_c} QD_i \quad (5)$$

where N_c is the number of classes (six in this study), p_{i+} and p_{+i} are the row and column totals of the confusion matrix, AD_i is the allocation disagreement of crop i and QD_i is the quantity disagreement of crop i . The sum of QD_i (QD) and AD_i (AD) are calculated and the total disagreement can be evaluated by the sum of QD and AD⁹¹.

In addition, three indicators including overall accuracy (OA, equations (6)), producer's accuracy (PA, equations (7)) and user's accuracy (UA, equations (8)) were calculated because they have widely been applied for assessing classification accuracies.

$$OA = \sum_{i=1}^N p_{ii} / N \quad (6)$$

$$PA = p_{ii} / R_i \quad (7)$$

$$UA = p_{ii} / C_i \quad (8)$$

where N is the number of fields, R_i and C_i represent the total number of crop i in the correct data and the total number from the classification results, respectively. McNemar's test⁹³ has been used to judge whether the differences between two given classification results were significant⁹⁴ and it was also applied in this study.

3. Results and Discussion

3.1. Classification accuracy

Crop classification maps are shown in Fig. 2, the maximum, minimum and averaged accuracies of ten repetitions and confusion matrices when all the repetitions were merged are shown in Table 3 and 4. Averaged OAs were 89.0% for RF, 90.6% for SVM and 91.6% for the ensemble machine learning method and the mean PAs and mean UAs derived using the machine learning algorithms were greater than 0.8, excepting those of RF (mean UA for maize was 0.797). All machine learning algorithms performed well in classifying croplands. Especially, the good accuracies were confirmed for the PAs and UAs for wheat (more than 93.8%) and beet (more than 89.9%). However, the chi-square values based on McNemar's tests were 12.02 – 40.60, 27.78 – 62.43 and 17.00 – 51.60 for R – SVM, RF – SL and SVM – SL, respectively. As the results, significant differences were confirmed among the results of three machine learning algorithms ($p < 0.05$).

Classification results by SL had the best OA and AD+QD (8.5%) and SVM had a slightly better PA of wheat (97.1%). On the contrary, identifying maize fields was difficult due to the similarity in their reflectance. Grasses cultivation employs fewer controls and then a lot of weeds were mixed with timothy and orchard grass in grasslands. As a result, variation in reflectance features were larger than in other crop types, causing misclassifications of relatively larger fields.

<Fig. 2. Crop classification map generated by (a) RF, (b) SVM and (c) SL.>

Table 3 Classification accuracies of each algorithm.

	RF			SVM			SL		
	Minimum	Maximum	Mean±std	Minimum	Maximum	Mean±std	Minimum	Maximum	Mean±std
PA									
Beans	80.6%	86.4%	83.4±1.6%	81.1%	90.5%	86.2±2.2%	84.7%	90.3%	87.6±1.4%
Beet	89.9%	94.8%	93.0±1.3%	91.0%	96.4%	94.5±1.5%	93.8%	96.1%	95.1±0.6%
Grassland	84.3%	88.3%	86.0±1.2%	86.7%	93.8%	89.4±2.5%	89.8%	94.3%	92.1±1.4%
Maize	78.8%	84.8%	80.8±1.7%	78.8%	87.6%	83.0±3.1%	81.2%	87.6%	84.6±1.8%
Potato	82.9%	89.7%	87.0±1.8%	83.5%	89.9%	87.6±1.9%	84.0%	89.7%	88.1±1.6%
Wheat	96.4%	97.9%	97.0±0.5%	96.3%	97.5%	97.1±0.4%	95.7%	97.5%	97.0±0.7%
UA									
Beans	84.9%	88.6%	86.8±1.1%	82.0%	91.4%	86.4±2.9%	83.4%	90.3%	88.6±2.0%
Beet	94.5%	96.9%	95.6±0.8%	94.3%	97.3%	95.7±0.9%	95.1%	97.1%	96.0±0.6%
Grassland	88.0%	93.3%	91.0±1.4%	89.9%	96.6%	94.0±2.3%	93.8%	97.7%	95.7±1.1%
Maize	77.8%	82.0%	79.7±1.3%	78.4%	87.3%	81.9±2.2%	81.4%	85.2%	83.6±1.4%
Potato	78.5%	83.1%	81.5±1.2%	82.1%	87.8%	85.2±1.9%	83.0%	86.8%	85.4±1.1%
Wheat	93.8%	96.1%	95.0±0.7%	94.5%	97.2%	95.9±0.8%	95.1%	97.2%	96.2±0.6%
OA	88.5%	89.4%	89.0±0.2%	89.3%	92.0%	90.6±0.9%	90.2%	92.2%	91.6±0.6%
κ	85.9%	87.0%	86.5±0.3%	86.8%	90.2%	88.4±1.1%	88.0%	90.5%	89.6±0.8%
AD	8.0%	9.9%	9.0±0.6%	6.5%	9.7%	7.9±1.0%	6.5%	8.8%	7.3±0.7%
QD	1.3%	2.8%	2.0±0.5%	0.7%	2.5%	1.5±0.6%	0.6%	2.3%	1.2±0.5%

Table 4 Confusion matrices for (a) RF, (b) SVM and (c) SL.

(a) RF		Reference data					
		Beans	Beetroot	Grasslands	Maize	Potato	Wheat
Classified data	Beans	4726	59	247	100	287	26
	Beet	48	3599	23	28	65	1
	Grasslands	172	65	4543	52	116	43
	Maize	139	21	128	2019	177	48
	Potato	503	119	230	235	5332	123
	Wheat	82	7	109	66	153	7919

(b) SVM		Reference data					
		Beans	Beetroot	Grasslands	Maize	Potato	Wheat
Classified data	Beans	4888	77	212	119	333	34
	Beet	61	3659	17	22	63	2
	Grasslands	110	34	4720	40	70	49
	Maize	112	14	130	2076	166	40

Potato	429	79	121	189	5368	115
Wheat	70	7	80	54	130	7920

(c) SL

		Reference data					
		Beans	Beetroot	Grasslands	Maize	Potato	Wheat
Classified data	Beans	4965	82	105	83	333	42
	Beet	61	3680	11	17	61	3
	Grasslands	59	17	4861	37	52	53
	Maize	85	8	121	2114	169	32
	Potato	426	77	113	200	5403	112
	Wheat	74	6	69	49	112	7918

Figure 3 shows the relationship between field area and misclassified fields for each algorithm after ten repetitions (i.e. the total number is ten times of that of the test data). More than 75% of the misclassified fields were less than 200 a in area for all algorithms, and 95.1% (RF), 95.5% (SVM) and 96.1% (SL) of misclassified fields were below 450 a. Applying stacking made the model more robust for classifying smaller fields and the number of misclassified croplands decreased (813 fields for smaller than 50 a) compared with the results by RF (909 fields for smaller than 50 a) and SVM (855 fields for smaller than 50 a). It was especially useful for identifying beans fields. It was not effective for identifying small grasslands since grass cultivation employs fewer controls and many weeds were present in grasslands. However, stacking was useful for identifying grasslands more than 500 a, which had a certain homogeneity with *Dactylis glomerata* or *Phleum pretense* in the MSI image.

<Fig.3 Relationship between field area and misclassified fields.>

3.2. Sensitive factor analysis

Reflectance values obtained from Sentinel-2A are shown in Fig. 4. Differences in reflectance were particularly clear between wheat and beans since the wheat harvest was finished on 11

August and the reflectance of wheat fields was similar to that of bare soil. Beetroot had the steepest gradient between Bands 5 and 6 and some differences in the reflectance values at Band 11 were confirmed between maize and potato. Differences in the reflectance patterns between grass and beans were not clear.

<Fig. 4 Reflectance spectra of each crop.>

To clarify which variables contributed to identifying each crop type, DSA was conducted for each algorithm and their importance values were calculated.

For identifying beans fields, Datt3 (6.0%, 6.6% and 6.3% for RF, SVM and SL, respectively) and REIP (6.4%, 8.2% and 7.3% for RF, SVM and SL, respectively) played important roles in the three algorithms. Some variables (the reflectance values at Bands 2 and 3, AFRI2.1, CVI and NDSI) possessed importance values of more than 5.0% in the RF-based model, while no variables except for Datt3 and REIP had importance values of more than 5.0% for SVM and SL. Even though the importance values of GEMI, Maccioni and MNSI in SVM were less than 5.0%, they were more than 5 times those in RF. AFRI1.6 and SIWSI were useful for identifying beetroot fields and AFRI1.6 occupied 11.1%, 6.8% and 9.0% and SIWSI occupied 10.6%, 7.1% and 8.9% of the importance for RF, SVM and SL, respectively. GEMI and NDSI also had importance values of more than 10% for RF, but were less than 5% for the others. In contrast, REIP was useful in SVM and it occupied 9.1% of the importance in SVM. AFRI1.6, REIP and MNSI were effective for identifying grassland for all algorithms, while SIWSI played an important role (7.8%) for RF and the reflectance at Band 6 played an important role (8.2%) for SVM. For identifying maize fields, no variable had importance values more than 5.0% for any algorithm, but the importance value of REIP was 25.3% for SVM (2.9% for RF). CRI550, CRI700 and MSBI were 9.1%, 12.9% and 5.6% in RF, respectively (those in SVM were 2.4%,

2.2% and 3.6%, respectively). REIP played the greatest role for identifying potato fields in all algorithms (12.8%, 6.9% and 9.9% for RF, SVM and SL, respectively). The importance values of CCCI and CVI were also high in RF (9.9%) but those in SVM were less than 3.0%. In contrast, Maccioni had an importance of 6.9% in SVM but in RF was 1.4%. REIP also played a great role for identifying wheat fields in SVM but 1.2% of the importance value was confirmed in RF while AVI occupied 15.1% in RF (1.2% in SVM). However, the original reflectance values possessed importance values of less than 1.0%.

In this season, the photosynthetic activities of each crop type were different; maize is a C4 plant, beans and beetroot were in their growing season, grassland was after second harvest, potato growth was inhibited by chemicals for easy harvesting and wheat fields were cultivated. Besides indices related to chlorophyll content, the additional use of shortwave infrared data contributed to the estimation of photosynthetic pigments, water, nitrogen, cellulose, lignin, phenols, and leaf mass per area (e.g. NDSI). As a result, vegetation indices had greater influence on the classification results than the original reflectance. However, there were differences among algorithms in which vegetation indices were more important. The importance values in SL were near the averaged values of RF and SVM. So, the differences in importance between RF and SVM were useful when stacking was applied, and the modification contributed to identifying croplands with higher accuracies.

4. Conclusions and future work

Cropland classifications were conducted using a single image from Sentinel-2 MSI and the suitability and accuracy of vegetation indices from the original reflectance data from Sentinel-2 MSI were assessed.

Of the two algorithms applied (RF and SVM), the accuracy of SVM was superior and 89.3–92.0% of OAs were confirmed. Furthermore, stacking contributed to higher OAs (90.2–92.2%) and significant differences were confirmed with the results of SVM. Based on DSA, the vegetation indices calculated from the original reflectance from Sentinel-2 MSI data were useful to identify the specific crop types. Although the vegetation indices that played the largest roles were different between RF and SVM, stacking helped to modify and reduce the importance of specific variables, which might prevent overfitting. Stacking should be utilized to monitor agricultural fields for improving classification accuracies.

The field is used as a basic unit in classification and some problems related to the borders of fields remain to be resolved. We are planning to evaluate the potential of geographic object-based image analysis in conjunction with MSI data and address this question in future work.

Disclosures

No potential conflicts of interest are reported by the authors.

Acknowledgments

The authors would like to thank Tokachi Nosai for providing the field data.

References

1. R. Sonobe, Y. Yamaya, H. Tani, X. F. Wang, N. Kobayashi, and K. I. Mochizuki, "Assessing the suitability of data from Sentinel-1A and 2A for crop classification," *GIScience & Remote Sensing* 54, 918-938 (2017) [doi: 10.1080/15481603.2017.1351149].
2. R. Sonobe, Y. Miura, T. Sano, and H. Horie, "Estimating leaf carotenoid contents of shade grown tea using hyperspectral indices and PROSPECT-D inversion," *International Journal of Remote Sensing* 39, 1306-1320 (2018) [doi:10.1080/01431161.2017.1407050].

3. C. Rankine, G. A. Sanchez-Azofeifa, J. A. Guzman, M. M. Espirito-Santo, and I. Sharp, "Comparing MODIS and near-surface vegetation indexes for monitoring tropical dry forest phenology along a successional gradient using optical phenology towers," *Environmental Research Letters* 12 (2017) [doi:10.1088/1748-9326/aa838c].
4. S. S. Liu, W. H. Zhao, H. F. Shen, and L. P. Zhang, "Regional-scale winter wheat phenology monitoring using multisensor spatio-temporal fusion in a South Central China growing area," *Journal of Applied Remote Sensing* 10, 16 (2016) [doi:10.1117/1.JRS.10.046029].
5. J. Vithanage, S. N. Miller, and K. Driese, "Land cover characterization for a watershed in Kenya using MODIS data and Fourier algorithms," *Journal of Applied Remote Sensing* 10, 13 (2016) [doi:10.1117/1.JRS.10.045015].
6. R. Sonobe, Y. Yamaya, H. Tani, X. F. Wang, N. Kobayashi, and K. I. Mochizuki, "Evaluating metrics derived from Landsat 8 OLI imagery to map crop cover," *Geocarto International* (2018) [doi:10.1080/10106049.2018.1425739].
7. G. P. Asner, "Biophysical and biochemical sources of variability in canopy reflectance," *Remote Sensing of Environment* 64, 234-253 (1998) [doi:10.1016/S0034-4257(98)00014-5].
8. M. A. Pena, R. Liao, and A. Brenning, "Using spectrotemporal indices to improve the fruit-tree crop classification accuracy," *ISPRS Journal of Photogrammetry and Remote Sensing* 128, 158-169 (2017) [doi:10.1016/j.isprsjprs.2017.03.019].
9. D. Bankestad, and T. Wik, "Growth tracking of basil by proximal remote sensing of chlorophyll fluorescence in growth chamber and greenhouse environments," *Computers and Electronics in Agriculture* 128, 77-86 (2016) [doi:10.1016/j.compag.2016.08.004].
10. Z. Wang, C. F. Xue, W. T. Quan, and H. J. He, "Spatiotemporal variations of forest phenology in the Qinling Mountains and its response to a critical temperature of 10 degrees C," *Journal of Applied Remote Sensing* 12, 15 (2018) [doi:10.1117/1.JRS.12.022202].
11. M. Morin, R. Lawrence, K. Repasky, T. Sterling, C. McCann, and S. Powell, "Agreement analysis and spatial sensitivity of multispectral and hyperspectral sensors in detecting vegetation stress at

- management scales," *Journal of Applied Remote Sensing* 11, 18 (2017) [doi:10.1117/1.JRS.11.046025].
12. C. J. Tucker, "Red and photographic infrared linear combinations for monitoring vegetation," *Remote Sensing of Environment* 8, 127-150 (1979) [doi:10.1016/0034-4257(79)90013-0].
 13. A. R. Huete, "A Soil-Adjusted Vegetation Index (SAVI)," *Remote Sensing of Environment* 25, 295-309 (1988) [doi:10.1016/0034-4257(88)90106-X].
 14. A. Huete, K. Didan, T. Miura, E. P. Rodriguez, X. Gao, and L. G. Ferreira, "Overview of the radiometric and biophysical performance of the MODIS vegetation indices," *Remote Sensing of Environment* 83, 195-213 (2002) [doi: 10.1016/S0034-4257(02)00096-2].
 15. C. E. Holden, and C. E. Woodcock, "An analysis of Landsat 7 and Landsat 8 underflight data and the implications for time series investigations," *Remote Sensing of Environment* 185, 16-36 (2016) [doi: 10.1016/j.rse.2016.02.052].
 16. M. Belgiu, and O. Csillik, "Sentinel-2 cropland mapping using pixel-based and object-based time-weighted dynamic time warping analysis," *Remote Sensing of Environment* 204, 509-523 (2018) [doi: 10.1016/j.rse.2017.10.005].
 17. M. Immitzer, F. Vuolo, and C. Atzberger, "First Experience with Sentinel-2 Data for Crop and Tree Species Classifications in Central Europe," *Remote Sensing* 8, 27 (2016) [doi: 10.3390/rs8030166].
 18. Y. Palchowdhuri, R. Valcarce-Dineiro, P. King, and M. Sanabria-Soto, "Classification of multi-temporal spectral indices for crop type mapping: a case study in Coalville, UK," *Journal of Agricultural Science* 156, 24-36 (2018) [doi: 10.1017/S0021859617000879].
 19. A. Novelli, M. A. Aguilar, A. Nemmaoui, F. J. Aguilar, and E. Tarantino, "Performance evaluation of object based greenhouse detection from Sentinel-2 MSI and Landsat 8 OLI data: A case study from Almeria (Spain)," *International Journal of Applied Earth Observation and Geoinformation* 52, 403-411 (2016) [doi: 10.1016/j.jag.2016.07.011].

20. Y. Du, Y. H. Zhang, F. Ling, Q. M. Wang, W. B. Li, and X. D. Li, "Water Bodies' Mapping from Sentinel-2 Imagery with Modified Normalized Difference Water Index at 10-m Spatial Resolution Produced by Sharpening the SWIR Band," *Remote Sensing* 8 (2016) [doi: 10.3390/rs8040354].
21. G. Biau, and E. Scornet, "A random forest guided tour," *Test* 25, 197-227 (2016) [doi: 10.1007/s11749-016-0481-7].
22. S. Ferrant, A. Selles, M. Le Page, P. A. Herrault, C. Pelletier, A. Al-Bitar, S. Mermoz, S. Gascoin, A. Bouvet, M. Saqalli, B. Dewandel, Y. Caballero, S. Ahmed, J. C. Marechal, and Y. Kerr, "Detection of Irrigated Crops from Sentinel-1 and Sentinel-2 Data to Estimate Seasonal Groundwater Use in South India," *Remote Sensing* 9 (2017) [doi: 10.3390/rs9111119].
23. A. O. Onojeghuo, G. A. Blackburn, Q. M. Wang, P. M. Atkinson, D. Kindred, and Y. X. Miao, "Mapping paddy rice fields by applying machine learning algorithms to multi-temporal Sentinel-1A and Landsat data," *International Journal of Remote Sensing* 39, 1042-1067 (2018) [doi:10.1080/01431161.2017.1395969].
24. R. Sonobe, H. Tani, X. Wang, N. Kobayashi, and H. Shimamura, "Discrimination of crop types with TerraSAR-X-derived information," *Physics and Chemistry of the Earth* 83-84, 2-13 (2015) [doi:10.1016/j.pce.2014.11.001].
25. J. K. Gilbertson, and A. van Niekerk, "Value of dimensionality reduction for crop differentiation with multi-temporal imagery and machine learning," *Computers and Electronics in Agriculture* 142, 50-58 (2017) [doi:10.1016/j.compag.2017.08.024].
26. R. Sonobe, H. Tani, X. Wang, N. Kobayashi, and H. Shimamura, "Random forest classification of crop type using multi- temporal TerraSAR- X dual- polarimetric data," *Remote Sensing Letters* 5, 157-164 (2014) [doi:10.1080/2150704X.2014.889863].
27. G. Camps-Valls, L. Gomez-Chova, J. Calpe-Maravilla, J. D. Martin-Guerrero, E. Soria-Olivas, L. Alonso-Chorda, and J. Moreno, "Robust support vector method for hyperspectral data classification and knowledge discovery," *IEEE Transactions on Geoscience and Remote Sensing* 42, 1530-1542 (2004) [doi: 10.1109/TGRS.2004.827262].

28. M. Drusch, U. Del Bello, S. Carlier, O. Colin, V. Fernandez, F. Gascon, B. Hoersch, C. Isola, P. Laberinti, P. Martimort, A. Meygret, F. Spoto, O. Sy, F. Marchese, and P. Bargellini, "Sentinel-2: ESA's Optical High-Resolution Mission for GMES Operational Services," *Remote Sensing of Environment* 120, 25-36 (2012) [doi:10.1016/j.rse.2011.11.026].
29. A. Chatziantoniou, G. P. Petropoulos, and E. Psomiadis, "Co-Orbital Sentinel 1 and 2 for LULC Mapping with Emphasis on Wetlands in a Mediterranean Setting Based on Machine Learning," *Remote Sensing* 9 (2017) [doi:10.3390/rs9121259].
30. E. M. D. Silveira, M. D. de Menezes, F. W. Acerbi, M. Terra, and J. M. de Mello, "Assessment of geostatistical features for object-based image classification of contrasted landscape vegetation cover," *Journal of Applied Remote Sensing* 11, 15 (2017) [doi:10.1117/1.JRS.11.036004].
31. A. Karnieli, Y. J. Kaufman, L. Remer, and A. Wald, "AFRI - aerosol free vegetation index," *Remote Sensing of Environment* 77, 10-21 (2001) [doi:10.1016/S0034-4257(01)00190-0].
32. A. A. Gitelson, O. B. Chivkunova, and M. N. Merzlyak, "Nondestructive estimation of anthocyanins and chlorophylls in anthocyanic leaves," *American Journal of Botany* 96, 1861-1868 (2009) [doi:10.3732/ajb.0800395].
33. Y. J. Kaufman, and D. Tanre, "Atmospherically resistant vegetation index (ARVI) for EOS-MODIS," *IEEE Transactions on Geoscience and Remote Sensing* 30, 261-270 (1992) [doi:10.1109/36.134076].
34. F. Baret, and G. Guyot, "Potentials and limits of vegetation indices for LAI and APAR assessment," *Remote Sensing of Environment* 35, 161-173 (1991) [doi:10.1016/0034-4257(91)90009-U].
35. P. Ashburn, "The vegetative index number and crop identification," *The LACIE Symposium Proceedings of the Technical Session*, 843-855 (1978).
36. C. G. Yang, J. H. Everitt, and J. M. Bradford, "Airborne hyperspectral imagery and linear spectral unmixing for mapping variation in crop yield," *Precision Agriculture* 8, 279-296 (2007) [doi:10.1007/s11119-007-9045-x].

37. O. B. Chivkunova, A. E. Solovchenko, S. G. Sokolova, M. N. Merzlyak, I. V. Reshetnikova, and A. A. Gitelson, "Reflectance Spectral Features and Detection of Superficial Scald -induced Browning in Storing Apple Fruit," *Journal of Russian Phytopathological Society* 2, 73-77 (2001).
38. D. W. Hancock, and C. T. Dougherty, "Relationships between blue- and red-based vegetation indices and leaf area and yield of alfalfa," *Crop Science* 47, 2547-2556 (2007)
[doi:10.2135/cropsci2007.01.0031].
39. M. S. Kim, C. S. T. Daughtry, E. W. Chappelle, J. E. McMurtrey, and C. L. Walthall, "The use of high spectral resolution bands for estimating absorbed photosynthetically active radiation (A_{par})," presented at the 6th International Symposium on Physical Measurements and Signatures in Remote Sensing Val D'Isere, France 1994.
40. D. M. El-Shikha, E. M. Barnes, T. R. Clarke, D. J. Hunsaker, J. A. Haberland, P. J. Pinter, P. M. Waller, and T. L. Thompson, "Remote sensing of cotton nitrogen status using the Canopy Chlorophyll Content Index (CCCI)," *Transactions of the ASABE* 51, 73-82 (2008)
[doi:10.2134/agronj2011.0124].
41. A. A. Gitelson, M. N. Merzlyak, and O. B. Chivkunova, "Optical properties and nondestructive estimation of anthocyanin content in plant leaves," *Photochemistry and Photobiology* 74, 38-45 (2001) [doi:0.1562/0031-8655(2001)074<0038:OPANEO>2.0.CO;2].
42. E. R. Hunt, C. S. T. Daughtry, J. U. H. Eitel, and D. S. Long, "Remote Sensing Leaf Chlorophyll Content Using a Visible Band Index," *Agronomy Journal* 103, 1090-1099 (2011)
[doi:10.1016/j.jag.2012.07.020].
43. B. Datt, "Remote sensing of water content in Eucalyptus leaves," *Australian Journal of Botany* 47, 909-923 (1999) [doi:10.1071/BT98042].
44. B. Datt, "Remote sensing of chlorophyll a, chlorophyll b, chlorophyll a+b, and total carotenoid content in eucalyptus leaves," *Remote Sensing of Environment* 66, 111-121 (1998)
[doi:10.1016/S0034-4257(98)00046-7].

45. A. J. Richardson, and C. L. Wiegand, "Distinguishing Vegetation from Soil Background Information," *Photogrammetric Engineering & Remote Sensing* 43, 1541-1552 (1978).
46. T. Miura, H. Yoshioka, K. Fujiwara, and H. Yamamoto, "Inter-comparison of ASTER and MODIS surface reflectance and vegetation index products for synergistic applications to natural resource monitoring," *Sensors* 8, 2480-2499 (2008) [doi:10.3390/s8042480].
47. Z. Y. Jiang, A. R. Huete, K. Didan, and T. Miura, "Development of a two-band enhanced vegetation index without a blue band," *Remote Sensing of Environment* 112, 3833-3845 (2008) [doi:10.1016/j.rse.2008.06.006].
48. A. A. Gitelson, Y. J. Kaufman, and M. N. Merzlyak, "Use of a green channel in remote sensing of global vegetation from EOS-MODIS," *Remote Sensing of Environment* 58, 289-298 (1996) [doi:10.1016/S0034-4257(96)00072-7].
49. F. M. Wang, J. F. Huang, and L. Chen, "Development of a vegetation index for estimation of leaf area index based on simulation modeling," *Journal of Plant Nutrition* 33, 328-338 (2010) [doi:10.1080/01904160903470380].
50. C. J. Tucker, "Monitoring corn and soybean crop development with hand-held radiometer spectral data," *Remote Sensing of Environment* 8, 237-248 (1979) [doi:10.1016/0034-4257(79)90004-X].
51. B. Pinty, and M. M. Verstraete, "GEMI: a non-linear index to monitor global vegetation from satellites," *Vegetatio* 101, 15-20 (1992).
52. N. Gobron, B. Pinty, M. M. Verstraete, and J. L. Widlowski, "Advanced vegetation indices optimized for up-coming sensors: Design, performance, and applications," *IEEE Transactions on Geoscience and Remote Sensing* 38, 2489-2505 (2000) [doi:10.1109/36.885197].
53. G. Rondeaux, M. Steven, and F. Baret, "Optimization of soil-adjusted vegetation indices," *Remote Sensing of Environment* 55, 95-107 (1996) [doi:10.1016/0034-4257(95)00186-7].
54. F.-M. Wang, J.-F. Huang, Y.-L. Tang, and X.-Z. Wang, "New Vegetation Index and Its Application in Estimating Leaf Area Index of Rice," *Rice Science* 14, 195-203 (2007) [doi:10.1016/S1672-6308(07)60027-4].

55. E. P. Glenn, P. L. Nagler, and A. R. Huete, "Vegetation Index Methods for Estimating Evapotranspiration by Remote Sensing," *Surveys in Geophysics* 31, 531-555 (2010) [doi:10.1007/s10712-010-9102-2].
56. R. Escadafal, A. Belghith, and H. Ben Moussa, "Indices spectraux pour la degradation des milieux naturels en Tunisie aride," in 6e Symposium international sur les mesures physiques et signatures en teledetection (Val d'Isere, France, 1994), pp. 253-259.
57. R. E. Crippen, "Calculating the vegetation index faster," *Remote Sensing of Environment* 34, 71-73 (1990) [doi:10.1016/0034-4257(90)90085-Z].
58. A. Maccioni, G. Agati, and P. Mazzinghi, "New vegetation indices for remote measurement of chlorophylls based on leaf directional reflectance spectra," *Journal of Photochemistry and Photobiology B-Biology* 61, 52-61 (2001) [doi:10.1016/S1011-1344(01)00145-2].
59. C. S. T. Daughtry, C. L. Walthall, M. S. Kim, E. B. de Colstoun, and J. E. McMurtrey, "Estimating corn leaf chlorophyll concentration from leaf and canopy reflectance," *Remote Sensing of Environment* 74, 229-239 (2000) [doi:10.1016/S0034-4257(00)00113-9].
60. J. U. H. Eitel, D. S. Long, P. E. Gessler, and A. M. S. Smith, "Using in-situ measurements to evaluate the new RapidEye (TM) satellite series for prediction of wheat nitrogen status," *International Journal of Remote Sensing* 28, 4183-4190 (2007) [doi:10.1080/01431160701422213].
61. D. Haboudane, J. R. Miller, E. Pattey, P. J. Zarco-Tejada, and I. B. Strachan, "Hyperspectral vegetation indices and novel algorithms for predicting green LAI of crop canopies: Modeling and validation in the context of precision agriculture," *Remote Sensing of Environment* 90, 337-352 (2004) [doi:10.1016/j.rse.2003.12.013].
62. P. N. Misra, S. G. Wheeler, and R. E. Oliver, "Kauth-Thomas brightness and greenness axes," *Contract NASA 9-14350*, 23-46 (1977).
63. R. Main, M. A. Cho, R. Mathieu, M. M. O'Kennedy, A. Ramoelo, and S. Koch, "An investigation into robust spectral indices for leaf chlorophyll estimation," *ISPRS Journal of Photogrammetry and Remote Sensing* 66, 751-761 (2011) [doi:10.1016/j.isprsjprs.2011.08.001].

64. J. Qi, A. Chehbouni, A. R. Huete, Y. H. Kerr, and S. Sorooshian, "A modified soil adjusted vegetation index," *Remote Sensing of Environment* 48, 119-126 (1994) [doi:10.1016/0034-4257(94)90134-1].
65. J. M. Chen, "Evaluation of Vegetation Indices and a Modified Simple Ratio for Boreal Applications," *Canadian Journal of Remote Sensing* 22, 229-242 (1996) [doi:10.1080/07038992.1996.10855178].
66. J. M. Chen, and J. Cihlar, "Retrieving leaf area index of boreal conifer forests using Landsat TM images," *Remote Sensing of Environment* 55, 153-162 (1996) [doi:10.1016/0034-4257(95)00195-6].
67. C. Key, and N. Benson, "Landscape assessment: Ground measure of severity; The Composite Burn Index, and remote sensing of severity, the Normalized Burn Index. Landscape assessment: Ground measure of severity; The Composite Burn Index, and remote sensing of severity, the Normalized Burn Index. ," in FIREMON: Fire Effects Monitoring and Inventory System, D. Lutes, R. Keane, J. Caratti, C. Key, N. Benson, S. Sutherland, and L. Gangi, eds. (Rocky Mountains Research Station, USDA Forest Service, Fort Collins, CO, USA, 2005), pp. 1-51.
68. P. J. Zarco-Tejada, J. R. Miller, T. L. Noland, G. H. Mohammed, and P. H. Sampson, "Scaling-up and model inversion methods with narrowband optical indices for chlorophyll content estimation in closed forest canopies with hyperspectral data," *IEEE Transactions on Geoscience and Remote Sensing* 39, 1491-1507 (2001) [doi:10.1109/36.934080].
69. M. A. Hardisky, V. Klemas, and R. M. Smart, "The influences of soil salinity, growth form, and leaf moisture on the spectral reflectance of *Spartina Alterniflora* canopies," *Photogrammetric Engineering & Remote Sensing* 49, 77-83 (1983).
70. E. M. Barnes, T. R. Clarke, S. E. Richards, P. D. Colaizzi, J. Haberl, M. Kostrzewski, P. Waller, C. Choi, E. Riley, T. Thompson, R. J. Lascano, H. Li, and M. S. Moran, "Coincident Detection of Crop Water Stress, Nitrogen Status and Canopy Density Using Ground Based Multispectral Data," in the Fifth International Conference on Precision Agriculture and Other Resource Management(ASA-CSSA-SSSA, Madison, WI, 2000).

71. A. Dehni, and M. Lounis, "Remote Sensing Techniques for Salt Affected Soil Mapping: Application to the Oran Region of Algeria," *Iswee'11* 33, 188-198 (2012).
72. D. Haboudane, J. R. Miller, N. Tremblay, P. J. Zarco-Tejada, and L. Dextraze, "Integrated narrow-band vegetation indices for prediction of crop chlorophyll content for application to precision agriculture," *Remote Sensing of Environment* 81, 416-426 (2002) [doi:10.1016/S0034-4257(02)00018-4].
73. G. Metternicht, "Vegetation indices derived from high-resolution airborne videography for precision crop management," *International Journal of Remote Sensing* 24, 2855-2877 (2003) [doi:10.1080/01431160210163074].
74. N. H. Broge, and E. Leblanc, "Comparing prediction power and stability of broadband and hyperspectral vegetation indices for estimation of green leaf area index and canopy chlorophyll density," *Remote Sensing of Environment* 76, 156-172 (2001) [doi:10.1016/S0034-4257(00)00197-8].
75. I. Herrmann, A. Pimstein, A. Karnieli, Y. Cohen, V. Alchanatis, and D. J. Bonfil, "LAI assessment of wheat and potato crops by VEN mu S and Sentinel-2 bands," *Remote Sensing of Environment* 115, 2141-2151 (2011) [doi:10.1016/j.rse.2011.04.018].
76. J. Clevers, S. M. De Jong, G. F. Epema, F. D. Van der Meer, W. H. Bakker, A. K. Skidmore, and K. H. Scholte, "Derivation of the red edge index using the MERIS standard band setting," *International Journal of Remote Sensing* 23, 3169-3184 (2002) [doi:10.1080/01431160110104647].
77. J. Penuelas, F. Baret, and I. Filella, "Semi-Empirical Indices to Assess Carotenoids/Chlorophyll-a Ratio from Leaf Spectral Reflectance," *Photosynthetica* 31, 221-230 (1995).
78. R. Fensholt, and I. Sandholt, "Derivation of a shortwave infrared water stress index from MODIS near- and shortwave infrared data in a semiarid environment," *Remote Sensing of Environment* 87, 111-121 (2003) [doi:10.1016/j.rse.2003.07.002].
79. L. Lyburner, P. J. Beggs, and C. R. Jacobson, "Estimation of canopy-average surface-specific leaf area using Landsat TM data," *Photogrammetric Engineering & Remote Sensing* 66, 183-191 (2000).

80. D. Haboudane, N. Tremblay, J. R. Miller, and P. Vigneault, "Remote estimation of crop chlorophyll content using spectral indices derived from hyperspectral data," *IEEE Transactions on Geoscience and Remote Sensing* 46, 423-437 (2008) [doi:10.1109/TGRS.2007.904836].
81. J. W. Rouse, R. H. Haas, J. A. Schell, and D. W. Deering, "Monitoring vegetation systems in the great plains with ERTS," in *Third Earth Resources Technology Satellite-1 Symposium*, S. C. Freden, E. P. Mercanti, and M. A. Becker, eds. (NASA, Washington, D.C, 1974), pp. 309-317.
82. A. A. Gitelson, M. N. Merzlyak, Y. Zur, R. Stark, and U. Gritz, "Non-destructive and remote sensing techniques for estimation of vegetation status," in *Third European Conference on Precision Agriculture*(Montpellier, France, 2001), pp. 301-306.
83. A. A. Gitelson, Y. J. Kaufman, R. Stark, and D. Rundquist, "Novel algorithms for remote estimation of vegetation fraction," *Remote Sensing of Environment* 80, 76-87 (2002) [doi:10.1016/S0034-4257(01)00289-9].
84. A. A. Gitelson, "Wide dynamic range vegetation index for remote quantification of biophysical characteristics of vegetation," *Journal of Plant Physiology* 161, 165-173 (2004) [doi:10.1078/0176-1617-01176].
85. T. Hastie, R. Tibshirani, and J. Friedman, *The Elements of Statistical Learning: Data Mining, Inference, and Prediction. Second Edition* (Springer-Verlag New York, 2009)
86. C. Cortes, and V. Vapnik, "Support-vector networks," *Machine Learning* 20, 273-297 (1995).
87. L. Breiman, "Random forests," *Machine Learning* 45, 5-32 (2001).
88. J. Bergstra, and Y. Bengio, "Random search for hyper-parameter optimization," *Journal of Machine Learning Research* 13, 281-305 (2012).
89. M. J. van der Laan, E. C. Polley, and A. E. Hubbard, "Super learner," *Statistical Applications in Genetics and Molecular Biology* 6 (2007) [doi:10.2202/1544-6115.1309].
90. P. Cortez, and M. J. Embrechts, "Using sensitivity analysis and visualization techniques to open black box data mining models," *Information Sciences* 225, 1-17 (2013) [doi:10.1016/j.ins.2012.10.039].

91. R. Pontius, and M. Millones, "Death to Kappa: birth of quantity disagreement and allocation disagreement for accuracy assessment," *International Journal of Remote Sensing* 32, 4407-4429 (2011) [doi:10.1080/01431161.2011.552923].
92. M. Story, and R. Congalton, "Accuracy assessment: A user's perspective," *Photogrammetric Engineering & Remote Sensing* 52, 397-399 (1986).
93. Q. McNemar, "Note on the sampling error of the difference between correlated proportions or percentages," *Psychometrika* 12, 153-157 (1947).
94. G. M. Foody, "Thematic map comparison: Evaluating the statistical significance of differences in classification accuracy," *Photogrammetric Engineering & Remote Sensing* 70, 627-633 (2004) [doi:10.14358/PERS.70.5.627].

Caption List

Fig. 1 Study area and the distribution of croplands (background map shows Sentinel-2A data obtained on August 11, 2016, R: Band 4, G: Band 3, B: Band 2).




Fig. 2 Crop classification map generated by (a) RF, (b) SVM and (c) SL.




Fig. 3 Relationship between field area and misclassified fields.




Fig. 4 Mean reflectance spectra and standard deviations of each crop.




Table 1 Vegetation indices calculated from Sentinel-2 MSI data.

Table 2 Crop type and number of fields.

Table 3 Classification accuracies of each algorithm.

Table 4 Confusion matrices for (a) RF, (b) SVM and (c) SL.