

UC San Diego

UC San Diego Previously Published Works

Title

Antidepressant Use and Subclinical Measures of Atherosclerosis

Permalink

<https://escholarship.org/uc/item/5c61t3nc>

Journal

Journal of Clinical Psychopharmacology, 36(4)

ISSN

0271-0749

Authors

Camacho, Álvaro
McClelland, Robyn L
Delaney, Joseph A
et al.

Publication Date

2016-08-01

DOI

10.1097/jcp.0000000000000518

Peer reviewed



HHS Public Access

Author manuscript

J Clin Psychopharmacol. Author manuscript; available in PMC 2017 August 01.

Published in final edited form as:

J Clin Psychopharmacol. 2016 August ; 36(4): 340–346. doi:10.1097/JCP.0000000000000518.

Antidepressant use and Subclinical Measures of Atherosclerosis: The Multi-Ethnic Study of Atherosclerosis (MESA)

Álvaro Camacho, MD, MPH¹, Robyn L. McClelland, PhD², Joseph A. Delaney, PhD³, Matthew A. Allison, MD, MPH⁴, Bruce M. Psaty, MD, PhD⁵, Dena E. Rifkin, MD, MS⁶, Stephen R. Rapp, PhD⁷, Moyses Szklo, MD, DrPH⁸, Murray B. Stein, MD, MPH¹, and Michael H. Criqui, MD, MPH⁴

¹Departments of Psychiatry, Family Medicine & Public Health. University of California, San Diego

²Department of Biostatistics. University of Washington, Seattle, WA

³Department of Epidemiology. University of Washington, Seattle, WA

⁴Department of Family Medicine & Public Health. University of California, San Diego

⁵Cardiovascular Health Research Unit. Departments of Medicine, Epidemiology & Health Services. University of Washington. Seattle, WA. Group Health Research Institute. Seattle, WA

⁶Departments of Medicine, Family & Public Health. University of California, San Diego

⁷Department of Psychiatry & Behavioral Medicine. Wake Forest School of Medicine. Winston-Salem, NC

⁸Department of Epidemiology. Johns Hopkins Bloomberg School of Public Health. Baltimore, MD

Abstract

Background—Antidepressants are commonly prescribed medications used in primary care. The cardiovascular safety profile of anti-depressant medications, in terms of subclinical atherosclerosis, is under-examined.

Methods—6,814 participants in the Multi-Ethnic Study of Atherosclerosis (MESA) were examined. At baseline, the mean age was 62 years with four race/ethnic groups represented: European-(38%), Hispanic-(23%), African-(28%) and Chinese-Americans (11%). Antidepressants were sub-grouped as serotonin reuptake inhibitors (SSRIs), tricyclic antidepressants (TCAs), serotonin-norepinephrine reuptake inhibitors (SNRIs), and “other” (bupropion, nefazodone, trazodone, mirtazapine). After adjusting for potential confounders, we estimated the association between antidepressant use and the following measures of subclinical atherosclerosis: coronary artery calcium (CAC), the ankle-brachial index (ABI), and carotid intima-media thickness (cIMT), both cross-sectionally and prospectively.

Corresponding Author: Alvaro Camacho, MD, MPH, Address: 2417 Marshall Rd, Ste 4, Imperial, CA. 92251, acamacho@ucsd.edu. Fax: 1-760-355-2596, Phone: 1-760-791-4018.

Author Disclosure Information: The authors declare no conflicts of interest.

Results—A total of 324 participants were exposed to SSRIs, 88 to TCAs, 41 to SNRIs and 123 to “other” antidepressants. For CAC incidence, the fully-adjusted longitudinal analyses revealed no consistent associations with SSRIs[RR: 0.99(0.71, 1.37)], SNRIs[RR=0.49(0.13, 1.86)], TCAs[RR=0.94(0.50, 1.77)], other antidepressant [RR=0.87(0.73, 1.03)] exposure and subclinical disease. Similar null results were obtained in the cross-sectional and longitudinal exposure to antidepressants with changes in baseline CAC>0, ABI and cIMT.

Conclusions—The results of the current study do not support an association between antidepressants and subclinical atherosclerosis.

Keywords

Multi-Ethnic Study of Atherosclerosis; antidepressant medication; sub-clinical disease; coronary artery calcium

Introduction

Antidepressants are among the most commonly prescribed medications given by primary care providers. Antidepressants are classified based on their postulated mechanism of action: serotonin reuptake inhibitors (SSRIs), serotonin-norepinephrine reuptake inhibitors (SNRIs), tricyclic antidepressants (TCAs), and other antidepressants with mixed dopaminergic and serotonergic modulation such as bupropion, trazodone and mirtazapine, among others. In general, and especially serotonin reuptake inhibitors (SSRIs), these medications have been associated with a reduction of major heart disease events and are considered to be a safe treatment for patients with known atherosclerotic risk factors. Conversely, there is evidence that TCAs and tetracyclic antidepressants like mirtazapine have been associated with cardiotoxic effects that affect conduction and rhythm. There are also reports that link SSRIs with increased risk of hemorrhagic stroke among postmenopausal women. Furthermore, open-label trials of 12 weeks duration and a meta-analysis have reported an association of exposure to SNRIs with increased blood pressure compared to non-SNRIs users. Even though such studies have suggested that antidepressants, especially SSRIs, are not associated with higher risk of atherosclerotic disease, other studies have reported an association between antidepressants and CVD, especially sudden cardiac death. To date, little is known about the association of different classes of antidepressants with measures of subclinical atherosclerotic disease in patients without a history of cardiovascular events.

Subclinical atherosclerosis is associated with incident CVD and is considered to be a chronic progressive inflammatory disease. Risk factors thought to contribute to the development of atherosclerosis include genetics, diabetes, obesity, smoking, dyslipidemia, physical inactivity, systematic inflammation, and negative emotional states such as depression and anxiety. Subclinical markers of atherosclerosis include coronary calcium (CAC), the ankle-brachial index (ABI) and carotid intima-media thickness (cIMT) [among others] which are measured with computed tomography, a hand-held Doppler and B-mode ultrasound respectively.

Since there is paucity in the literature, and to better understand the safety profile of antidepressants for sub-clinical CVD, we conducted a longitudinal study among a diverse cohort of participants free from cardiovascular disease at baseline and with repeat measures of subclinical measures of atherosclerosis at regular intervals. In this study we tested the hypothesis that individuals exposed to different classes of antidepressants would show a protective association with different measures of subclinical atherosclerosis as measured by CAC, ABI and cIMT.

Methods

Study Population

The design and goals of the MESA have been described in previous literature. Briefly, the MESA is a multi-site cohort study that included 6,814 participants with the principal objective of identifying risk factors for the progression of subclinical atherosclerosis to clinical CVD. The MESA cohort includes men and women ages 45-84 years recruited in 2000-2002 from 6 field centers (Columbia University, New York; Johns Hopkins University, Baltimore; Northwestern University, Chicago; University of California, Los Angeles; University of Minnesota, Minneapolis-St. Paul; and Wake Forest University, Winston Salem) and followed for approximately 10 years. All participants were free from known CVD at baseline. Approximately 38% of the participants are European-American, 28% African-American, 23% Hispanic-American, and 11% Chinese-American. All field centers and the data coordinating center had independent institutional review board approval.

Data Collection

Standardized questionnaires were used at each clinic visit to collect demographics, ethnicity, educational attainment, income and health insurance status. Information about medications, and antidepressant medications in particular, was collected through a previously validated medication inventory approach during each examination. The medication inventory approach is more accurate than self-report for assessing baseline exposure to medications in a cohort study. The use of antidepressant medications was classified according to their pharmacological properties into SSRIs, SNRIs, TCAs, and "other" antidepressants (bupropion, nefazodone, trazodone, mirtazapine). A total of 501 (7.3%) participants reported taking anti-depressant medications at baseline. These are standard categories and are considered exhaustive within the current classification of antidepressants. Concordance use of antidepressant was also determined between baseline and follow-up visits. Between the first and second MESA exams, on average 18 months apart, the concordance was 79% for non-TCA use and 65% for TCA use, indicating reasonably consistent antidepressant use over time.

Covariates included systolic blood pressure, body mass index (kg/m^2) (BMI), total cholesterol, HDL cholesterol, C-reactive protein, interleukin-6 and fibrinogen. Psychologic variables included depressive symptom severity as reported by the Center for Epidemiologic Studies Depression scale (CES-D) and anxiety state reported by the Spielberger State-Trait Anxiety Inventory (STAI). Behavioral variables included smoking and alcohol use (never, former, current), and self-reported physical activity. Additional covariates included the

presence of diabetes defined by the 2003 American Diabetes Association's criteria of fasting plasma glucose >126 mg/dl or taking medication for diabetes, lipid-lowering medications, anti-hypertensive medications and aspirin use.

Measures of Subclinical Atherosclerosis

Coronary Artery Calcium (CAC)—CAC was measured using electron-beam computed tomography or multi-detector computed tomography. CAC is a strong predictor of coronary heart disease and provides predictive information for cardiovascular events. At Exam 1 each participant was scanned twice consecutively and the results averaged. At Exam 5 only one scan was performed. Scans were read independently at a centralized reading center (22,23). The amount of calcium was quantified using the Agatston scoring method (24). Calcium scores were adjusted using a standard calcium phantom (25). CAC presence was defined as an Agatston score >0.

Ankle-Brachial Index (ABI)—The ABI was measured at a number of MESA exams, including both exam 1 and 5, as the ratio of the systolic blood pressure in the ankle to that in the arm, as described by previous MESA reports. The participant's ABI was the lower of the left and right legs. Participants with abnormal high ABI >1.40 in either leg were excluded from the analyses because of the possibility of stiff ankle arteries that may lead to an artificially elevated systolic blood pressure. A low-ABI was defined as <0.9, while normal was 0.90-1.40. A low-ABI has been recognized as a good indicator of systemic atherosclerosis and preclinical CVD as well as a good predictor of cardiovascular disease.

Carotid Intima-Media Thickness (cIMT)—At the baseline exam and 10 years later (at exam 5), B-mode ultrasound was used to image the near and far walls of the left distal common carotid artery using a Logiq-700 ultrasound system (General Electric Medical Systems, 13-MHz transducer). The literature has described that one standard deviation increase in cIMT is significantly associated with myocardial infarction and stroke, establishing that high measures of cIMT a good predictor cardiovascular events. Images were read at the University of Wisconsin, and the exact position of the measures at follow-up was determined using anatomical landmarks. All readings were done centrally at the University of Wisconsin and blinded to participant characteristics.

Statistical Analyses

Continuous variables were described by means/standard deviations (SD) and categorical variables were described as frequencies/percentages where applicable (Table 1). Both unadjusted and adjusted analyses were performed for all estimates of association between antidepressant medications and subclinical CVD.

After initial unadjusted analysis, we used a staged model approach with model 1 controlling for demographics only. Model 2 additionally controlled for biological, psychological and behavioral covariates. Model 3 was fully adjusted, additionally controlling for medications and comorbidities. Absolute differences and prevalence ratios (PR) were estimated by linear and prevalence ratio regression models, respectively to determine the effect size of

antidepressant exposure with continuous and dichotomous measures of subclinical atherosclerosis.

We also examined the longitudinal association of antidepressant use with subclinical measures of atherosclerosis. The longitudinal association was determined using a generalized linear model to determine the association of antidepressant exposure at baseline with measures of subclinical atherosclerosis over a 10 year period of study follow-up. Relative risk was calculated for CAC incidence and linear regression estimates were reported for change in CAC, in continuous ABI, and in changes in left distal common carotid mean IMT.

Results

The mean age of participants was 62.1 (10.2) and 56% were women (Table 1). The mean CES-D score was 7.5 (7.5) and the mean STAI state score was 15.8 (4.5). Of the 501 participants taking anti-depressants at baseline, 56% (324) reported taking a SSRI, 15% (88) reported taking a TCA, 7% (41) reported taking a SNRI and 21% (123) reporting taking “other” antidepressant (bupropion, nefazodone, trazodone and/or mirtazapine). Anti-hypertensive medications were the most frequently used concomitant medication in our sample. Thirty nine percent of SSRI users, 37% of SNRI users, 50% of TCA users, and 38% of users of “other” antidepressant were also taking an antihypertensive medication. Table 1 summarizes these findings.

Additional matched controlled analyses between those with significant (CES-D>16) and normal depressive symptoms Vs. those exposed and not exposed to antidepressants, did not show a significant difference between the mean scores of the different measures of subclinical disease. Table 2 summarized this analyses.

Coronary Artery Calcium (CAC)

Compared to non-antidepressant medication users, in fully adjusted models, there were no significant cross-sectional associations between SSRIs, TCAs, SNRIs and “other” antidepressant users with the presence of CAC (Table 2). Similarly, there was no significant association between antidepressant medications and CAC measured as a continuous variable [$\log(\text{CAC})$] (Table 3).

In the longitudinal analyses, in fully adjusted models, there was no significant relationship between exposure to SSRIs, TCAs, SNRIs and “other” antidepressants with incident CAC (change from CAC=0 at baseline) (Table 4). Compared to non-antidepressant users, there was also no significant relationship between exposure to antidepressants and changes in CAC>0 at baseline (Table 4).

Ankle Brachial Index (ABI)

In the fully adjusted model there was no significant association between exposure to SSRIs, TCAs and SNRIs with the ABI as a continuous variable (Table 3), but there was a significant positive association between “other” antidepressant exposure ($\beta=0.023$;95%CI:0.004,0.04) with continuous ABI measures (Table 3).

In longitudinal analyses (Table 4), the fully adjusted model showed no association between “other” antidepressant exposure and change in ABI from baseline over the 10-year observation period. There were also no significant associations between SSRIs, TCAs and SNRIs and change in ABI during the 10-year observation period. Table 4 summarizes these findings.

Carotid Intima-Media Thickness (cIMT)

The estimates of the cross-sectional association in the fully adjusted models were non-significant for the associations between SSRIs, TCAs, SNRIs or for “other” antidepressants and the left distal common carotid mean IMT compared to non-antidepressant medication use (Table 3).

Similarly, the longitudinal analyses in the fully adjusted models (Table 4) showed non-significant findings on the associations between SSRIs, TCAs, SNRIs and “other” antidepressant medication with left distal common carotid mean IMT compared to non-antidepressant medication use.

Discussion

In this longitudinal study of MESA participants free from clinical cardiovascular disease at baseline, there were no significant associations between antidepressant use and measures of subclinical atherosclerosis. The single, cross-sectional association between “other” antidepressant use and ABI was compatible with chance given that we tested 4 medication classes, 2 time-points, and 3 measures of sub-clinical CVD (24 tests). Furthermore, the longitudinal association between these two variables was non-significant, leading us to presume that it was not a real association.

To our knowledge, this is the first study that included longitudinal analyses of different antidepressants that are commonly used in primary care settings with subclinical measures of atherosclerosis. A previous report from the Look AHEAD clinical trial, which studied the long-term health outcomes of a comprehensive weight loss intervention among patients with diabetes and obesity, found a significant cross-sectional association of antidepressant use with low-ABI that was non-significant after controlling for multiple covariates. The association of antidepressants with low-ABI was not reported in the longitudinal analysis of the Look AHEAD clinical trial. Additionally, in the Look AHEAD study the association between different classes of antidepressants and ABI measures was not reported.

Depressive and anxiety disorders are well-described risk factors for the development of CVD. In this study we controlled for symptoms of depression and anxiety, yet the scales that were used (CES-D and STAI) are not frequently used in clinical practice and participants in the MESA cohort were not assessed with structure clinical interviews to discern a diagnosis of a major depressive or anxiety disorder. Even though, cross-sectional studies have demonstrated an association between depressive and anxiety disorders with subclinical atherosclerosis as measured by the ABI, our manuscript was not intended to study the association of depressive symptoms with subclinical measures of atherosclerosis. To what extent the use of antidepressants might protect against the development of

atherosclerosis remains to be further examined. The literature has reported an association between antidepressant exposure and cardiovascular events such as sudden cardiac death. In our study, we did not test for the association of antidepressant use and cardiovascular events since our main focus was to test the association of antidepressants with subclinical measures of atherosclerosis. Also, our study did not have the power to test for the association of antidepressant exposure and cardiovascular events.

Studies in rodents have shown that SSRIs reduce atherosclerotic changes in the endothelium as measured by decreased expression in cell adhesion molecules (VCAM-1). In humans, a randomized clinical trial demonstrated that patients with congestive heart failure taking sertraline, an SSRI, had a reduction in the expression of cell endothelial adhesion molecules (VCAM and ICAM) after a 3 month exposure compared to patients receiving placebo indicating a possible anti-inflammatory effect. On the other hand, a cross-sectional population study from the Netherlands found no association between SSRIs, TCAs or “other” antidepressants with measures of ABI. Our longitudinal study further complements the literature by demonstrating a null relationship between SSRIs, SNRIs, TCAs and “other” antidepressants not only with measures of ABI, but also measures of CAC and cIMT over a 10-year observational period.

Our study has some limitations. We are not able to examine the association between different doses of antidepressants and measures of subclinical atherosclerosis. Additionally, we did not have information of pre-baseline exposure to antidepressants to have a continuous parameter for temporal exposure, that means we did not have duration of exposure prior to baseline assessments. Antidepressant exposure was too rare to estimate dose or agent specific associations. Because of the prevalence use of antidepressants was low, the power to detect associations is accordingly limited. The study was, in addition, observational and thus residual confounding in our estimates is possible.

However, this study has many strengths including: longitudinal data, presence of multiple classes of antidepressants, having gold-standard measures of subclinical atherosclerosis, and the multi-ethnic nature of our sample. Additionally, this study could adjust for a broad range of potential confounders, including a number of covariates that may be important in this context.

In conclusion, we found no significant associations between different classes of antidepressants and any measure of subclinical atherosclerosis. Clinical trials are needed to demonstrate any definitive statement of medication safety, but such trials seem unlikely given the current evidence. However, given its longitudinal component, this study does provide a moderate to high level of evidence that these medications are not strongly associated with subclinical atherosclerosis.

Acknowledgments

This research was supported by contracts N01-HC-95159, N01-HC-95160, N01-HC-95161, N01-HC-95162, N01-HC-95163, N01-HC-95164, N01-HC-95165, N01-HC-95166, N01-HC-95167, N01-HC-95168 and N01-HC-95169 from the National Heart, Lung, and Blood Institute and by grants UL1-TR-000040 and UL1-TR-001079 from NCCRR. The authors thank the other investigators, the staff, and the participants of the MESA study for their valuable contributions. A full list of participating MESA investigators and institutions can be found at <http://>

www.mesa-nhlbi.org. This publication was developed under a STAR research assistance agreement, No. RD831697 (MESA Air), awarded by the U.S Environmental protection Agency. It has not been formally reviewed by the EPA. The views expressed in this document are solely those of the authors and the EPA does not endorse any products or commercial services mentioned in this publication. There was also support from the T32HL07989 UCSD Integrated Cardiovascular Epidemiology Fellowship and R25HL105430 Mentoring Latinos in Health Disparities Research from the National Heart, Lung and Blood Institute (NHLBI).

References

1. Pincus H, Tanielian T, Marcus SC, Olfson M, Zarin DA, Thompson J, Magno Zito J. Prescribing trends in psychotropic medications: Primary care, psychiatry, and other medical specialties. *JAMA*. 1998; 279:526–531. [PubMed: 9480363]
2. Simon GE. Evidence review: efficacy and effectiveness of antidepressant treatment in primary care. *Gen Hosp Psychiatry*. 2002; 24:213–24. [PubMed: 12100832]
3. Stahl SM, Grady MM, Moret C, Briley M. SNRIs: their pharmacology, clinical efficacy, and tolerability in comparison with other classes of antidepressants. *CNS Spectr*. 2005 Sep; 10(9):732–47. *CNS Spectr*;10:732-47. [PubMed: 16142213]
4. Glassman AH, O'Connor CM, Califf RM, Swedberg K, Schwartz P, Bigger JT Jr, Krishnan KR, van Zyl LT, Swenson JR, Finkel MS, Landau C, Shapiro PA, Pepine CJ, Mardekian J, Harrison WM, Barton D, M M. Sertraline Antidepressant Heart Attack Randomized Trial (SADHEART) Group. Sertraline treatment of major depression in patients with acute MI or unstable angina. *JAMA*. 2002; 288:701–9. [PubMed: 12169073]
5. Scherrer JF, Garfield LD, Lustman PJ, Hauptman PJ, Chrusciel T, Zeringue A, Carney RM, Freedland KE, Bucholz KK, Owen R, Newcomer JW, True WR. Antidepressant Drug Compliance: Reduced Risk of MI and Mortality in Depressed Patients. *Am J Med*. 2011; 124:318–324. [PubMed: 21435422]
6. Gordon NF, Habib A, Salmon RD, Bishop KL, Drimmer A, Reid KS, Wright BS, Faircloth GC, Gordon TL, Franklin BA, Rubenfire M, Gracik T, Sperling LS. Effect of Exercise-Based Cardiac Rehabilitation on Multiple Atherosclerotic Risk Factors in Patients Taking Antidepressant Medication. *Am J Cardiol*. 2013; 111:346–351. [PubMed: 23178050]
7. Cohen HW, Gibson G, Alderman MH. Excess risk of myocardial infarction in patients treated with antidepressant medications: association with use of tricyclic agents. *Am J Med*. 2000; 108:2–8. [PubMed: 11059434]
8. Leonard CE, Bilker WB, Newcomb C, Kimmel SE, Hennessy S. Antidepressants and the Risk of Sudden Cardiac Death and Ventricular Arrhythmia. *Pharmacoepidemiol Drug Saf*. 2011; 20:903–913. [PubMed: 21796718]
9. Smoller JW, Allison M, Cochrane BB, Curb JD, Perlis RH, Robinson JG, Rosal MC, Wenger NK, Wassertheil-Smoller S. Antidepressant Use and Risk of Incident Cardiovascular Morbidity and Mortality Among Postmenopausal Women in the Women's Health Initiative Study. *Arch Intern Med*. 2009; 169:2128–2139. [PubMed: 20008698]
10. Smoller JW. Do antidepressants raise the risk of stroke? *Am J Psychiatry*. 2011; 168:457–9. [PubMed: 21536699]
11. Wu CS, Wang SC, Cheng YC, G SF. Association of cerebrovascular events with antidepressant use: a case cross-over study. *Am J Psychiatry*. 2011; 168:457–9. [PubMed: 21536699]
12. Johnson EM, Whyte E, Mulsant BH, Pollock BG, Weber E, Begley AE, Reynolds CF. Cardiovascular changes associated with venlafaxine in the treatment of late-life depression. *Am J Geriatr Psychiatry*. 2006; 14:796–802. [PubMed: 16943176]
13. Thase ME. Effects of venlafaxine on blood pressure: a meta-analysis of original data from 3744 depressed patients. *J Clin Psychiatry*. 1998; 59:502–8. [PubMed: 9818630]
14. Serebruany VL, Glassman AH, Malinin AI, Nemeroff CB, Musselman DL, van Zyl LT, Finkel MS, Krishnan KR, Gaffney M, Harrison W, Califf RM, O'Connor CM. Sertraline AntiDepressant Heart Attack Randomized Trial Study Group. Platelet/endothelial biomarkers in depressed patients treated with the selective serotonin reuptake inhibitor sertraline after acute coronary events: the Sertraline AntiDepressant Heart Attack Randomized Trial (SADHART) Platelet Substudy. *Circulation*. 2003; 108:939–44. [PubMed: 12912814]

15. Wozniak G, Toska A, Saridi M, Mouzas O. Serotonin reuptake inhibitor antidepressants (SSRIs) against atherosclerosis. *Med Sci Monit.* 2011; 17:RA205–14. [PubMed: 21873959]
16. Narayan SM, Stein MB. Do Depression or Antidepressants Increase Cardiovascular Mortality? The Absence of Proof Might Be More Important Than the Proof of Absence*. *J Am Coll Cardiol.* 2009; 53:959–961. [PubMed: 19281926]
17. Allison MA, Budoff MJ, Wong ND, Blumenthal RS, Schreiner PJ, Criqui MH. Prevalence of and risk factors for subclinical cardiovascular disease in selected US Hispanic ethnic groups: the Multi-Ethnic Study of Atherosclerosis. *Am J Epidemiol.* 2008; 167:962–9. [PubMed: 18283034]
18. Toth PP. Subclinical atherosclerosis: what it is, what it means and what we can do about it. *Int J Clin Pract.* 2008; 62:1246–54. [PubMed: 18564201]
19. Tiemeier H, van Dijk W, Hofman A, Witteman JC, Stijnen T, Breteler MM. Relationship between atherosclerosis and late-life depression: the Rotterdam Study. *Arch Gen Psychiatry.* 2004; 61:369–76. [PubMed: 15066895]
20. Bertoni AG, Goff DC Jr, D'Agostino RB Jr, Liu K, Hundley WG, Lima JA, Polak JF, Saad MF, Szklo M, Tracy RP, Siscovick DS. Diabetic cardiomyopathy and subclinical cardiovascular disease: the Multi-Ethnic Study of Atherosclerosis (MESA). *Diabetes Care.* 29:588–94. [PubMed: 16505511]
21. Timóteo AT, Toste A, Araújo A, Ferreira RC. Expression of subclinical atherosclerosis for different cardiovascular risk factors in young populations. *Rev Port Cardiol.* 2010; 29:1181–90. [PubMed: 21066970]
22. Bild DE, Bluemke DA, Burke GL, Detrano R, Diez Roux AV, Folsom AR, Greenland P, Jacobs DR Jr, Kronmal R, Liu K, Nelson JC, O'Leary D, Saad MF, Shea S, Szklo M, Tracy RP. Multi-Ethnic Study of Atherosclerosis: Objectives and Design. *Am J Epidemiol.* 2002; 156:871–881. [PubMed: 12397006]
23. Roy B, Diez-Roux AV, Seeman T, Ranjit N, Shea S, Cushman M. Association of optimism and pessimism with inflammation and hemostasis in the Multi-Ethnic Study of Atherosclerosis (MESA). *Psychosom Med.* 2010; 72:134–40. [PubMed: 20100888]
24. Psaty BM, Lee M, Savage PJ, Rutan GH, German PS, Lyles M. HSCRG. Assessing the use of medications in the elderly: methods and initial experience in the Cardiovascular Health Study. *The Cardiovascular. J Clin Epidemiol.* 1992; 45:683–92. [PubMed: 1607909]
25. Delaney JA, Oddson BE, McClelland RL, Psaty BM. Estimating ethnic differences in self-reported new use of antidepressant medications: results from the Multi-Ethnic Study of Atherosclerosis. *Pharmacoepidemiol Drug Saf.* 2009; 18:545–53. [PubMed: 19399919]
26. Delaney JA, Oddson BE, Kramer H, Shea S, Psaty BM, McClelland RL. Baseline depressive symptoms are not associated with clinically important levels of incident hypertension during two years of follow-up: the multi-ethnic study of atherosclerosis. *Hypertension.* 2010; 55:408–14. [PubMed: 20065156]
27. Cipriani A, Furukawa TA, Salanti G, Geddes JR, Higgins JPT, Churchill R, Watanabe N, Nakagawa A, Omori IM, McGuire H, Tansella M, Barbui C. Comparative efficacy and acceptability of 12 new-generation antidepressants: a multiple-treatments meta-analysis. *The Lancet.* 373:746–758.
28. Stahl SM. Enhancing outcomes from major depression: using antidepressant combination therapies with multifunctional pharmacologic mechanisms from the initiation of treatment. *CNS Spectr.* 2010; 15:79–94. [PubMed: 20414154]
29. American Diabetes Association. Position statement: diagnosis and classification of diabetes mellitus. *Diabetes Care.* 2008; 31:S55–S60. [PubMed: 18165338]
30. Detrano R, Guerci AD, Carr JJ, Bild DE, Burke G, Folsom AR, Liu K, Shea S, Szklo M, Bluemke DA, O'Leary DH, Tracy R, Watson K, Wong ND, Kronmal RA. Coronary Calcium as a Predictor of Coronary Events in Four Racial or Ethnic Groups. *New England Journal of Medicine.* 2008; 358:1336–1345. [PubMed: 18367736]
31. Criqui MH, McClelland RL, McDermott MM, Allison MA, Blumenthal RS, Aboyans V, Ix JH, Burke GL, Liu K, Shea S. The ankle-brachial index and incident cardiovascular events in the MESA (Multi-Ethnic Study of Atherosclerosis). *J Am Coll Cardiol.* 2010; 56:1506–12. [PubMed: 20951328]

32. Shadman R, Criqui MH, Bundens WP, Fronck A, Denenberg JO, Gamst AC, McDermott MM. Subclavian artery stenosis: prevalence, risk factors, and association with cardiovascular diseases. *J Am Coll Cardiol*. 2004; 44:618–23. [PubMed: 15358030]
33. Aboyans V, McClelland RL, Allison MA, McDermott MM, Blumenthal RS, Macura K, Criqui MH. Lower extremity peripheral artery disease in the absence of traditional risk factors. The Multi-Ethnic Study of Atherosclerosis. *Atherosclerosis*. 2011; 214:169–73. [PubMed: 21067754]
34. Criqui MH, Vargas V, Denenberg JO, Ho E, Allison M, Langer RD, Gamst A, Bundens WP, Fronck A. Ethnicity and peripheral arterial disease: the San Diego Population Study. *Circulation*. 2005; 112:2703–7. [PubMed: 16246968]
35. Lijmer JG, Hunink MG, van den Dungen JJ, Loonstra J, Smit AJ. ROC analysis of noninvasive tests for peripheral arterial disease. *Ultrasound Med Biol*. 22:391–8. 196. [PubMed: 8795165]
36. Newman AB, Siscovick DS, Manolio TA, Polak J, Fried LP, Borhani NO. Ankle-arm index as a marker of atherosclerosis in the Cardiovascular Health Study. Cardiovascular Health Study (CHS) Collaborative Research Group. *Circulation*. 1993; 88
37. van den Oord SCH, Sijbrands EJG, ten Kate GL, van Klaveren D, van Domburg RT, van der Steen AFW, Schinkel AFL. Carotid intima-media thickness for cardiovascular risk assessment: Systematic review and meta-analysis. *Atherosclerosis*. 2013; 228:1–11. [PubMed: 23395523]
38. Folsom AR, Kronmal RA, Detrano RC, et al. Coronary artery calcification compared with carotid intima-media thickness in the prediction of cardiovascular disease incidence: The multi-ethnic study of atherosclerosis (mesa). *Archives of Internal Medicine*. 2008; 168:1333–1339. [PubMed: 18574091]
39. Gepner AD, Young R, Delaney JC, et al. A Comparison of Coronary Artery Calcium Presence, Carotid Plaque Presence, and Carotid Intima-Media Thickness for Cardiovascular Disease Prediction in the Multi-Ethnic Study of Atherosclerosis (MESA). *Circ Cardiovasc Imaging*. 2015; 8(1)doi: 10.1161/CIRCIMAGING.114.002262
40. Rubin RR, Gaussoin SA, Peyrot M, DiLillo V, Miller K, Wadden TA, West DS, Wing RR, Knowler WC. Look AHEAD Research Group. Cardiovascular disease risk factors, depression symptoms and antidepressant medicine use in the Look AHEAD (Action for Health in Diabetes) clinical trial of weight loss in diabetes. *Diabetologia*. 2010; 53:1581–9. [PubMed: 20422396]
41. Rubin RR, Peyrot M, Gaussoin SA, Espeland MA, Williamson D, Faulconbridge LF, Wadden TA, Ewing L, Safford M, Evans-Hudnall G, Wing RR, Knowler WC. Look AHEAD Research Group. Four-year analysis of cardiovascular disease risk factors, depression symptoms, and antidepressant medicine use in the Look AHEAD (Action for Health in Diabetes) clinical trial of weight loss in diabetes. *Diabetes Care*. 2013; 36:1088–94. [PubMed: 23359362]
42. Wassertheil-Smoller S, Shumaker S, Ockene J, Talavera GA, Greenland P, Cochrane B, Robbins J, Aragaki A, Dunbar-Jacob J. Depression and cardiovascular sequelae in postmenopausal women: The women's health initiative study (WHI). *Arch Intern Med*. 2004; 164:289–298. [PubMed: 14769624]
43. Roest AM, Martens EJ, de Jonge P, Denollet J. Anxiety and Risk of Incident Coronary Heart Disease: A Meta-Analysis. *J Am Coll Cardiol*. 2010; 56:38–46. [PubMed: 20620715]
44. Seldenrijk A, Vogelzangs N, van Hout HPJ, van Marwijk HWJ, Diamant M, Penninx BWJH. Depressive and anxiety disorders and risk of subclinical atherosclerosis: Findings from the Netherlands Study of Depression and Anxiety (NESDA). *J Psychosom Res*. 2010; 69:203–210. [PubMed: 20624520]
45. Unis A, Abdelbary A, Hamza M. Comparison of the effects of escitalopram and atorvastatin on diet-induced atherosclerosis in rats. *Can J Physiol Pharmacol*. 2013; 92:226–233. [PubMed: 24593787]
46. Lekakis J, Ikonomidis I, Papoutsis Z, Moutsatsou P, Nikolaou M, Parissis J, Kremastinos DT. Selective serotonin re-uptake inhibitors decrease the cytokine-induced endothelial adhesion molecule expression, the endothelial adhesiveness to monocytes and the circulating levels of vascular adhesion molecules. *Int J Cardiol*. 2010; 139:150–158. [PubMed: 19004511]
47. Arnaud C, Burger F, Steffens S, Veillard NR, Nguyen TH, Trono D, Mach F. Statins reduce interleukin-6-induced C-reactive protein in human hepatocytes: new evidence for direct

antiinflammatory effects of statins. *Arterioscler Thromb Vasc Biol.* 2005; 25:1231–6. [PubMed: 15790934]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

Participant characteristics by baseline use of antidepressant medications. Data from the Multi-Ethnic Study of Atherosclerosis (MESA).

	SSRIs (n=324)	TCAs (n=88)	SNRIs (n=41)	Other (n=123)	No Antidepressant (n=6298)
Age (years)	61	62	60	57	62
Male	30%	20%	34%	41%	49%
Race/Ethnicity					
Caucasian	63%	58%	78%	74%	36%
Chinese	5%	3%	2%	1%	12%
African-American	13%	20%	2%	12%	29%
Hispanic	19%	18%	17%	13%	22%
Education					
<High School	11%	15%	7%	7%	19%
High School	45%	56%	51%	47%	47%
College	23%	11%	10%	22%	17%
Graduate School	21%	18%	32%	25%	18%
Household Income					
<\$25K	30%	37%	21%	25%	32%
\$25K and < \$50	33%	31%	21%	30%	29%
\$50K and < \$100	24%	29%	38%	28%	26%
\$100K	13%	2%	21%	17%	14%
No Health Insurance	4%	5%	12%	5%	9%
BMI (kg/m ²)	29	30	30	29	28
Total Cholesterol (mg/dL)	200	192	193	191	194
HDL Cholesterol (mg/dL)	54	53	52	52	51
Systolic BP (mmHG)	124	124	130	121	127
C-reactive protein (mg/dl)	4.9	6.1	4.4	4.2	3.7
Fibrinogen Antigen (mg/dl)	346	360	342	337	347
IL-6 (pg/mL)	1.67	1.95	1.76	1.54	1.54

	SSRIs (n=324)	TCAs (n=88)	SNRIs (n=41)	Other (n=123)	No Antidepressant (n=6298)
Depression Score (CES-D)	11	10	11	12	7
Anxiety Score (STAI)	18	17	18	19	16
Smoking: Current	12%	19%	17%	20%	13%
Smoking: Former	42%	40%	44%	43%	36%
Alcohol Consumption: Current	57%	50%	66%	50%	55%
Alcohol Consumption: Former	28%	27%	24%	39%	24%
Intentional Exercise (met-min/wk)	1680	1312	1119	1474	1548
Diabetes	13%	16%	10%	8%	13%
Lipid Lowering Medications	23%	25%	34%	23%	16%
Antihypertensive Medications	39%	50%	37%	38%	37%
Aspirin Use	26%	24%	39%	32%	25%
Coronary Artery Calcium					
Prevalent CAC>0	46%	44%	51%	37%	50%
incident CAC>0	28%	23%	14%	26%	26%
ABI Baseline	1.12	1.12	1.11	1.14	1.11
Change E1 to E5	-0.014	-0.016	-0.024	-0.005	-0.007
Carotid IMT (mm) Baseline	0.74	0.73	0.67	0.70	0.76
Change E1 to E5	0.11	0.13	0.10	0.10	0.11

IL-6= Interleukin-6; CES-D: Center for Epidemiologic Studies Depression Scale; STAI= Spielberger Trait Anxiety; CCA=Common Carotid Artery; IMT=Intima-Media Thickness; ABI=Ankle-Brachial Index.

Table 2
Mean Differences of Subclinical Measures of Atherosclerosis among participants exposed to antidepressants and CES-D symptoms

CES-D > 16	Sub-Clinical Measure of Atherosclerosis	N	Mean	SD	Minimum	Maximum
No Antidepressants & Not depressed	CCA IMT (mm)	5578	0.874	0.195	0.400	2.453
	ANKLE-BRACHIAL INDEX	5592	1.114	0.122	0.306	1.983
	CAC Score	5655	150.238	424.993	0.000	6315.910
No Antidepressants & Depressed	CCA IMT (mm)	635	0.858	0.192	0.470	1.858
	ANKLE-BRACHIAL INDEX	632	1.104	0.115	0.607	1.888
	CAC Score	641	127.418	394.309	0.000	4506.495
On Antidepressants & Not depressed	CCA IMT (mm)	379	0.845	0.183	0.475	1.625
	ANKLE-BRACHIAL INDEX	377	1.122	0.105	0.712	1.455
	CAC Score	384	131.011	376.286	0.000	2333.970
On Antidepressants & Depressed	CCA IMT (mm)	119	0.852	0.194	0.530	1.528
	ANKLE-BRACHIAL INDEX	119	1.108	0.125	0.452	1.500
	CAC Score	119	109.897	292.459	0.000	1697.730

CES-D: Center of Epidemiologic Studies Depression Scale; CCA-IMT: Common Carotid Artery-Intima Media Thickness; CAC: Coronary Artery Calcification; SD: Standard Deviation

Table 3
Cross-Sectional Association of antidepressants with subclinical atherosclerosis

Coronary Artery Calcium Prevalence (CAC>0)						
	SSRIs PR (95% CI)	TCAs PR (95% CI)	SNRIs PR (95% CI)	Other PR (95% CI)	Non Users	
Model 1	1.05 (0.95, 1.15)	0.99 (0.82, 1.20)	1.05 (0.82, 1.34)	0.87 (0.73, 1.04)	Ref	
Model 2	1.04 (0.94, 1.14)	0.99 (0.83, 1.19)	1.03 (0.86, 1.25)	0.88 (0.74, 1.05)	Ref	
Model 3	1.01 (0.92, 1.11)	0.95 (0.78, 1.14)	0.98 (0.82, 1.17)	0.87 (0.73, 1.03)	Ref	
Coronary Artery Calcium Amount: log(CAC) among those with CAC>0						
	SSRIs Coef (95% CI)	TCAs Coef (95% CI)	SNRIs Coef (95% CI)	Other Coef (95% CI)	Non Users	
Model 1	-0.04 (-0.33, 0.25)	0.20 (-0.36, 0.75)	-0.02 (-0.80, 0.75)	-0.09 (-0.62, 0.43)	Ref	
Model 2	-0.10 (-0.39, 0.20)	0.23 (-0.32, 0.78)	-0.19 (-0.96, 0.58)	-0.07 (-0.59, 0.46)	Ref	
Model 3	-0.17 (-0.47, 0.12)	0.17 (-0.38, 0.72)	-0.28 (-1.05, 0.48)	-0.09 (-0.61, 0.43)	Ref	
Continuous ABI excludes participants with ABI > 1.40						
	SSRIs Coef (95% CI)	TCAs Coef (95% CI)	SNRIs Coef (95% CI)	Other Coef (95% CI)	Non Users	
Model 1	0.005 (-0.01, 0.02)	0.014 (-0.01, 0.04)	-0.018 (-0.05, 0.02)	0.017 (-0.003, 0.04)	Ref	
Model 2	0.007 (-0.01, 0.02)	0.016 (-0.01, 0.04)	-0.013 (-0.05, 0.02)	0.022 (0.003, 0.04) ‡	Ref	
Model 3	0.009 (-0.003, 0.02)	0.019 (-0.004, 0.04)	-0.010 (-0.04, 0.02)	0.023 (0.004, 0.04) ‡	Ref	
Carotid IMT (left distal common carotid mean IMT)						
	SSRIs Coef (95% CI)	TCAs Coef (95% CI)	SNRIs Coef (95% CI)	Other Coef (95% CI)	Non Users	
Model 1	0.011 (-0.02, 0.05)	-0.026 (-0.10, 0.04)	-0.027 (-0.13, 0.08)	0.00 (-0.05, 0.05)	Ref	
Model 2	-0.000 (-0.03, 0.03)	-0.028 (-0.10, 0.04)	-0.035 (-0.14, 0.07)	-0.001 (-0.05, 0.05)	Ref	
Model 3	-0.002 (-0.04, 0.03)	-0.031 (-0.10, 0.04)	-0.038 (-0.14, 0.06)	-0.004 (-0.06, 0.05)	Ref	

Model 1 (Demographics): Age, gender, ethnicity, education, income, health insurance; **Model 2 (Biological, Psychological & Behaviors):** BMI, total cholesterol, HDL cholesterol, systolic blood pressure, CRP, IL-6, Fibrinogen, CES-D, STAI, smoking, alcohol use, and physical activity; **Model 3: Medications and Comorbidities:** Diabetes, lipid-lowering medication, antihypertensives, aspirin use. **CAC:** Coronary calcium. **ABI:** Ankle-brachial index. **IMT:** intima-media thickness. **SSRIs:** Serotonin reuptake inhibitors. **SNRIs:** Tricyclic antidepressants. **TCAs:** Serotonin reuptake inhibitors. **Other=** bupropion, nefazodone, mirtazapine. **PR:** prevalence ratio

Table 4
Longitudinal Association of antidepressants with subclinical atherosclerosis

Coronary Artery Calcium Incidence (CAC>0)						
	SSRIs RR (95% CI)	TCAs RR (95% CI)	SNRIs RR (95% CI)	Other RR (95% CI)	Non Users	
Model 1	1.07 (0.81, 1.42)	0.88 (0.48, 1.64)	0.52 (0.14, 1.94)	1.02 (0.65, 1.60)	Ref	
Model 2	0.99 (0.72, 1.37)	0.93 (0.50, 1.73)	0.49 (0.13, 1.86)	1.11 (0.70, 1.76)	Ref	
Model 3	0.99 (0.71, 1.37)	0.94 (0.50, 1.77)	0.49 (0.13, 1.86)	1.10 (0.69, 1.76)	Ref	
Change in CAC Amount (Agatston; among those with CAC>0 at baseline)						
	SSRIs Coef (95% CI)	TCAs Coef (95% CI)	SNRIs Coef (95% CI)	Other Coef (95% CI)	Non Users	
Model 1	3 (-123, 128)	149 (-95, 394)	-187 (-552, 178)	-1 (-216, 214)	Ref	
Model 2	-36 (-164, 92)	124 (-120, 369)	-258 (-622, 106)	-34 (-249, 181)	Ref	
Model 3	-54 (-178, 70)	137 (-101, 374)	-220 (-574, 133)	-58 (-266, 151)	Ref	
Change in Continuous ABI excludes participants with ABI> 1.40						
	SSRIs Coef (95% CI)	TCAs Coef (95% CI)	SNRIs Coef (95% CI)	Other Coef (95% CI)	Non Users	
Model 1	-0.01 (-0.03, 0.01)	-0.004 (-0.04, 0.03)	-0.01 (-0.06, 0.04)	-0.004 (-0.03, 0.02)	Ref	
Model 2	-0.01 (-0.03, 0.01)	-0.001 (-0.04, 0.03)	-0.01 (-0.06, 0.04)	-0.004 (-0.03, 0.02)	Ref	
Model 3	-0.01 (-0.03, 0.01)	-0.004 (-0.04, 0.03)	-0.01 (-0.06, 0.04)	-0.005 (-0.03, 0.02)	Ref	
Change in Carotid IMT (left distal common carotid mean IMT)						
	SSRIs Coef (95% CI)	TCAs Coef (95% CI)	SNRIs Coef (95% CI)	Other Coef (95% CI)	Non Users	
Model 1	-0.003 (-0.03, 0.03)	0.03 (-0.03, 0.08)	-0.01 (-0.10, 0.08)	-0.002 (-0.05, 0.04)	Ref	
Model 2	-0.007 (-0.04, 0.02)	0.03 (-0.03, 0.09)	-0.01 (-0.10, 0.08)	-0.002 (-0.05, 0.04)	Ref	
Model 3	-0.009 (-0.04, 0.02)	0.03 (-0.03, 0.09)	-0.02 (-0.10, 0.07)	0.001 (-0.04, 0.05)	Ref	

Model 1 (Demographics): Age, gender, ethnicity, education, income, health insurance; **Model 2 (Biological, Psychological & Behaviors):** BMI, total cholesterol, HDL cholesterol, systolic blood pressure, CRP, IL-6, Fibrinogen, CES-D, STAI, smoking, alcohol use, and physical activity; **Model 3 (Medications and Comorbidities):** Diabetes, lipid-lowering medication, antihypertensives, aspirin use. **CAC:** Coronary calcium. **ABI:** Ankle-brachial index. **IMT:** intima-media thickness. **SSRIs:** Serotonin reuptake inhibitors. **SNRIs:** Tricyclic antidepressants. **TCAs:** Tricyclic antidepressants. **Other:** bupropion, nefazadone, trazodone, mirtazapine. **RR:** relative risk

* p<0.05