

UC Irvine

Faculty Publications

Title

Feasibility of reconstructing paleoatmospheric records of selected alkanes, methyl halides, and sulfur gases from Greenland ice cores

Permalink

<https://escholarship.org/uc/item/9wz0z76j>

Journal

Journal of Geophysical Research, 112(D7)

ISSN

0148-0227

Authors

Aydin, M.
Williams, M. B
Saltzman, E. S

Publication Date

2007-04-16

DOI

10.1029/2006JD008027

Supplemental Material

<https://escholarship.org/uc/item/9wz0z76j#supplemental>

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Feasibility of reconstructing paleoatmospheric records of selected alkanes, methyl halides, and sulfur gases from Greenland ice cores

M. Aydin,¹ M. B. Williams,¹ and E. S. Saltzman¹

Received 7 September 2006; revised 29 December 2006; accepted 15 January 2007; published 12 April 2007.

[1] Seven short-lived atmospheric trace gases were measured in 25 ice core samples from Summit, Greenland. Samples were selected from contemporaneous sections of fluid- and dry-drilled ice cores to examine what effects using *n*-butyl acetate as the drill fluid would have on the measurements. The gases include three light alkanes, C₂H₆, C₃H₈, and *n*-C₄H₁₀; two methyl halides, CH₃Cl and CH₃Br; and two sulfur compounds, OCS and CS₂, with gas ages from 125 to 325 years before present. Alkane levels are comparable to measurements in modern Arctic air, although C₂H₆ exhibits greater variability than expected compared with C₃H₈ and *n*-C₄H₁₀. These results are not consistent with the idea that the alkanes are primarily of anthropogenic origin, suggesting that the ice cores may not truly record a paleoatmospheric signal with respect to these gases. The CH₃Br results are consistent with previous observations of “excess” CH₃Br in Greenland firn air. In situ production processes appear to overwhelm the paleoatmospheric signal of this gas. CH₃Cl exhibits the same effect to a lesser extent. OCS levels are similar to those in Antarctic ice cores and appear to reflect paleoatmospheric levels. CS₂ results are similar to the limited database of modern atmospheric measurements. Only C₃H₈ and *n*-C₄H₁₀ exhibit clear evidence of contamination because of the presence of the drill fluid. The results indicate that it is possible to analyze many trace gases in fluid- and dry-drilled ice samples. However, it appears that in situ production may significantly alter the levels of some trace gases in Greenland ice cores.

Citation: Aydin, M., M. B. Williams, and E. S. Saltzman (2007), Feasibility of reconstructing paleoatmospheric records of selected alkanes, methyl halides, and sulfur gases from Greenland ice cores, *J. Geophys. Res.*, 112, D07312, doi:10.1029/2006JD008027.

1. Introduction

[2] Air bubbles trapped inside ice cores are a valuable archive of the ancient atmosphere. Although paleoatmospheric histories of greenhouse gases like CO₂ and CH₄ have been reconstructed from ice cores, much less information exists about the atmospheric variability of lower abundance trace gases such as halocarbons, hydrocarbons, and sulfur gases that have relatively short atmospheric lifetimes, ranging from days to a few years. In this study, we report measurements of C₂-C₄ alkanes (ethane, C₂H₆; propane, C₃H₈; *n*-butane, *n*-C₄H₁₀), two methyl halides (CH₃Cl and CH₃Br), and two sulfur compounds (OCS and CS₂) in air extracted from Greenland ice cores. Thirty samples were analyzed from B and D cores drilled to depths of 200 and 3000 m, respectively, as part of the Greenland Ice Sheet Project (GISP2) at Summit, Greenland (72.6°N, 38.5°W, 3200 m asl). The shallow B core was dry-drilled, while *n*-butyl acetate (C₆H₁₂O₂) was used as the drilling fluid for the D core. The samples from both cores have contemporaneous gas ages as they were chosen from similar depths. The primary goal of this study is to assess the

feasibility of reconstructing paleoatmospheric levels of short-lived trace gases from dry- and fluid-drilled Greenland ice cores.

[3] There have been two studies of low-molecular-weight alkanes in firn air. C₂H₆ and C₃H₈ results from an Antarctic shallow firn [Kaspers *et al.*, 2004] and C₂H₆, C₃H₈, and C₄H₁₀ measurements in firn air from Summit, Greenland [Worton, 2006], suggest that alkanes are stable in firn and are not prone to in situ artifacts. It has also been shown that there is no photochemical production of alkanes in Greenland snow [Swanson *et al.*, 2002]. The available evidence, albeit limited, suggests that it may be possible to reconstruct a paleoatmospheric record of alkanes from ice core measurements.

[4] Measurements of CH₃Cl in firn air from Greenland and Antarctica suggest that this gas is preserved in the polar firn air from both hemispheres [Butler *et al.*, 1999; Kaspers *et al.*, 2004; Trudinger *et al.*, 2004]. Trudinger *et al.* [2004] argue that subtle discrepancies exist between certain Antarctic firn records, which may implicate in situ chemistry. A 300-year atmospheric history of CH₃Cl was developed based on measurements in Siple Dome ice cores from Antarctica [Aydin *et al.*, 2004]. CH₃Cl has also been measured in a limited number of samples from another Antarctic ice core [Saito *et al.*, 2006]. The results indicate that CH₃Cl is stable in polar ice matrices over centennial timescales.

¹Department of Earth System Science, University of California, Irvine, California, USA.

[5] CH_3Br exhibits different behavior in firm air from the two hemispheres. Antarctic firm air appears to record paleo-atmospheric CH_3Br levels. Firm air profiles from several Antarctic sites show CH_3Br levels increasing through the twentieth century in reasonable agreement with estimates of anthropogenic emissions [Butler *et al.*, 1999; Sturges *et al.*, 2001a; Trudinger *et al.*, 2004]. Antarctic ice core measurements in Siple Dome demonstrate that CH_3Br is stable in Antarctic ice on timescales of hundreds of years and exhibits levels consistent with those in firm air [Saltzman *et al.*, 2004]. In contrast, results from Tunu, Greenland, and Devon Island, Canada, indicate enrichment of CH_3Br in firm air relative to atmospheric levels [Butler *et al.*, 1999; Sturges *et al.*, 2001a]. So far, the enrichment observed in the Northern Hemisphere firm sites has not been explained, but Swanson *et al.* [2002] observed photochemical production of CH_3Br in Greenland surface snow, providing the first direct evidence of possible artifacts because of in situ chemistry.

[6] OCS is the only sulfur gas that has been measured in polar firm air and ice cores. Firm air results from Antarctica and Canada suggest that an atmospheric OCS record is preserved in both hemispheres [Sturges *et al.*, 2001b]. More recently, Antarctic ice core and firm air measurements have been used to reconstruct the atmospheric history of OCS over the last 400 years [Aydin *et al.*, 2002; Montzka *et al.*, 2004]. To date, there are no published measurements of CS_2 in firm air or ice cores.

2. Methods

2.1. Ice Core Dating

[7] At polar glacial sites, the vertical transport of gases in firm is determined primarily by molecular diffusion, gravitational settling, and to a lesser degree by wind pumping and convection [Schwander *et al.*, 1993]. The aging of various gases in the firm occurs at slightly different rates that can be estimated from their respective molecular weights and diffusivities [Schwander *et al.*, 1993, 1997]. Below the bubble close-off depth, all gases age at the same rate, determined by the snow accumulation rate. A gas age at bubble close off is not a true age but rather a mean diffusive age that is always younger than the surrounding ice at close off. Lighter, more diffusive gases have a younger mean age and a narrower gas age distribution profile than heavier, less diffusive ones. In the calculation of the mean gas ages in this study, we determine an ice age-gas age difference (delta age) specific to each gas at pore close-off depth and subtract this delta age from the ice ages through the length of the core.

[8] The ice core samples analyzed in this study are from a depth range of 98.7–137.1 m for GISP2 B and 112.9–145.9 m for GISP2 D. At the D core site, the pore close off is at 72 m, and the mean methane age at this depth is 9 years with a standard deviation of 5.5 years for the age distribution [Schwander *et al.*, 1997]. The species-specific gas ages and the standard deviations of the age distributions at pore close off are calculated by scaling the mean methane age with the diffusivity of each gas relative to methane [Schwander *et al.*, 1997]. Subtracting these gas ages from the ice age at close off yields delta ages of 194, 191, 188,

193, 191, 192, and 189 years for C_2H_6 , C_3H_8 , $n\text{-C}_4\text{H}_{10}$, CH_3Cl , CH_3Br , OCS, and CS_2 , respectively. The standard deviations of age distributions are 8, 11, 12, 9, 11, 9, and 12 years for these gases in the same order. The gas ages at sampling depths for the length of the core are calculated by subtracting the delta ages from the ice ages for the GISP2 D core determined by layer counting [Alley *et al.*, 1993; Meese *et al.*, 1994; Meese *et al.*, special CRREL report, 94-1 US], resulting in a slightly different mean age for each gas species at the same depth. The ice ages for the GISP2 B are calculated by adding a 5-year offset to the ice ages of the D core at the same depth. This offset is based on a comparison of ice ages determined for both cores by continuous measurements of various major ions [Mayewski *et al.*, 1990, 1997; Taylor *et al.*, 1996; Yang *et al.*, 1995]. Calculated gas ages are reported as years before present (BP) referenced to the calendar year 2000.

[9] In the discussion of OCS results, GISP2 measurements are compared with previously published results from Siple Dome, Antarctica. The Siple Dome gas ages were obtained with a similar approach to what is used in this study. Details of the age calculations for that core are given in the works by Aydin *et al.* [2002] and Montzka *et al.* [2004]. Those gas ages are used unchanged in the current study.

2.2. Ice Core Extraction

[10] The air is liberated from the ice core samples using a dry-extraction technique, in which ice is mechanically shredded under vacuum [Etheridge *et al.*, 1988; Sowers and Jubenville, 2000]. The extraction chamber is a cylindrical electropolished stainless steel tube, which is 14.5 cm in diameter and 20 cm in length. The chamber ends are conflat flanges sealed by copper gaskets, and a bellows valve is welded to the flange on one end for transferring gas. A flat stainless steel grater is mounted lengthwise in the chamber. Since dry-extraction requires that the samples stay frozen during extraction, the chamber is mounted inside a chest freezer and connected to a motorized piston providing a linear motion of 13 cm at 1 Hz. When not in motion, the extraction chamber is connected to a stainless steel vacuum line via a flexible stainless steel bellows. All valves on the vacuum line are stainless steel bellows valves with alloy or copper valve tips.

[11] In preparation for the extraction, the ice core sample and the extraction chamber are precooled to -40 or -50°C . We found that the extraction process worked successfully at either temperature for the gases presented here. The outer 2–3 mm are shaved off, and the sample is placed on the grater inside the chamber. After loading the sample, the chamber is pumped down to 10^{-2} torr and repeatedly flushed with clean N_2 to remove residual air. The chamber is then disconnected from the vacuum line, and the motorized piston is operated for 15 min. The ice core shreds as it slides back and forth over the grater, liberating 70–80% of the air trapped in the sample. The GISP2 samples were 300–600 g, yielding 30–60 cm^3 standard temperature and pressure (STP) of air. After extraction, the chamber is reconnected to the vacuum line, and the air is condensed into a 76-cm long, 1/4" outside diameter (OD) stainless steel tube immersed in liquid helium. It takes about 4 min to

cryogenically pump >95% of the sample into the tube. The analysis of the extracted air starts 10 min after recovery.

2.3. Gas Analysis

[12] The air extracted from the ice cores is analyzed using gas chromatography (GC) with mass spectrometric detection (HP 5890/Micromass Autospec Ultima). The sample is cryogenically focused on a 1/8" OD glass bead trap immersed in liquid nitrogen. The size of the sample is determined by measuring the pressure change on a calibrated volume downstream of the cryotrap. An aliquot of isotope-labeled gas standard (30 cm³ STP) is added to the cryotrap to provide an internal reference for quantitation. The mixture of condensable gases from the sample and the internal standard are then transferred in a helium stream to a fused silica loop in liquid nitrogen and thermally injected onto the GC column. The GC oven is programmed to stay isothermal at -50°C for 3 min, ramp to -40°C at 5°C/min, followed by a 15°C/min ramp to 30°C and a 30°C/min ramp to 120°C.

[13] The mass spectrometer is operated in selected ion mode at a mass resolution of 6000 ($M/\Delta M$ at 5%), using ion fragments of deuterium (D-) labeled dodecane as the lock masses. At this mass resolution, halocarbons are fully resolved from hydrocarbons at the same mass to charge ratio [Engen *et al.*, 1998, 1999]. D-labeled dodecane is used because its ion fragments do not interfere with fragments from undeuterated atmospheric hydrocarbons, from ¹³C-labeled hydrocarbons used as internal standards, or from halocarbons.

[14] The primary standards used for the calibration of the system are prepared at University of California, Irvine (UCI), in Aculife-treated aluminum cylinders (Scott Specialty Gases) pressurized with dry N₂. Each standard cylinder contains a parts per billion (ppb)- level mixture of all the unlabeled trace gases, which are added into the cylinders by volumetric injection of commercially purchased pure compounds. One exception is CS₂. Since pure CS₂ is in liquid phase at laboratory temperature and pressure, it is prediluted to parts per thousand levels in a separate aluminum cylinder, which is used in the preparation of the ppb-level standards. Currently, there are three of these primary standards that were prepared in 2003, 2004, and 2005 calendar years. All three cylinders are used to calibrate a fourth standard cylinder, prepared at ppb-level in the same manner as the primary standards but containing the isotope-labeled analogues of the same trace gases. The isotope-labeled gas standard contains ¹³C-labeled C₂H₆, C₃H₈, *n*-C₄H₁₀, OCS, and CS₂ as well as fully deuterated CH₃Cl and CH₃Br. CFC-12 is also measured but not calibrated by isotope dilution. The stability of the trace gases in the pressurized cylinders is monitored by cross calibrating the standard tanks twice a year.

[15] System calibration and routine analyses with isotope dilution are carried out, using parts per trillion (ppt)- level working standards. Both the unlabeled and labeled working standards are prepared at near ambient levels by diluting the ppb-level primary standards with humid N₂ in electropolished stainless steel flasks. New working standards are prepared every 3–4 weeks to avoid compromising their stability in the flasks. The use of isotope-labeled internal standards compensates for drift in the mass spectrometer

response and run-to-run variations in chromatographic peak shape.

3. Data

[16] Thirty ice core samples were analyzed as part of this project with 15 samples each from the GISP2 B and D cores. All data are corrected for gravitational effects that act to enrich the air in the bottom of the firn with respect to gases that have higher molecular weights than air. In the order of size, the gravitational corrections are 0.04%, 0.5%, 0.7%, 1.0%, 1.1%, 1.7%, and 2.3% for C₂H₆, C₃H₈, CH₃Cl, *n*-C₄H₁₀, OCS, CS₂, and CH₃Br, respectively. In addition to the gases presented here, CFC-12 was also measured as an indicator of contamination by modern air due either to air trapped in fractures in the ice sample or to the leaks during the handling of the gas sample. Three GISP2 B and two GISP2 D samples had exceptionally high mixing ratios of CFC-12 (>10 ppt) and were not included in this data set, lowering the total number of samples to 25. The average CFC-12 level is 2 ± 3 ppt (1σ) for the 12 GISP2 B samples and less than 1 ± 2 ppt (1σ) for the 13 GISP2 D samples. Nitrogen blank runs were always CFC-12 free, and barometric leak checks did not indicate any problems with the integrity of the vacuum systems during extraction and analyses. We suggest that the low levels of CFC-12 measured in most ice core samples may reflect diffusion of air along small fractures in the ice core, with the presumption that the preindustrial atmosphere was CFC free.

[17] Since CFC-12 currently measures over 500 ppt in the atmosphere [Walker *et al.*, 2000] and potentially higher in storage freezers where ice cores are stored, even a minute amount of modern air trapped in a fracture can cause a nonzero CFC-12 measurement. For example, 100 μl STP of modern air at 500 ppt mixed with 50 cm³ STP of CFC-free air from an ice core will result in a 1 ppt measurement. If the ice core and modern air mixing ratios were 100 ppt and 1000 ppt, respectively, for a given gas, the same amount of intrusion would result in a 101.8 ppt measurement. Thus gases with large anthropogenic components should be the most sensitive to contamination by modern air. In this study, none of the gases, including the alkanes, display any correlation with the nonzero CFC-12 measurements. We conclude that the existence of CFC-12 in some ice core samples has no significant impact on the data presented here.

[18] Analytical uncertainties for all mixing ratios in the subsequent sections and tables are reported as ±1σ. One factor that contributes to the analytical uncertainty is the variability in the background levels of each trace gas in the ice core extraction system as determined by N₂ blank runs, acquired before and after an ice core sample is shredded. For this purpose, about 30 cm³ STP of clean N₂ is isolated in the extraction chamber for the same amount of time it takes to shred an ice core sample. The size of the blank correction for each sample is determined from the average of N₂ blanks taken before and after shredding the sample. The preshredded N₂ blanks are isolated in the chamber over a whole piece of ice core sample, and the postshredded blanks are exposed to the shredded sample. The extraction and analyses of the N₂ blanks are carried out following exactly the same procedures as a sample to capture the combined background level

Table 1. Molar Ratio of Analytical Blank to Ice Core Signal for Various Trace Gases in Percentages

	GISP2 B	GISP2 D	Overall
C ₂ H ₆	3.3 ± 0.9	3.5 ± 0.6	3.4 ± 0.8
C ₃ H ₈	8.3 ± 1.4	7.8 ± 1.9	8.0 ± 1.6
<i>n</i> -C ₄ H ₁₀	8.5 ± 1.0	7.8 ± 2.4	8.2 ± 1.9
CH ₃ Cl	2.0 ± 1.1	2.6 ± 2.8	2.3 ± 2.1
CH ₃ Br	1.4 ± 1.1	0.8 ± 0.6	1.0 ± 0.9
OCS	2.7 ± 0.4	6.7 ± 4.5	4.8 ± 3.8
CS ₂	13.3 ± 7.2	13.7 ± 6.7	13.5 ± 6.8

contamination in both the extraction and analytical systems as they apply to an ice core sample. Ice core samples, preshred, and postshred blanks are all wet, containing roughly the same amount of humidity because of sublimation from the ice surface. The samples are analyzed without any chemical or physical dehumidification procedure.

[19] The size of the blank correction is only 1 and 2% of the signal for the two methyl halides, over 3% for C₂H₆, about 8% for C₃H₈ and *n*-C₄H₁₀, 5% for OCS, and close to 14% for CS₂ (Table 1). The blank correction for CS₂ is large and quite variable. For the remainder of the compounds, the blank corrections have a minor impact on the overall analytical uncertainty. For example, the C₃H₈ and *n*-C₄H₁₀ blanks are relatively high compared to signal size, but they are reproducible. CH₃Cl, CH₃Br, and OCS blanks are highly variable but equal only a small fraction of the signal. There is no difference in the blanks between the dry-drilled B core and the fluid-drilled D core except for OCS. OCS blanks for the D core samples are higher on average and display more variability. On the basis of the blank variability observed for the GISP2 data set, the detection limits (3σ) of our system for an ice core sample of average size (~ 35 cm³ STP of air) are 92, 69, 43, 30, 2.1, 31, and 3.3 ppt for C₂H₆, C₃H₈, C₄H₁₀, CH₃Cl, CH₃Br, OCS, and CS₂, respectively.

[20] Analytical precision is the second factor that contributes significantly to the overall uncertainties reported for the current data set. The precision is determined from the reproducibility of the calibration runs for the analytical system as well as the variation in factors inherent to the gas extraction process such as ice core sample size, extraction efficiency, and the concentration of each particular gas in the extracted air.

[21] The accuracy of the analysis includes uncertainties arising from standard preparation in addition to the factors that affect the precision. The accuracy of the standards that are used in calibration of the system is estimated at 2–3% on the basis of the reproducibility of primary standard tanks prepared at the UCI. Informal intercalibration efforts between our laboratory at the UCI and NOAA/ESRL-GMD for OCS, CH₃Cl, and CH₃Br show agreement to within 2–3% [Aydin et al., 2004; Montzka et al., 2004] (S. A. Montzka, personal communication) or within the range of our estimated accuracy.

4. Results and Discussion

4.1. Alkanes

4.1.1. Ethane

[22] Modern atmospheric C₂H₆ levels in the Northern Hemisphere range from 0.3 to 3.5 ppb, with the highest mixing ratios occurring at high latitudes [Kanakidou et al.,

1991; Rudolph, 1995; Boissard et al., 1996]. Rudolph [1995] estimates the modern annual average C₂H₆ mixing ratio in the Arctic lower troposphere at 2.0–2.5 ppb, with a steep latitudinal gradient between midlatitudes and the Arctic. Land-based measurements at Fraserdale (50°N, 82°W) and Alert (82°N, 62°W), Canada, Rorvik (57°N, 12°E), Sweden, and Pallas (68°N, 24°E), Norway, show wintertime average levels in the 2.2–2.7 ppb range and mean summertime levels in the 0.8–1.0 ppb range [Gautrois et al., 2003; Jobson et al., 1994; Laurila and Hakola, 1996; Lindsog and Moldanova, 1994]. Airborne measurements from the lower troposphere in the high northern latitudes show wintertime C₂H₆ levels of 2.0–2.5 ppb [Anderson et al., 1993; Blake et al., 2003; Penkett et al., 1993] and summertime levels at Summit, Greenland are 0.7 ppb [Swanson et al., 2002]. C₂H₆ is removed from the atmosphere by reaction with hydroxyl radical (OH), with a mean annual lifetime of about a few months [Blake et al., 2003; Boissard et al., 1996]. The atmospheric burden of C₂H₆ is consistent with the estimated emissions from biomass burning and natural gas losses, with minor contributions from wetlands, soils, oceans, and automotive as well as industrial emissions [Rudolph, 1995].

[23] C₂H₆ levels in the GISP2 ice core samples range from 0.96 to 2.53 ppb with a mean of 1.62 ± 0.43 ppb (Figure 1 and Table 2). In five pairs of contemporaneous samples from the B and D cores at 180, 192, 204, 222, and 283 BP (given ages are for D core samples), the agreement is very good. Only the pair at 275 BP displays a large difference, with one fluid-drilled sample having 2.04 ppb of C₂H₆ compared with 1.15 and 1.25 ppb of C₂H₆ in the two adjacent dry-drilled samples. The highest C₃H₈ and *n*-C₄H₁₀ mixing ratios in the entire data set are also observed in the same fluid-drilled sample from 275 BP (Figure 1). C₃H₈ and *n*-C₄H₁₀ measurements appear to be more sensitive to the effects of drill-fluid contamination as discussed in the following sections. It is likely that this sample represents an extreme case of drill-fluid contamination.

4.1.2. Propane

[24] Propane levels in the modern atmosphere are generally about 30% of atmospheric ethane levels, although data are more limited for propane, especially at high latitudes of the Northern Hemisphere [Poisson et al., 2000]. C₃H₈ levels at Fraserdale, Canada (50°N, 82°W) averaged 0.08–1.22 ppb from summer to winter over the period 1990–1992 [Jobson et al., 1994], while at Alert (82°N, 62°W), Canada over the period 1989–1996, the mean C₃H₈ levels were about 0.71 ppb with a summer to winter range of 0.16–1.42 ppb [Gautrois et al., 2003]. At Rorvik (57°N, 12°E), Sweden, wintertime C₃H₈ levels in 1989–1990 averaged about 1.27 ppb and dropped to 0.24 ppb in summer [Lindsog and Moldanova, 1994], and at Pallas (68°N, 24°E), Norway in 1993–1994, the mean C₃H₈ level was 0.6 ppb, with a variability of 0.14–1.20 ppb between summer and winter [Laurila and Hakola, 1996]. Airborne measurements from the high northern latitudes show wintertime C₃H₈ levels near 1 ppb, suggesting mean annual levels of roughly 0.5 ppb [Anderson et al., 1993; Blake et al., 2003; Penkett et al., 1993]. C₃H₈ has an atmospheric lifetime of a few weeks [Blake et al., 2003].

[25] The range of C₃H₈ levels in the dry- and fluid-drilled ice core samples are 0.32–0.49 and 0.28–1.0 ppb,




Figure 1. C_2H_6 , C_3H_8 , and $n-C_4H_{10}$ mixing ratios in Greenland ice core samples from the dry-drilled GISP2 B core (●) and the fluid-drilled GISP2 D core (○). Error bars are shown only when they are larger than the symbols.

respectively (Figure 1). There are several samples for which there is excellent agreement between the two ice cores. However, there is an overall tendency for C_3H_8 in the fluid-drilled samples to be higher than contemporary samples of the dry-drilled core. The presence of the drilling fluid, which probably penetrated the ice along fractures that later healed, appear to contaminate C_3H_8 as well as the $n-C_4H_{10}$ results. For the B core, the C_3H_8 levels are roughly 20–30% of the corresponding C_2H_6 measurements, similar to their modern relative abundance in the Northern Hemisphere, and exhibit a standard deviation of only 15% unlike the highly variable C_2H_6 (Table 2). Overall, ice core C_3H_8 levels and C_3H_8/C_2H_6 ratios are similar to those in modern high-latitude Northern Hemispheric air.

4.1.3. *n*-Butane

[26] $n-C_4H_{10}$ levels in the Arctic lower troposphere range from about 0.3–0.5 ppb during winter to less than 0.2 ppb during summertime based on airborne studies [Anderson *et al.*, 1993; Blake *et al.*, 2003; Penkett *et al.*, 1993]. Land-based measurements at high latitudes vary from 0.4–0.8 ppb in winter to 40–50 ppt in summer [Gautrois *et al.*, 2003; Jobson *et al.*, 1994; Laurila and Hakola, 1996; Lindskog and Moldanova, 1994]. $n-C_4H_{10}$ is short-lived, with a mean annual lifetime of about 10 days to a few weeks [Blake *et*

al., 2003]. $n-C_4H_{10}$ levels in the dry- and wet-drilled ice core samples range from 0.2 to 0.3 ppb and 0.2 to 0.6 ppb, respectively (Figure 1). The pattern of variability of $n-C_4H_{10}$ in the ice is very similar to that of C_3H_8 , with a similar tendency for the fluid-drilled core to have elevated levels (Table 2).

[27] C_2H_6 exhibits a surprisingly high level of variability in the ice that is not shared by C_3H_8 and $n-C_4H_{10}$. This variability in the C_2H_6 signal is apparently a real feature of the ice chemistry, as evidenced by the covariance between the two ice cores. There is no evidence of severe analytical artifacts as the signal for the alkanes in the samples are well above the analytical blanks, and there are no obvious problems associated with extraction of the gas from the ice. There is evidence for drill-fluid contamination of C_3H_8 and $n-C_4H_{10}$ in some samples, but this is not severe.

[28] The observed levels of C_2H_6 , C_3H_8 , and $n-C_4H_{10}$ are well within the range observed in the modern atmosphere. The ratios of the various alkanes are also generally consistent with an atmospheric origin for the ice core measurements. The atmospheric ratios of these gases presumably reflect the combined effect of the relative strength of their sources and their OH rate constants. If the ice core signal were a result of some photochemical or biological process in firm or ice, one would expect to observe anomalous alkane ratios in the ice bubbles. In an analysis of firm air from Greenland, Worton [2006] reported that the amount of C_2H_6 , C_3H_8 , and $n-C_4H_{10}$ over Greenland peaked in the 1990s at levels up to two times higher than 1950 values. Worton [2006] also noted a considerable but relatively smaller drop in light alkane levels since the 1990s, which they attribute to changes in hydrocarbon emissions. However, current budget estimates attribute most of the C_2H_6 in the Arctic region to natural gas losses [Olivier *et al.*, 1996; Rudolph, 1995], and global natural gas consumption has nearly doubled between 1980 and 2004 (<http://www.eia.doe.gov/emeu/international/>). Assuming that the nonanthropogenic source terms in the C_2H_6 budget have not changed, it is expected that preindustrial atmospheric C_2H_6 levels should have been considerably lower than modern levels. In this regard, the GISP2 ice core results are not consistent with current budget estimates.

4.2. Methyl Chloride and Methyl Bromide

[29] CH_3Cl is the most abundant halocarbon in the troposphere, with a global average mixing ratio of 550 ± 30 ppt and an atmospheric lifetime estimated at 1.3 years [Keene *et al.*, 1996; Khalil and Rasmussen, 1999, 2000; Montzka *et al.*, 2003a; Moore *et al.*, 1996]. Firm air measurements from Antarctica and Greenland suggest that the atmospheric levels of CH_3Cl remained within 10% of

Table 2. Average Mixing Ratio of All Compounds Reported as ppt With $\pm 1\sigma$ of Measured Quantities

	GISP2 B	GISP2 D	Overall
C_2H_6	1606 ± 491	1627 ± 393	1617 ± 434
C_3H_8	390 ± 60	486 ± 191	440 ± 149
$n-C_4H_{10}$	226 ± 33	278 ± 115	253 ± 89
CH_3Cl	692 ± 299	717 ± 384	705 ± 339
CH_3Br	59 ± 34	105 ± 198	83 ± 144
OCS	334 ± 19	317 ± 23	325 ± 23
CS_2	10 ± 5	10 ± 8	10 ± 6




Figure 2. CH_3Cl and CH_3Br mixing ratios in Greenland ice core samples from the dry-drilled GISP2 B core (●) and the fluid-drilled GISP2 D core (○). The CH_3Br mixing ratio for the D core sample at 220 BP is off-scale at 760 ppt. The scale of the y axis is set at 125 ppt to better represent the data set as a whole. Error bars are shown only when they are larger than the symbols.

the current levels over the last few decades [Aydin *et al.*, 2004; Butler *et al.*, 1999; Kaspers *et al.*, 2004; Trudinger *et al.*, 2004]. Ice core measurements from Antarctica indicate that the average CH_3Cl mixing ratio during the last 300 years was 499 ± 28 ppt [Aydin *et al.*, 2004]. These measurements support the modern budget which asserts that the anthropogenic emission sources for CH_3Cl are much smaller than the natural sources [Montzka *et al.*, 2003a].

[30] The average CH_3Cl mixing ratio measured in GISP2 ice cores is 705 ± 339 ppt (Table 2), exceeding both the mean level and variability observed in Antarctic ice cores (Figure 2). Despite the high mean value, the CH_3Cl levels are around 500 ppt during the youngest 60 years of the GISP2 record, in accord with what would be expected from firn records. However, there are several measurements greater than 600 ppt and five scattered at or above 1 ppb between 185 and 300 BP, suggesting that the data may be impacted by in situ CH_3Cl production. If the elevated points are ignored, there appears to be a “baseline” CH_3Cl level of about 500 ppt.

[31] Episodic in situ production may be linked to particulate deposition [Sturges *et al.*, 2001a] or other environmental factors, such as extreme warming and melting events, that are linked to the ice itself rather than the air trapped in the bubbles. The ice ages of the samples with elevated CH_3Cl measurements correspond to a 150-year period between 1500 and 1650 AD (approximate gas ages of 1700–1850 BP; Figure 2) that is marked by a warming climate at Summit, as evidenced by a 10% increase in the snow accumulation rate [Meese *et al.*, 1994]. However,

there are no melt layers [Meese *et al.*, 1994] or obvious anomalies in ice chemistry [Mayewski *et al.*, 1990, 1997; Taylor *et al.*, 1996; Yang *et al.*, 1995] that correlate with the elevated CH_3Cl during this period. There is also no indication that the drill fluid has any effect on CH_3Cl measurements (Figure 2). The B and D core averages are similar at 692 ± 299 and 717 ± 384 ppt, respectively (Table 2), yet only three of the six sample pairs show good agreement between B and D cores. If in situ production is responsible for this scatter, the processes responsible do not appear to be continuous and ongoing since the highest measurements are concentrated in the middle of the record around 220 BP. It seems more likely that the production occurred in the firn or ended some time after bubble close off and that a warmer local climate might have triggered it. The results place doubts on the prospects of recovering a paleoatmospheric CH_3Cl record from Greenland ice cores.

[32] CH_3Br is a precursor of stratospheric ozone depleting bromine and has a tropospheric lifetime of 0.7 years [Colman *et al.*, 1998; Kurylo *et al.*, 1999; Shorter *et al.*, 1995; Warwick *et al.*, 2006; Yvon-Lewis and Butler, 1997]. Tropospheric CH_3Br peaked at about 10.5 ppt in the Northern Hemisphere during the late 1990's, with an inter-hemispheric gradient slightly higher than 2 ppt [Montzka *et al.*, 2003b]. CH_3Br levels have been dropping since then primarily because of the restrictions under the Montreal Protocol [Montzka *et al.*, 2003b; Yokouchi *et al.*, 2002]. Model simulations based on Antarctic firn air and ice core measurements suggest that preindustrial levels of CH_3Br were lower in both hemispheres with a smaller interhemispheric gradient [Reeves, 2003; Saltzman *et al.*, 2004].

[33] In the GISP ice cores, CH_3Br levels are well above ambient levels and highly scattered, with an average mixing ratio of 83 ± 144 ppt (Table 2 and Figure 2). The data strongly suggest that CH_3Br undergoes in situ enrichment similar to what has been found in Northern Hemisphere firn air [Butler *et al.*, 1999; Sturges *et al.*, 2001a]. The enrichment of CH_3Br can exceed the ambient levels by more than an order of magnitude and it is not restricted to a particular section of the ice cores. There does appear to be some relationship between the enrichments of CH_3Br and CH_3Cl . The minimum and maximum D core CH_3Br measurements are 12 and 760 ppt at 195 and 220 BP, respectively, both measured in samples with extreme CH_3Cl values. In the B core, the minimum CH_3Br and CH_3Cl values of 12 and 476 ppt also occur in the same sample from 226 BP. The maximum CH_3Br measurement from the B core is 115 ppt at 276 BP, but the CH_3Cl value for this sample is only slightly elevated at 660 ppt. For all 25 data points, omitting one very high measurement for both compounds at 220 BP, the correlation coefficient between CH_3Br and CH_3Cl is 0.37 with a p value of 0.072, not showing significance at 95% level. There is likely more than one mechanism that leads to the production of methyl halides, as the enrichment of the brominated species appears to be more widespread.

4.3. Carbonyl Sulfide and Carbon Disulfide

[34] OCS is the most abundant sulfur species in the troposphere with a mixing ratio of about 500 ppt and a




Figure 3. OCS and CS₂ mixing ratios in Greenland ice core samples from the dry-drilled GISP2 B core (●) and the fluid-drilled GISP2 D core (○). Error bars are shown only when they are larger than the symbols.

lifetime of 2–4 years [Chin and Davis, 1995; Montzka et al., in print, JGR 2006]. Results of the first OCS measurements in firn air from Canada and Antarctica suggest that this compound is stable in the firn with middle to late twentieth century levels scattered around 500 ppt in both hemispheres [Sturges et al., 2001b]. Subsequent measurements in Antarctic firn air and ice cores extended the length of the record to 350 years, with OCS levels varying between 300 and 400 ppt [Aydin et al., 2002; Montzka et al., 2004].

[35] The results from GISP B and D cores display steady OCS levels with a mean of 325 ± 23 ppt for the Northern Hemisphere between 125 and 325 BP (Table 2 and Figure 3). The OCS levels between 215 and 270 BP are marginally lower than the rest at 300–310 ppt. The data from both cores agree within errors where they overlap, suggesting that the OCS signal in the GISP cores is robust.

[36] In Figure 4, the OCS data from the GISP core are compared with the 350 year record from a Siple Dome Antarctic ice core [Montzka et al., 2004]. The overall agreement between the Greenland and Antarctic OCS records strongly suggest that ice cores from both hemispheres can be used to develop paleo records for this trace gas over centennial timescales. Between 300 and 320 BP, Antarctic OCS levels are about 75–100 ppt greater than those measured at Greenland (Figure 4). If confirmed by other ice cores, such an interhemispheric gradient would likely indicate either an increase in the Southern Hemisphere oceanic source or a decrease in the Northern Hemisphere terrestrial sink during this period.

[37] CS₂ is a precursor to atmospheric OCS, with both natural and anthropogenic sources and an atmospheric

lifetime of several days. Ambient levels of CS₂ in marine air are typically in the range of 0.5–10 ppt, although there are relatively few measurements to date [Bandy et al., 1993; Johnson and Bates, 1993; Kim and Andreae, 1992; Kim and Andreae, 1987; Tucker et al., 1985]. Over the central and eastern North Pacific where the effects of the anthropogenic emissions diminish, CS₂ levels are close to 4 ppt in the boundary layer and drop to about 2 ppt at 4 km altitude [Blake et al., 2004]. CS₂ levels in the GISP ice core samples are similar to those in ambient air with an overall average of 10 ± 6 ppt (Table 2 and Figure 3). There is reasonably good agreement between the two cores, with the notable exception of an outlier from the D core with a mixing ratio of 34 ppt at 209 BP. The OCS measurement for this sample also looks slightly anomalous. In our laboratory, CS₂ contamination is usually accompanied by elevated OCS levels. We attribute this to contamination from sulfur cross-linked polymers. In the ice core samples, most of the samples with higher than average CS₂ levels are apparently not associated with elevated OCS. It is possible that there is a stronger emission of CS₂ from the contamination source, or the OCS contamination is masked by its relatively high abundance. Contamination at low ppt levels can essentially remain less than the measurement uncertainty against an OCS background that is at 300 ppt and above.

[38] Current budget estimates for CS₂ suggest that the oceans should be the dominant source of CS₂ to the preindustrial atmosphere [Andreae and Ferek, 1992; Chin and Davis, 1993; Kettle et al., 2002; Kjellström, 1998; Watts, 2000]. Although their atmospheric lifetimes are very different, one might expect some correlation between the atmospheric levels of OCS and CS₂. It is interesting that the lowest OCS mixing ratios in the GISP record are roughly from the same time period between 215 and 275 BP as the lowest CS₂ measurements (Figure 3). Currently, there are no firn or surface air measurements of CS₂ from Summit, Greenland with which the ice core measurements can be compared. However, the data them-




Figure 4. OCS mixing ratios in contemporaneous Greenland and Antarctic ice core samples. Ice core samples from the dry-drilled GISP2 B core (●), the fluid-drilled GISP2 D core (○), and the dry-drilled Siple Dome C core (□); [Aydin et al., 2002; Montzka et al., 2004].

selves are consistent with an atmospheric origin of the ice core CS₂ signal.

5. Conclusions

[39] This study demonstrates the successful extraction and analysis of several short-lived trace gas species in Greenland ice cores. For all the gases measured, the signal derived from air extracted from ice core bubbles is above our detection limits. Only a few of the C₃H₈ and *n*-C₄H₁₀ samples appear to be elevated because of drill-fluid contamination. In future studies, the presence of drill fluid in the ice core samples can probably be minimized by careful sample selection. Monitoring additional hydrocarbons that are more sensitive to drill-fluid contamination could facilitate detection and possibly quantification of the drill-fluid contamination.

[40] The interpretation of the alkane measurements is challenging. Overall, the alkane mixing ratios in the ice are consistent with those in the modern Arctic atmosphere, and do not provide a priori evidence of in situ production. However, a paleoatmospheric interpretation of the data conflicts with current atmospheric budgets of the alkanes. Emissions because of natural gas and oil use are thought to constitute a major part of the total burden [Boissard *et al.*, 1996; Katzenstein *et al.*, 2003; Rudolph, 1995], suggesting that there should have been lower atmospheric levels during preindustrial times. We consider the origin of the current levels of alkanes in Greenland ice to be an open issue, which will require additional measurements to resolve.

[41] The methyl halide measurements indicate that in situ production processes, rather than paleoatmospheric levels, dominate the CH₃Br signal in GISP ice cores and may influence to a lesser extent the ice core record of CH₃Cl. The GISP2 OCS measurements are in good agreement with those from Antarctic ice cores, and it appears likely that the ice core signals reflect paleoatmospheric mixing ratios. This raises the possibility of reconstructing a record of the inter-hemispheric gradient for OCS. The OCS budgets in the two hemispheres are quite different, with terrestrial processes dominating in the north and oceanic processes controlling in the south [Chin and Davis, 1993; Kettle *et al.*, 2002; Kjellström, 1998; Watts, 2000]. The land sink for atmospheric OCS is related to gross primary productivity [Montzka *et al.*, in press, JGR 2007] through uptake by plants [Goldan *et al.*, 1988], and the Greenland ice core record may provide insight into the variability of the Northern Hemisphere terrestrial biosphere on short timescales. The CS₂ measurements in the GISP cores appear reasonable in light of current atmospheric levels, but our knowledge of the modern atmospheric distribution of this compound is insufficient to draw a firm conclusion as to the origin of this signal.

[42] **Acknowledgments.** We would like to thank all PIs, the scientific staff, and the drilling team that contributed to the realization of the GISP project. We also would like to thank National Ice Core Laboratories for their assistance in sampling the GISP cores and Stephen A. Montzka for insightful discussions. This work was supported by NSF-Office of Polar Programs grant OPP0221480 and by the Gary Comer Science and Education Foundation.

References

- Alley, R. B., *et al.* (1993), Abrupt increase in Greenland snow accumulation at the end of the Younger Dryas event, *Nature*, 362, 527–529.
- Anderson, B. E., J. E. Collins, Jr., G. W. Sachse, G. W. Whiting, D. R. Blake, and F. S. Rowland (1993), AASE-II observations of trace carbon species distributions in the mid to upper troposphere, *Geophys. Res. Lett.*, 20, 2539–2542.
- Andreae, M. Ö., and R. R. Ferek (1992), Photochemical production of carbonyl sulfide in seawater and its emissions to the atmosphere, *Global Biogeochem. Cycles*, 6, 175–183.
- Aydin, M., W. J. DeBruyn, and E. S. Saltzman (2002), Preindustrial atmospheric carbonyl sulfide (OCS) from an Antarctic ice core, *Geophys. Res. Lett.*, 2(9), 1359, doi:10.1029/2002GL014796.
- Aydin, M., W. J. DeBruyn, E. S. Saltzman, S. A. Montzka, and J. H. Butler (2004), Atmospheric variability of methyl chloride during the last 300 years from an Antarctic ice core and firn air, *Geophys. Res. Lett.*, 31, L02109, doi:10.1029/2003GL018750.
- Bandy, A. R., D. C. Thornton, and J. E. Johnson (1993), Carbon disulfide measurements in the atmosphere of the western North Atlantic and the northwestern South Atlantic Oceans, *J. Geophys. Res.*, 98(D12), 23,449–23,457.
- Blake, N. J., D. R. Blake, B. C. Sive, A. S. Katzenstein, S. Meinardi, O. W. Wingenter, E. L. Atlas, F. Flocke, B. A. Ridley, and F. S. Rowland (2003), The seasonal evolution of NMHCs and light alkyl nitrates at middle to high northern latitudes during TOPSE, *J. Geophys. Res.*, 108(D4), 8359, doi:10.1029/2001JD001467.
- Blake, N. J., *et al.* (2004), Carbonyl sulfide and carbon disulfide: Large-scale distributions over the western Pacific and emissions from Asia during TRACE-P, *J. Geophys. Res.*, 109(D15S05), doi:10.1029/2003JD004259.
- Boissard, C., B. Bonsang, M. Kanakidou, and G. Lambert (1996), TROPOZ II: Global distributions and budgets of methane and light hydrocarbons, *J. Atmos. Chem.*, 25(2), 115–148.
- Butler, J. H., M. Battle, M. L. Bender, S. A. Montzka, A. D. Clarke, E. S. Saltzman, C. M. Sucher, J. P. Severinghaus, and J. W. Elkins (1999), A record of atmospheric halocarbons during the twentieth century from polar firn air, *Nature*, 399, 749–755.
- Chin, M., and D. D. Davis (1993), Global sources and sinks of OCS and CS₂ and their distributions, *Global Biogeochem. Cycles*, 7, 321–337.
- Chin, M., and D. D. Davis (1995), A reanalysis of carbonyl sulfide as a source of stratospheric background sulfur aerosol, *J. Geophys. Res.*, 100(D5), 8993–9006.
- Colman, J. J., D. R. Blake, and F. S. Rowland (1998), Atmospheric residence time of CH₃Br estimated from the Junge spatial variability relation, *Science*, 281, 392–396.
- Engen, M. A., V. A. Wagner, L. J. Sears, and E. P. Grimsrud (1998), Detection of CF₃-containing compounds in background air by gas chromatography/high-resolution mass spectrometry, *J. Geophys. Res.*, 103, 25,287–25,297.
- Engen, M. A., J. A. Culbertson, and E. P. Grimsrud (1999), Analysis of brominated compounds in background air by gas chromatography-high resolution mass spectrometry, *J. Chromatogr. A*, 848, 261–277.
- Etheridge, D. M., G. I. Pearman, and F. de Silva (1988), Atmospheric trace-gas variations as revealed by air trapped in an ice core from Law Dome, Antarctica, *Ann. Glaciol.*, 10, 28–33.
- Gautrois, M., T. Brauers, R. Koppmann, F. Rohrer, O. Stein, and J. Rudolph (2003), Seasonal variability and trends of volatile organic compounds in the lower polar troposphere, *J. Geophys. Res.*, 108(D13), 4393, doi:10.1029/2002JD002765.
- Goldan, P. D., R. Fall, W. C. Kuster, and F. C. Fehsenfeld (1988), Uptake of COS by growing vegetation: A major tropospheric sink, *J. Geophys. Res.*, 93, 14,186–14,192.
- Jobson, B. T., H. Niki, Y. Yokouchi, J. Bottenheim, F. Hopper, and R. Leaitch (1994), Measurements of C₂–C₆ hydrocarbons during the Polar Sunrise 1992 Experiment: Evidence for Cl atom and Br atom chemistry, *J. Geophys. Res.*, 99(D12), 25,355–25,368.
- Johnson, J. E., and T. S. Bates (1993), Atmospheric measurements of carbonyl sulfide, dimethyl sulfide, and carbon disulfide using the electron capture sulfur detector, *J. Geophys. Res.*, 98(D12), 23,411–23,421.
- Kanakidou, M., K. M. Valentin, P. J. Crutzen, and H. B. Singh (1991), A two-dimensional study of ethane and propane oxidation in the troposphere, *J. Geophys. Res.*, 96, 15,395–15,413.
- Kaspers, K. A., *et al.* (2004), Analyses of firn gas samples from Dronning Maud Land, Antarctica: Study of nonmethane hydrocarbons and methyl chloride, *J. Geophys. Res.*, 109(D2), D02307, doi:10.1029/2003JD003950.
- Katzenstein, A. S., L. A. Doezeza, I. J. Simpson, D. R. Blake, and F. S. Rowland (2003), Extensive regional atmospheric hydrocarbon pollution in the southwestern United States, *Proc. Natl. Acad. Sci. U. S. A.*, 100(21), 11,975–11,979.
- Keene, W. C., *et al.* (1996), Composite global emissions of reactive chlorine from anthropogenic and natural sources: Reactive chlorine emissions inventory, *J. Geophys. Res.*, 104, 8429–8440.

- Kettle, A. J., U. Kuhn, M. von Hobe, J. Kesselmeier, and M. Ö. Andreae (2002), Global budget of atmospheric carbonyl sulfide: Temporal and spatial variations of the dominant sources and sinks, *J. Geophys. Res.*, *107*(D22), 4658, doi:10.1029/2002JD002187.
- Khalil, M. A. K., and R. A. Rasmussen (1999), Atmospheric methyl chloride, *Atmos. Environ.*, *33*, 1305–1321.
- Khalil, M. A. K., and R. A. Rasmussen (2000), Soil-atmosphere exchange of radiatively and chemically active gases, *Environ. Sci. Pollut. Res. Int.*, *7*, 79–82.
- Kim, K.-H., and M. Ö. Andreae (1987), Carbon disulfide in seawater and the marine atmosphere over the North Atlantic, *J. Geophys. Res.*, *92*, 14,733–14,738.
- Kim, K.-H., and M. Ö. Andreae (1992), Carbon disulfide in the estuarine, coastal, and oceanic environments, *Mar. Chem.*, *40*, 179–197.
- Kjellström, E. (1998), A three-dimensional global model study of Carbonyl Sulfide in the troposphere and the lower stratosphere, *J. Atmos. Chem.*, *29*, 151–177.
- Kurylo, M. J., et al. (1999), Short-lived ozone-related compounds, in *Scientific Assessment of Ozone Depletion: 1998*, chapter 2, Global Ozone Research and Monitoring Project—Report 44, World Meteorological Organization, Geneva.
- Laurila, T., and H. Hakola (1996), Seasonal cycle of C₂-C₅ hydrocarbons over the Baltic Sea and northern Finland, *Atmos. Environ.*, *30*(10/11), 1597–1607.
- Lindskog, A., and J. Moldanova (1994), The influence of the origin, season, and time of the day on the distribution of individual NMHC measured at Rorvik, Sweden, *Atmos. Environ.*, *28*(15), 2383–2398.
- Mayewski, P. A., W. B. Lyons, M. J. Spencer, M. S. Twickler, C. F. Buck, and S. I. Whitlow (1990), An ice core record of atmospheric response to anthropogenic sulphate and nitrate, *Nature*, *346*, 554–556.
- Mayewski, P. A., L. D. Meeker, M. S. Twickler, S. I. Whitlow, Q. Yang, W. B. Lyons, and M. Prentice (1997), Major features and forcing of high-latitude Northern Hemisphere atmospheric circulation using a 110,000-year-long glaciochemical series, *J. Geophys. Res.*, *102*, 26,345–26,366.
- Meese, D. A., A. J. Gow, P. Grootes, P. A. Mayewski, M. Ram, M. Stuiver, K. C. Taylor, E. D. Waddington, and G. A. Zielinski (1994), The accumulation record from the GISP2 core as an indicator of climate change through the Holocene, *Science*, *266*(5191), 1680–1682.
- Montzka, S. A., et al. (2003a), Controlled substances and other source gases, in *Scientific Assessment of Ozone Depletion: 2002*, chapter 1, Global Ozone Research and Monitoring Project—Report 47, World Meteorological Organization, Geneva.
- Montzka, S. A., J. H. Butler, B. D. Hall, D. J. Mondeel, and J. W. Elkins (2003b), A decline in tropospheric organic bromine, *Geophys. Res. Lett.*, *30*(15), 1826, doi:10.1029/2003GL017745.
- Montzka, S. A., M. Aydin, J. H. Butler, M. Battle, E. S. Saltzman, G. S. Dutton, B. D. Hall, A. D. Clarke, D. Mondeel, and J. W. Elkins (2004), A 350-year atmospheric history for carbonyl sulfide inferred from Antarctic firm air and air trapped in ice, *J. Geophys. Res.*, *109*(D22302), doi:10.1029/2004JD004686.
- Moore, R. M., W. Groszko, and S. J. Niven (1996), Ocean-atmosphere exchange of methyl chloride: Results from NW Atlantic and Pacific Ocean studies, *J. Geophys. Res.*, *101*, 28,529–28,538.
- Olivier, J. G. J., A. F. Bouwman, C. W. M. Van der Maas, J. J. M. Berdowski, C. Veldt, J. P. J. Bloos, A. J. H. Visschedijk, P. Y. J. Zandveld, and J. L. Haverlag (1996), Description of EDGAR Version 2.0: A set of global emission inventories of greenhouse gases and ozone-depleting substances for all anthropogenic and most natural sources on a per country basis and on 1° × 1° grid, National Institute of Public Health and the Environment (RIVM), Bilthoven, Report 771060 002/TNO-MEP report R96/110.
- Penkett, S. A., N. J. Blake, P. Lightman, A. R. W. Marsh, P. Anwyl, and G. Butcher (1993), The seasonal variation of non-methane hydrocarbons in the free troposphere over the North Atlantic Ocean: Possible evidence for extensive reaction of hydrocarbons with the nitrate radical, *J. Geophys. Res.*, *98*, 2865–2885.
- Poisson, N., M. Kanakidou, and P. J. Crutzen (2000), Impact of NMHCs on tropospheric chemistry and the oxidizing power of the global troposphere: 3-dimensional modeling results, *J. Atmos. Chem.*, *36*, 157–230.
- Reeves, C. (2003), Atmospheric budget implications of the temporal and spatial trends in methyl bromide concentration, *J. Geophys. Res.*, *108*(D11), 4343, doi:10.1029/2002JD002943.
- Rudolph, J. (1995), The tropospheric distribution and budget of ethane, *J. Geophys. Res.*, *100*(D6), 11,369–11,381.
- Saito, T., Y. Yokouchi, S. Aoki, T. Nakazawa, Y. Fujii, and O. Watanabe (2006), A method for determination of methyl chloride concentration in air trapped in ice cores, *Chemosphere*, *63*, 1209–1213.
- Saltzman, E. S., M. Aydin, W. J. DeBruyn, D. B. King, and S. A. Yvon-Lewis (2004), Methyl bromide in preindustrial air: Measurements from an Antarctic ice core, *J. Geophys. Res.*, *109*(D05301), doi:10.1029/2003JD004157.
- Schwander, J., J. M. Barnola, C. Andrie, M. Leuenberger, A. Ludin, D. Raynaud, and B. Stauffer (1993), The age of the air in the firm and ice at Summit, Greenland, *J. Geophys. Res.*, *98*, 2831–2838.
- Schwander, J., T. Sowers, J.-M. Barnola, T. Blunier, A. Fuchs, and B. Malaize (1997), Age scale of the air in the summit ice: Implication for glacial-interglacial temperature change, *J. Geophys. Res.*, *102*(D16), 19,483–19,493.
- Shorter, J. H., C. E. Kolb, P. M. Crill, R. A. Kerwin, R. W. Talbot, M. E. Hines, and R. C. Harriss (1995), Rapid degradation of atmospheric methyl bromide in soils, *Nature*, *377*, 717–719.
- Sowers, T., and J. Jubenville (2000), A modified extraction technique for liberating occluded gases from ice cores, *J. Geophys. Res.*, *105*, 29,155–29,164.
- Sturges, W. T., H. P. McIntyre, S. A. Penkett, J. A. Chappellaz, J.-M. Barnola, R. Mulvaney, E. Atlas, and V. Stroud (2001a), Methyl bromide, other brominated methanes and methyl iodide in polar firm air, *J. Geophys. Res.*, *106*, 1595–1606.
- Sturges, W. T., S. A. Penkett, J. M. Barnola, J. Chappellaz, E. Atlas, and V. Stroud (2001b), A long-term measurement record of carbonyl sulfide (COS) in two hemispheres from firm air measurements, *Geophys. Res. Lett.*, *28*, 4095–4098.
- Swanson, A. L., N. J. Blake, J. E. Dibb, M. A. Albert, D. R. Blake, and F. S. Rowland (2002), Photochemically induced production of CH₃Br, CH₃I, C₂H₅I, ethene, and propene within surface snow, *Atmos. Environ.*, *36*, 2671–2682.
- Taylor, K. C., P. A. Mayewski, M. S. Twickler, and S. I. Whitlow (1996), Biomass burning recorded in the GISP2 ice core: A record from eastern Canada?, *Holocene*, *6*(1), 1–6.
- Trudinger, C. M., D. M. Etheridge, G. A. Sturrock, P. J. Fraser, P. B. Krummel, and A. McCulloch (2004), Atmospheric histories of halocarbons from analysis of Antarctic firm air: Methyl bromide, methyl chloride, chloroform, and dichloromethane, *J. Geophys. Res.*, *109*(D22310), doi:10.1029/2004JD004932.
- Tucker, B. J., P. J. Maroulis, and A. R. Bandy (1985), Free tropospheric measurements of CS₂ over a 45°N to 45°S latitude range, *Geophys. Res. Lett.*, *12*, 9–11.
- Walker, S. J., R. F. Weiss, and P. K. Salameh (2000), Reconstructed histories of the annual mean atmospheric mole fractions for the halocarbons CFC-11, CFC-12, CFC-113 and carbon tetrachloride, *J. Geophys. Res.*, *105*, 14,285–14,296.
- Warwick, N. J., J. A. Pyle, and D. E. Shallcross (2006), Global modeling of the Atmospheric Methyl Bromide Budget, *J. Atmos. Chem.*, *54*, 133–159.
- Watts, S. F. (2000), The mass budgets of carbonyl sulfide, dimethyl sulfide, carbon disulfide and hydrogen sulfide, *Atmos. Environ.*, *34*, 761–779.
- Worton, D. R. (2006), Alkyl nitrates (C1-C5), trihalomethanes and related compounds in contemporary air and air preserved in polar firm and ice, Ph.D. thesis, University of East Anglia, Norwich, UK.
- Yang, Q., P. A. Mayewski, S. I. Whitlow, M. S. Twickler, M. C. Morrison, R. W. Talbot, J. E. Dibb, and E. Linder (1995), Global perspective of nitrate flux in ice cores, *J. Geophys. Res.*, *100*, 5113–5121.
- Yokouchi, Y., D. Toom-Saundry, K. Yazawa, T. Inagaki, and T. Tamaru (2002), Recent decline of methyl bromide in the troposphere, *Atmos. Environ.*, *36*, 4985–4989.
- Yvon-Lewis, S. A., and J. H. Butler (1997), The potential effect of biological degradation on the lifetime of atmospheric CH₃Br, *Geophys. Res. Lett.*, *24*, 1227–1230.

M. Aydin, E. S. Saltzman, and M. B. Williams, Department of Earth System Science, University of California, Croul Hall 3206, Irvine, CA 92697, USA. (maydin@uci.edu)