

## Research 1395<sup>OA</sup> Unusually effective microRNA targeting within repeat-rich coding regions of mammalian mRNAs Michael Schnall-Levin, Olivia S. Rissland, Wendy K. Johnston, Norbert Perrimon, David P. Bartel, and Bonnie Berger Coactivation of GR and NFKB alters the repertoire of their binding sites and target genes 1404 Nagesha A.S. Rao, Melysia T. McCalman, Panagiotis Moulos, Kees-Jan Francoijs, Aristotelis Chatziioannou, Fragiskos N. Kolisis, Michael N. Alexis, Dimitra J. Mitsiou, and Hendrik G. Stunnenberg 1417 Alterations in TCF7L2 expression define its role as a key regulator of glucose metabolism Daniel Savic, Honggang Ye, Ivy Aneas, Soo-Young Park, Graeme I. Bell, and Marcelo A. Nobrega 1426 Histone H3 trimethylation at lysine 36 is associated with constitutive and facultative heterochromatin Sophie Chantalat, Arnaud Depaux, Patrick Héry, Sophie Barral, Jean-Yves Thuret, Stefan Dimitrov, and Matthieu Gérard Genome-scale analysis of metazoan replication origins reveals their organization in specific 1438 but flexible sites defined by conserved features Christelle Cayrou, Philippe Coulombe, Alice Vigneron, Slavica Stanojcic, Olivier Ganier, Isabelle Peiffer, Eric Rivals, Aurore Puy, Sabine Laurent-Chabalier, Romain Desprat, and Marcel Méchali 1450 Post-transcriptional generation of miRNA variants by multiple nucleotidyl transferases contributes to miRNA transcriptome complexity Stacia K. Wyman, Emily C. Knouf, Rachael K. Parkin, Brian R. Fritz, Daniel W. Lin, Lucas M. Dennis, Michael A. Krouse, Philippa J. Webster, and Muneesh Tewari Deep small RNA sequencing from the nematode Ascaris reveals conservation, functional 1462 diversification, and novel developmental profiles Jianbin Wang, Benjamin Czech, Amanda Crunk, Adam Wallace, Makedonka Mitreva, Gregory J. Hannon, and Richard E. Davis 1478 Transcriptome dynamics through alternative polyadenylation in developmental and environmental responses in plants revealed by deep sequencing Yingjia Shen, R.C. Venu, Kan Nobuta, Xiaohui Wu, Varun Notibala, Caghan Demirci, Blake C. Meyers, Guo-Liang Wang, Guoli Ji, and Qingshun Q. Li

Genome-wide detection of novel regulatory RNAs in <i>E. coli</i> Rahul Raghavan, Eduardo A. Groisman, and Howard Ochman	1487
Methods	
Accurate and comprehensive sequencing of personal genomes Subramanian S. Ajay, Stephen C.J. Parker, Hatice Ozel Abaan, Karin V. Fuentes Fajardo, and Elliott H. Margulies	1498 <sup>OA</sup>
Barcoding bias in high-throughput multiplex sequencing of miRNA Shahar Alon, Francois Vigneault, Seda Eminaga, Danos C. Christodoulou, Jonathan G. Seidman, George M. Church, and Eli Eisenberg	1506
Resources	
Cactus: Algorithms for genome multiple sequence alignment Benedict Paten, Dent Earl, Ngan Nguyen, Mark Diekhans, Daniel Zerbino, and David Haussler	1512 <sup>OA</sup>
A probabilistic disease-gene finder for personal genomes  Mark Yandell, Chad Huff, Hao Hu, Marc Singleton, Barry Moore, Jinchuan Xing, Lynn B. Jorde, and Martin G. Reese	1529 <sup>OA</sup>
Synthetic spike-in standards for RNA-seq experiments Lichun Jiang, Felix Schlesinger, Carrie A. Davis, Yu Zhang, Renhua Li, Marc Salit, Thomas R. Gingeras, and Brian Oliver	1543 <sup>OA</sup>
Integrative analysis of environmental sequences using MEGAN4 Daniel H. Huson, Suparna Mitra, Hans-Joachim Ruscheweyh, Nico Weber, and Stephan C. Schuster	1552 <sup>OA</sup>
Erratum	1561

OAOpen Access paper.



**Cover** A messenger RNA is descended on by numerous microRNAs that bind to a repeated sequence motif in its coding region. Short, highly repeated sequences with complementarity to microRNAs have arisen in a number of gene families as a result of evolutionary duplication processes. Genes with such repeats are frequently targeted by microRNAs outside of their 3' UTRs. (Cover illustration by Leslie Gaffney and Lauren Solomon, Broad Institute. [For details, see Schnall-Levin et al., pp. 1395–1403.])