

SERS analysis of bacterial strains: Escherichia coli and Staphylococcus epidermidis

Original

SERS analysis of bacterial strains: Escherichia coli and Staphylococcus epidermidis / Paccotti, Niccolò; Chiado', Alessandro; Novara, Chiara; Giorgis, Fabrizio; Geobaldo, Francesco; Boschetto, Francesco; Marin, Elia; Horiguchi, Satoshi; Pezzotti, Giuseppe. - ELETTRONICO. - (2019), pp. 152-152. (Intervento presentato al convegno I3S 7th International Symposium on Sensor Science tenutosi a Napoli nel 9-11/05/2019).

Availability:

This version is available at: 11583/2749053 since: 2021-03-19T10:53:26Z

Publisher:

MDPI

Published

DOI:

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Abstract

134. SERS Analysis of Bacterial Strains: *Escherichia coli* and *Staphylococcus epidermidis*

Niccolò Paccotti ^{1,*}, Alessandro Chiadò ¹, Chiara Novara ¹, Fabrizio Giorgis ¹, Francesco Geobaldo ¹,
Francesco Boschetto ^{2,3}, Elia Marin ^{2,3}, Satoshi Horiguchi ^{3,4} and Giuseppe Pezzotti ^{2,3,5,6}

¹ Department of Applied Science and Technology, Politecnico di Torino, Torino, Italy

² Ceramic Physics Laboratory, Kyoto Institute of Technology, Kyoto 606-8585, Japan

³ Department of Immunology, Kyoto Prefectural University of Medicine, Kyoto 602-0841, Japan

⁴ Department of Dental Medicine, Kyoto Prefectural University of Medicine, Kyoto 602-0841, Japan

⁵ Department of Orthopedic Surgery, Tokyo Medical University, Tokyo 160-8402, Japan

⁶ The Center for Advanced Medical Engineering and Informatics, Osaka University, Osaka 565-0871, Japan

* Correspondence: niccolo.paccotti@polito.it

Bacteria are prokaryotic microorganism whose pathogenic activity can affect every aspect of daily life. In this context, a powerful tool for their detection is provided by the Surface Enhanced Raman Spectroscopy (SERS), that can reveal the cellular composition of different bacterial strains with high sensitivity and intrinsic specificity. In this study, mesoporous silicon-based SERS substrates decorated with silver nanoparticles were used to investigate the vibrational pattern of *S. epidermidis* and *E. coli*, representative Gram-positive and Gram-negative bacteria, to provide a reproducible method to discriminate between distinct bacterial strains. At first, each spectrum was completely characterized in order to evaluate the contribution of each vibrational mode. The SERS spectra showed species-related features, arising from the different composition of the cell wall as well as their distinctive biofilm matrix and metabolic pathways. Furthermore, a life cycle analysis was carried out monitoring the evolution of the main SERS bands over time, analyzing the bacteria population after 12, 24 and 48 h of culture. The results pointed out an increase of the intensity of the bands after 24 h, while a successive decrease at 48 h was observed, in agreement with the bacterial growth profile.



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).