

## Multimedia Appendix 7

*This is a Multimedia Appendix to a full manuscript published in the J Med Internet Res. For full copyright and citation information see <http://dx.doi.org/10.2196/jmir.45754>.*

It was found that three narratives could not be recommended (i.e., did not appear in NarraGive's top-10 list for any participant) when NarraGive was trained on any combination of NEON Trial data, NEON-O Trial data, Hopefulness rating set, either of the two Similarity rating sets, and any of the three filtering algorithms.

Accesses to narratives 1, 2 and 3 are summarized in Table S1.

<b>Narrative</b>	<b>Hope</b>	<b>Similarity to Narrator</b>	<b>Similarity to Narrative</b>	<b>Learning</b>	<b>Empathy</b>	<b>Trial</b>
1	0	Not given	Not given	Not given	Not given	NEON
1	-1	Not given	Not given	Not given	Not given	NEON-O
1	-1	Not given	Not given	Not given	Not given	NEON-O
1	1	Not given	Not given	Not given	Not given	NEON
2	1	Not given	Not given	Not given	Not given	NEON
2	0	Not given	Not given	Not given	Not given	NEON-O
3	1	1	1	1	1	NEON-O
3	0	Not given	Not given	Not given	Not given	NEON-O
3	1	Not given	Not given	Not given	Not given	NEON-O
3	0	Not given	Not given	Not given	Not given	NEON

*Table S1 - Narrative ratings for the three unrecommendable narratives using synthetic narrative IDs to avoid identification.*

Each row represents a rating given by a participant. The table shows all ratings for the three narratives.