

Publisher's Note: Rheology of Semiflexible Bundle Networks with Transient Linkers
[Phys. Rev. Lett. 112, 238102 (2014)]

Kei W. Müller, Robijn F. Bruinsma, Oliver Lieleg, Andreas R. Bausch, Wolfgang A. Wall, and Alex J. Levine
(Received 14 July 2014; published 22 July 2014)

DOI: 10.1103/PhysRevLett.113.049901

PACS numbers: 87.15.A-, 83.60.Bc, 87.15.La, 87.16.Ln, 99.10.Fg

This Letter was published online on 10 June 2014 with a production error resulting in an incorrect reference citation in Fig. 2. Figure 2 has been replaced online as of 11 July 2014. The figure is incorrect in the printed version of the journal; therefore, for the benefit of the print readership, the figure has been replicated below.




FIG. 2 (color online). Comparison of simulated rheological spectra to experiments. Simulation parameters are $c_f = 4 \mu\text{M}$, $c_l/c_f = 0.017$, $k_{\text{on}} = 90 \text{ s}^{-1}$, and $k_{\text{off}} = \{2, 0.07\} \text{ s}^{-1}$. We encounter three distinct regimes. At high frequencies ($\omega > \omega_1$), we see scaling behaviors $G' \sim \omega^{0.93}$ and $G'' \sim \omega^{0.55}$. The inset shows G' and G'' of the same system without linkers. The expected $\omega^{3/4}$ scaling relation is recovered. At intermediate frequencies ($\omega_2 < \omega < \omega_1$), there is a local maximum in G'' and large sample-to-sample variations. At low frequencies, there is again a power-law regime.