

Scan A4121  
- 4123

Wagner A670

Reprint

add to 4 seqs

Ref ~~SEC 11 3 80~~  
Archie Mark

reprint

A4121  
-4123  
A670

f91  
indexed

ENUMERATION OF GENERALIZED WEAK ORDERS  
Carl G. Wagner\*

University of Tennessee, Knoxville, TN 37916

1. Introduction. Let  $R$  be a binary relation on  $X$  and let  
 $aR = \{(x, y): (x, y) \in R \text{ and } (y, x) \notin R\}$ ,  $cR = \{(x, y): (x, y) \notin R\}$ ,  
and  $sR = \{(x, y): (x, y) \in R \text{ and } (y, x) \in R\}$ . We term  $R$  negatively  
transitive when  $cR$  is transitive.

$R$  is called a generalized weak order if  $caR$  is transitive. Generalized weak orders were introduced by P.C. Fishburn [4], and include as special cases the asymmetric weak orders (asymmetric, negatively transitive relations) and complete weak orders (complete, reflexive, transitive relations) familiar to economists as models, respectively, of strict preference and preference-or-indifference.

Generalizing the familiar 1-1 correspondence between asymmetric weak orders, complete weak orders, and linearly ordered partitions, one may exhibit a 1-1 correspondence between generalized weak orders on a set and linearly ordered partitions of that set, the blocks of which are equipped with arbitrary symmetric relations [4, p. 165]. The essential details of this correspondence are as follows: A generalized weak order  $R$  on a set  $X$  partitions  $X$  by the equivalence relation  $scaR$ , and  $\hat{X} = X/scaR$  is linearly ordered by  $>$ , where for all  $A, B \in \hat{X}$ ,  $A > B$  iff  $(x, y) \in aR$  for all  $x \in A$  and  $y \in B$ . The symmetric relation attached to each  $A \in \hat{X}$  is simply the restriction of  $R$  to  $A$ . Conversely, if  $\tilde{X}$  is a partition of  $X$  linearly ordered by  $>$ , and each  $A \in \tilde{X}$  is equipped with a symmetric relation  $R_A$ , then  $R = R^{(s)} \cup R^{(a)}$  is a generalized weak order on  $X$ , where  $R^{(s)} = \bigcup_{A \in \tilde{X}} R_A$

and  $R^{(a)} = \bigcup_{\substack{A, B \in \tilde{X} \\ A > B}} A \times B$ . In the remainder of this paper we exploit the

foregoing correspondence to enumerate various classes of generalized weak orders.

\* Portions of this work were completed while the author was a Fellow at the Center for Advanced Study in the Behavioral Sciences, supported by grants from the National Science Foundation (BNS 76-22943 A 02), the Andrew W. Mellon Foundation, and the University of Tennessee.

2. Recurrence Relations. Let  $W(n)$  denote the number of generalized weak orders on an  $n$ -set, equivalently, the number of ordered partitions of an  $n$ -set, each block of which is equipped with an arbitrary symmetric relation. The number of such partitions with initial block of size  $k$  is

$\binom{n}{k} 2^{\binom{k+1}{2}} W(n-k)$ , and hence,

$$(1) \quad W(n) = \sum_{k=1}^n \binom{n}{k} 2^{\binom{k+1}{2}} W(n-k); \quad W(0) = 1.$$

A generalized weak order need not be transitive or negatively transitive. Indeed, the transitive (resp., negatively transitive) generalized weak orders are just those corresponding to ordered partitions with transitive (resp., negatively transitive) symmetric relations attached to each block [4, Theorems 5 and 6].

Since the map  $R \rightarrow cR$  is a bijection from transitive to negatively transitive generalized weak orders, we restrict attention to determining  $T(n)$ , the number of transitive generalized weak orders on an  $n$ -set. Proceeding as in the derivation of (1) above, we need to determine the number of transitive, symmetric relations on a block of size  $k$ . Now such a relation  $R$  on a block  $A$ , when restricted to  $\{x \in A: (x, x) \in R\}$ , is an equivalence. As is familiar, the number of equivalence relations on a  $j$ -set is given by  $B(j)$ , the  $j$ -th Bell number. Hence the number of transitive, symmetric relations on a  $k$ -set is  $\sum_{j=0}^k \binom{k}{j} B(j) = B(k+1)$ , by a familiar recurrence for the Bell numbers [6]. It follows that

$$(2) \quad T(n) = \sum_{k=1}^n \binom{n}{k} B(k+1) T(n-k); \quad T(0) = 1.$$

Next, let  $M(n)$  denote the number of generalized weak orders on an  $n$ -set which are both transitive and negatively transitive. (Remark. Fishburn [4, Lemma 5] has proved that transitivity of  $R$  and  $cR$  imply transitivity of  $caR$ . Hence  $M(n)$  is, more simply, the number of transitive, negatively transitive relations on an  $n$ -set.) It is easy to see that a transitive, negatively transitive, symmetric relation on a block is either the empty or universal relation on that block. Hence, arguing as in the case of (1) and (2) above, it follows that

$$(3) \quad M(n) = \sum_{k=1}^n 2 \binom{n}{k} M(n-k) ; M(0) = 1 .$$

Finally, let  $P(n)$  denote the number of asymmetric (equivalently, complete) weak orders on an  $n$ -set. As noted above,  $P(n)$  is simply the number of linearly ordered partitions on an  $n$ -set. (In terms of the correspondence which we have exploited above, asymmetric weak orders correspond to linearly ordered partitions whose blocks are all equipped with the empty relation, and complete weak orders with those whose blocks are all equipped with the universal relation.) By a now familiar argument, we have

$$(4) \quad P(n) = \sum_{k=0}^n \binom{n}{k} P(n-k) ; P(0) = 1 ,$$

which is derived in [5] by a more elaborate argument.

3. Generating Functions. Let  $F$  and  $G$  be real valued arithmetic functions on the non-negative integers, and let

$$F + G(n) = F(n) + G(n)$$

and

$$F * G(n) = \sum_{k=0}^n \binom{n}{k} F(k) G(n-k) .$$

Equipped with these operations, the set of arithmetic functions is an integral domain with multiplicative identity  $I$ , where  $I(0) = 1$  and  $I(n) = 0$  if  $n > 0$ .  $F$  has a multiplicative inverse iff  $F(0) \neq 0$ . As is familiar, this ring of arithmetic functions is isomorphic with the ring of formal exponential series

$$\sum_{n=0}^{\infty} a_n \frac{x^n}{n!}$$

equipped with ordinary addition and multiplication, the isomorphism being given by the mapping  $\psi$ , where

$$\psi(F) = \sum_{n=0}^{\infty} F(n) \frac{x^n}{n!}$$

is the exponential generating function of the arithmetic function  $F$ . We determine in this section the exponential generating functions of  $T$ ,  $M$ , and  $P$ .

Define arithmetic functions  $Z$ ,  $\bar{B}$ , and  $E$  by  $Z(n) = 1$  for all  $n \geq 0$ ,  $\bar{B}(n) = B(n+1)$ , the  $n+1$ <sup>st</sup> Bell number, and  $E(n) = 2^{\binom{n+1}{2}}$ . Then we may express the recurrence relations (1), (2), (3), and (4) as equations in the ring of arithmetic functions as follows:

$$(5) \quad W = (E - I) * W + I$$

$$(6) \quad T = (\bar{B} - I) * T + I$$

$$(7) \quad M = (2Z - 2I) * M + I$$

$$(8) \quad P = (Z - I) * P + I.$$

Solving the above for  $W$ ,  $T$ ,  $M$ , and  $P$  we have

$$(9) \quad W = I/(2I - E)$$

$$(10) \quad T = I/(2I - \bar{B})$$

$$(11) \quad M = I/(3I - 2Z)$$

$$(12) \quad P = I/(2I - Z).$$

The numbers  $W(n)$  grow too rapidly to possess a non-trivially convergent exponential generating function. In order to determine  $\psi(T)$ , we need to determine

$$\psi(\bar{B}) = \sum_{n=0}^{\infty} B(n+1) \frac{x^n}{n!}.$$

Now [6],

$$\psi(B) = \sum_{n=0}^{\infty} B(n) \frac{x^n}{n!} = e^{e^x - 1},$$

and since  $\psi(\bar{B}) = D_x \psi(B)$ , it follows that  $\psi(\bar{B}) = e^{e^x + x - 1}$ . Since  $\psi(I) = 1$ , (10) implies that

$$(13) \quad \psi(T) = \sum_{n=0}^{\infty} T(n) \frac{x^n}{n!} = (2 - e^{e^x + x - 1})^{-1}.$$

And, since  $\psi(Z) = e^x$ , (11) and (12) imply that

$$(14) \quad \psi(M) = \sum_{n=0}^{\infty} M(n) \frac{x^n}{n!} = (3 - 2e^x)^{-1}$$

and

$$(15) \quad \psi(P) = \sum_{n=0}^{\infty} P(n) \frac{x^n}{n!} = (2 - e^x)^{-1},$$

the latter result having been derived by a more elaborate argument in [5].

4. Dobinski-type Formulas. Over a century ago Dobinski [3] established the following beautiful formula for the Bell numbers:

$$B(n+1) = \frac{1}{e} \sum_{k=1}^{\infty} \frac{k^n}{(k-1)!}.$$

In this section we establish Dobinski-type formulas for  $P(n)$ ,  $M(n)$ , and  $T(n)$ .

For  $\lambda \neq 1$ , define  $H_n[\lambda]$  by

$$(16) \quad \frac{\lambda - 1}{\lambda - e^x} = \sum_{n=0}^{\infty} H_n[\lambda] \frac{x^n}{n!}.$$

The numbers  $H_n[\lambda]$  are called Eulerian numbers. (See [1], [2].) From (14) and (15) it follows that  $M(n) = H_n[3/2]$  and  $P(n) = H_n[2]$ . A Dobinski-type formula for  $H_n[\lambda]$  may be derived as follows:

$$\begin{aligned} H_n[\lambda] &= (\lambda - 1) D^n (\lambda - e^x)^{-1} \Big|_{x=0} = \frac{\lambda-1}{\lambda} D^n (1 - \frac{1}{\lambda} e^x)^{-1} \Big|_{x=0} \\ (17) \quad &= \frac{\lambda-1}{\lambda} D^n \sum_{k=0}^{\infty} (\frac{1}{\lambda} e^x)^k \Big|_{x=0} = \frac{\lambda-1}{\lambda} \sum_{k=0}^{\infty} D^n (\frac{1}{\lambda} e^x)^k \Big|_{x=0} \\ &= \frac{\lambda-1}{\lambda} \sum_{k=0}^{\infty} \lambda^{-k} k^n. \end{aligned}$$

In particular,

$$(18) \quad M(n) = H_n[3/2] = \frac{1}{3} \sum_{k=0}^{\infty} (\frac{2}{3})^k k^n$$

and

$$(19) P(n) = H_n[2] = \frac{1}{2} \sum_{k=0}^{\infty} \left(\frac{1}{2}\right)^k k^n$$

the latter result having appeared in [5].

As for the numbers  $T(n)$ , we have by (13),

$$T(n) = \frac{1}{2} D^n (1 - \frac{1}{2} e^{e^x+x-1})^{-1} \Big|_{x=0} = \frac{1}{2} D^n \sum_{k=0}^{\infty} \left(\frac{1}{2} e^{e^x+x-1}\right)^k \Big|_{x=0}$$

$$(20) = \frac{1}{2} \sum_{k=0}^{\infty} \left(\frac{1}{2}\right)^k D^n (e^{ke^x+kx-k}) \Big|_{x=0}$$

$$= \frac{1}{2} \sum_{k=0}^{\infty} \left(\frac{1}{2}\right)^k f_n(k),$$

where  $f_n(k) = D^n (e^{ke^x+kx-k}) \Big|_{x=0}$ . Now,  $f_0(k) = 1$ ,  $f_1(k) = 2k$ ,  $f_2(k) = 4k^2 + k$ ,  $f_3(k) = 8k^3 + 6k^2 + k$ , and in general,  $f_n(k)$  is a polynomial in  $k$  of degree  $n$ , as seen from the recurrence

$$(21) f_n(k) = \sum_{j=0}^{n-2} \binom{n-1}{j} k f_j(k) + 2k f_{n-1}(k),$$

which may be derived by writing  $f_n(k) = D^{n-1} (e^{ke^x+kx-k} (ke^x + k)) \Big|_{x=0}$ , and by applying Newton's formula.

5. Tables. Initial values of  $W(n)$ ,  $T(n)$ ,  $M(n)$  and  $P(n)$ , calculated from (1), (2), (3), and (4), appear below.

$n$	$W(n)$	$T(n)$	$M(n)$	$P(n)$
1	2	2	2	1
2	16	13	10	3
3	208	123	74	13
4	3,968	1,546	730	75
5	109,568	24,283	9,002	541
6	4,793,344	457,699	133,210	4,683
7	*	10,064,848	2,299,754	47,293
8	*	252,945,467	45,375,130	545,835

A670

omit

new

new

new

have

We remark in conclusion that the numbers  $M(n)$  and  $P(n)$  satisfy some interesting congruences. It follows from (16) that

Please enter 3.

Name: Generalized weak orders.

Ref: SEC 11 ? 80.

$$(22) \quad H_n[\lambda] = \sum_{k=0}^n (\lambda - 1)^{-k} \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} j^n [1] .$$

Hence,  $H_n[\lambda]$  is integral for all  $n$  iff  $(\lambda - 1)^{-1}$  is integral. If  $(\lambda - 1)^{-1}$  is integral, it follows from (22) and Fermat's Theorem that, for  $p$  an odd prime,

$$(23) \quad H_{n+p-1}[\lambda] \equiv H_n[\lambda] \pmod{2p(\lambda - 1)^{-2}}$$

Since  $H_n[3/2] = M(n)$  and  $H_n[2] = P(n)$ , it follows from

(23) that for  $p$  an odd prime,

$$(24) \quad M(n + p - 1) \equiv M(n) \pmod{8p}$$

and

$$(25) \quad P(n + p - 1) \equiv P(n) \pmod{2p} ,$$

the latter result appearing in [5].

#### References

1. L. Carlitz, Eulerian Numbers and Polynomials, Mathematics Magazine 33(1959), 247-260.
2. L. Carlitz, Extended Bernoulli and Eulerian Numbers, Duke Mathematical Journal 31(1964), 667-689.
3. G. Dobinski, Grunert's Archiv 61(1877), 333-336.
4. P. C. Fishburn, Transitivity, The Review of Economic Studies 46(1), No. 142(1979), 163-173.
5. A. O. Gross, Preferential Arrangements, American Mathematical Monthly 69(1962), 4-8.
6. Gian-Carlo Rota, The Number of Partitions of a Set. American Mathematical Monthly 71(1963), 498-504.