

Datasheet for ABIN934717
Thyroperoxidase Protein

2 Validations

2 Publications



[Go to Product page](#)

Overview

Quantity: 100 µg

Target: Thyroperoxidase (TPO)

Origin: Human

Source: Human

Protein Type: Native

Product Details

Characteristics: Purified native Human Thyroid Peroxidase protein
Protein Source: Human thyroid glands

Purity: > 95 % pure

Target Details

Target: Thyroperoxidase (TPO)

Alternative Name: Thyroid Peroxidase ([TPO Products](#))

Background: Thyroid peroxidase or thyroperoxidase (TPO) is an enzyme expressed mainly in the thyroid that liberates iodine for addition onto tyrosine residues on thyroglobulin for the production of thyroxine (T4) or triiodothyronine (T3), thyroid hormones. In humans, thyroperoxidase is encoded by the TPO gene.

Description: Human thyroid glands.

Alternative Names: TPO protein, MSA protein, TPX protein

Pathways: [Thyroid Hormone Synthesis](#)

Application Details

Application Notes: Each Investigator should determine their own optimal working dilution for specific applications.

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Reconstitute in distilled water.

Buffer: Lyophilized in 20 mM TRIS, with 50 mM KCl, no preservatives.

Preservative: Without preservative

Handling Advice: Avoid repeated freeze/thaw cycles.

Storage: -20 °C

Storage Comment: Aliquot and store at -20 °C.

Publications

Product cited in: Ali, Wolfe, Mukherjee, Radford, Patel, White, Milojevic, Capretta, Nair, Brennan: "A sputum bioassay for airway eosinophilia using an eosinophil peroxidase aptamer." in: **Scientific reports**, Vol. 12, Issue 1, pp. 22476, (2022) ([PubMed](#)).

Rahnama, Mahmoudi, Kazemnejad, Salehi, Ghahiri, Soltanghoraee, Vafaei, Rezaei, Zarnani: "Thyroid peroxidase in human endometrium and placenta: a potential target for anti-TPO antibodies." in: **Clinical and experimental medicine**, (2020) ([PubMed](#)).



Successfully validated (SDS-PAGE (SDS))


by [in.vent Diagnostica GmbH](#), Hennigsdorf, Germany

Report Number: 102793

Date: Apr 17 2018

Target:	TPO
Lot Number:	A17091521
Method validated:	SDS-PAGE (SDS)
Positive Control:	Molecular weight (Uniprot P07202: 8 possible isoforms of TPO with a molecular weight between 103 and 67 kDa)
Negative Control:	none
Notes:	Passed. Native human TPO protein ABIN934717 appears as a band with a molecular weight of approximately 100kDa after separation on an SDS-PAGE gel.
Primary Antibody:	ABIN934717
Protocol:	<ul style="list-style-type: none">• Dilute 2µg of native human TPO (antibodies-online, ABIN934717, lot A17091521) in 2x sample buffer (126mM Tris-HCl pH6.8, 20% glycerol, 4% SDS, 0.02% bromophenol blue, 0.1M DTT).• Boil samples for 5min at 95°C.• Separate the sample on a denaturing SERVAGel TG PRIME 12% precast gel (Serva, 43266, V170144) in a Electrophoresis chamber (Hoefer Inc., SE250) with Laemmli buffer for 1h at 250V, 50mA/gel along side 4µl Roti-Mark Tricolor (Carl Roth, 8271, lot 326248051) molecular weight marker.• Staining of the separated proteins with InstantBlue Protein Stain (Expedeon, ISB1L, lot 170616330) according to the manufacturer's recommendations.• Image acquisition: 600dpi scan and increasing of color saturation.
Experimental Notes:	The manufacturer's information concerning the protein's purity and molecular weight can be confirmed.

**Validation image no. 1 for Thyroid Peroxidase (TPO)
protein (ABIN934717)**



Coomassie staining subsequently to SDS-PAGE separation
of human TPO1 ABIN934717.



Successfully validated (Lipid Interaction Assay (LIA))

by [in.vent Diagnostica GmbH, Hennigsdorf, Germany](#)

Report Number: 103248

Date: Apr 17 2018

Target:	TPO
Lot Number:	A17091521
Method validated:	Lipid Interaction Assay (LIA)
Positive Control:	Protein concentration (manufacturer's specification) Standards and controls for LIA based of purified TPO out of human thyroid
Negative Control:	Dilution Buffer for LIA Deionized water for absorbance at 280nm
Notes:	Passed. Protein concentration of native human TPO protein ABIN934717 can be confirmed by absorbance at 280nm and by LIA.
Primary Antibody:	ABIN934717
Protocol:	<ul style="list-style-type: none">• Measuring protein concentration using absorbance at 280nm:<ul style="list-style-type: none">◦ 100µl sample in a 96-Well UV Microplate (Thermo Fisher Scientific, 8404)◦ Measuring at 280nm◦ Calculation with d of 0.28cm◦ Extinction coefficient assuming all pairs of Cys residues form cysteines: Abs 0.1% (=1g/l)= 1.304• Measuring TPO concentration using an in-house coated tube LIA:<ul style="list-style-type: none">◦ Sample dilution with assay dilution buffer in 6 steps (10000ng/ml and 100ng/ml as pre-dilution; 50, 25, 12.5, 6.25ng/ml as samples)◦ Pipette 300µl tracer and 200µl standards, controls, or samples per tube in duplicates.◦ Incubation ON at RT.◦ Add 1ml washing solution to each coated tube prior to decanting off the liquid.◦ Add 1 ml washing solution to each coated tube three times and decant off the liquid. Turn the tubes upside down for 5–10min and adsorb remaining liquid with blotting paper.◦ Measurement in a luminometer by automatic injection of LIA reagents (sodium hydrate, azotic acid). Measurement time: 1sec.• Calculate the concentration based on the standard curve.
Experimental Notes:	<ul style="list-style-type: none">• The LIA measures human TPO in serum, plasma and other fluids, using one anti-TPO antibody at solid phase and another anti-TPO antibody as tracer. The measurement range is

