



# Pharmacological characterization of human recombinant melatonin $mt_1$ and $MT_2$ receptors

\*<sup>1</sup>Christopher Browning, <sup>1</sup>Isabel Beresford, <sup>2</sup>Neil Fraser & <sup>1</sup>Heather Giles

<sup>1</sup>Receptor Pharmacology Glaxo Wellcome Medicines Research Centre, Gunnels Wood Road, Stevenage, Herts, SG1 2NY and

<sup>2</sup>Molecular Pharmacology, Glaxo Wellcome Medicines Research Centre, Gunnels Wood Road, Stevenage, Herts, SG1 2NY

**1** We have pharmacologically characterized recombinant human  $mt_1$  and  $MT_2$  receptors, stably expressed in Chinese hamster ovary cells (CHO- $mt_1$  and CHO- $MT_2$ ), by measurement of [<sup>3</sup>H]-melatonin binding and forskolin-stimulated cyclic AMP (cAMP) production.

**2** [<sup>3</sup>H]-melatonin bound to  $mt_1$  and  $MT_2$  receptors with  $pK_D$  values of 9.89 and 9.56 and  $B_{max}$  values of 1.20 and 0.82 pmol  $mg^{-1}$  protein, respectively. Whilst most melatonin receptor agonists had similar affinities for  $mt_1$  and  $MT_2$  receptors, a number of putative antagonists had substantially higher affinities for  $MT_2$  receptors, including luzindole (11 fold), GR128107 (23 fold) and 4-P-PDOT (61 fold).

**3** In both CHO- $mt_1$  and CHO- $MT_2$  cells, melatonin inhibited forskolin-stimulated accumulation of cyclic AMP in a concentration-dependent manner ( $pIC_{50}$  9.53 and 9.74, respectively) causing 83 and 64% inhibition of cyclic AMP production at 100 nM, respectively. The potencies of a range of melatonin receptor agonists were determined. At  $MT_2$  receptors, melatonin, 2-iodomelatonin and 6-chloromelatonin were essentially equipotent, whilst at the  $mt_1$  receptor these agonists gave the rank order of potency of 2-iodomelatonin > melatonin > 6-chloromelatonin.

**4** In both CHO- $mt_1$  and CHO- $MT_2$  cells, melatonin-induced inhibition of forskolin-stimulated cyclic AMP production was antagonized in a concentration-dependent manner by the melatonin receptor antagonist luzindole, with  $pA_2$  values of 5.75 and 7.64, respectively. Melatonin-mediated responses were abolished by pre-treatment of cells with pertussis toxin, consistent with activation of  $G_i/G_o$  G-proteins.

**5** This is the first report of the use of [<sup>3</sup>H]-melatonin for the characterization of recombinant  $mt_1$  and  $MT_2$  receptors. Our results demonstrate that these receptor subtypes have distinct pharmacological profiles.

*British Journal of Pharmacology* (2000) **129**, 877–886

**Keywords:** Melatonin;  $mt_1$  receptor;  $MT_2$  receptor; [<sup>3</sup>H]-melatonin; 2-[<sup>125</sup>I]-iodomelatonin; receptor classification; subtypes; cyclic AMP; luzindole

**Abbreviations:** cAMP, cyclic adenosine 3':5' monophosphate; CHO, Chinese hamster ovary; 5-HT, 5-hydroxytryptamine, MT, melatonin; 8-M-PDOT, 8-methoxy-2-propionamidotetraline; 4-P-PDOT, 4-phenyl-2-propionamidotetraline

## Introduction

Melatonin is the principle hormone of the pineal gland and is believed to have a central role in the regulation of the mammalian circadian system and reproductive function in seasonally breeding animals (for a review see Arendt, 1995). To date, two human melatonin receptors have been cloned, termed  $mt_1$  and  $MT_2$ , and have been shown to be seven transmembrane G-protein coupled receptors (Reppert *et al.*, 1994; 1995). When expressed in immortalized mammalian cell lines, both  $mt_1$  and  $MT_2$  receptors bind melatonin with high affinity and couple to the inhibition of adenylate cyclase (Reppert *et al.*, 1994; 1995).

Radioligand binding studies using 2-[<sup>125</sup>I]-iodomelatonin have identified high affinity binding sites in rat and human suprachiasmatic nucleus, sheep pars tuberalis and chicken retina (for a review see Morgan *et al.*, 1994). *In situ* hybridization studies have shown the presence of  $mt_1$  mRNA in the human suprachiasmatic nucleus (Weaver & Reppert, 1996), whilst  $MT_2$  mRNA is found in human retina and brain (Reppert *et al.*, 1995). Subtype selective agonists and antagonists would greatly assist the elucidation of the relative contributions of melatonin receptor subtypes

to the physiological actions of melatonin, such as its effects in the suprachiasmatic nucleus to regulate the circadian system. The expression of  $mt_1$  and  $MT_2$  receptors as isolated, single populations of receptors allows the investigation of melatonin receptor pharmacology and the search for subtype selective ligands which cannot be achieved in physiological systems where mixed populations of melatonin receptors may exist.

Thus, we have stably expressed  $mt_1$  and  $MT_2$  receptors in Chinese hamster ovary (CHO) cells and characterized the pharmacology of these receptors using both radioligand binding assays and measurement of the inhibition of forskolin-stimulated cyclic adenosine 3':5' monophosphate (cyclic AMP). Previous melatonin receptor binding studies have generally employed 2-[<sup>125</sup>I]-iodomelatonin as a radioligand. However, high receptor expression levels have enabled us to develop a radioligand binding assay utilising [<sup>3</sup>H]-melatonin. This radioligand has advantages over 2-[<sup>125</sup>I]-iodomelatonin, as it is safer, more economical to use, and is chemically identical to the endogenous hormone. We have used a number of melatonin receptor agonists, including the indolene GR196429 (Beresford *et al.*, 1998a) and the naphthalenic compound S-20098 (Yous *et al.*, 1992) to characterize CHO- $mt_1$  and CHO- $MT_2$  receptors. In addition,

\*Author for correspondence at: National Institute for Medical Research, The Ridgeway, Mill Hill, London, NW7 1AA.

a number of putative melatonin receptor partial agonists and antagonists with selectivity for the  $MT_2$  receptor were evaluated, including GR128107, 5-MCA-NAT, 8-M-PDOT and 4-P-PDOT (Dubocovich *et al.*, 1997). The most selective of these compounds, the amidotetraline 4-P-PDOT, was compared to the previously characterized melatonin receptor antagonist luzindole (Dubocovich, 1988) for its ability to antagonize melatonin mediated inhibition of forskolin-stimulated cyclic AMP production. Preliminary accounts of this work have already been presented (Browning *et al.*, 1997; 1998).

The nomenclature and classification of melatonin receptors used here was recently approved by the Nomenclature Committee of the International Union of Pharmacology (Dubocovich *et al.*, 1998a). The denomination 'mt<sub>1</sub>' corresponds to that of the recombinant receptor previously termed Mel<sub>1a</sub>.  $MT_2$  refers to native functional receptors with pharmacological characteristics similar to that of the recombinant receptor mt<sub>2</sub>, previously termed Mel<sub>1b</sub>.  $MT_3$  corresponds to the pharmacologically defined melatonin receptor subtype, with unknown gene sequence, previously referred to as ML<sub>2</sub>.

## Methods

### Generation of CHO-mt<sub>1</sub> and CHO-MT<sub>2</sub> cells

Clones representing the sequence of the human mt<sub>1</sub> receptor were amplified using degenerate primers based on the sequence of the *Xenopus laevis* melatonin receptor (Ebisawa *et al.*, 1994). These clones defined the genomic sequence flanking the intron within the coding sequence of the receptor and enabled the coding exons to be amplified from genomic DNA independently and reassembled in frame using an engineered site. Cloning was in pBluescript (Stratagene). The human  $MT_2$  sequence was cloned in a similar manner using the sequence described by Reppert and co-workers (1995). The nucleotide sequences obtained encoded for the same amino acid sequences as described by Reppert *et al.* (1994; 1995). The sequences encoding the receptor were cloned into the mammalian expression vector pcDNA3 (Invitrogen) and introduced into CHO cells by conventional calcium phosphate precipitation techniques, which were then placed under G418 selection (1 mg ml<sup>-1</sup>). Several G418 resistant mt<sub>1</sub> and  $MT_2$  cell lines were selected for [<sup>3</sup>H]-melatonin saturation binding studies and measurement of melatonin-mediated inhibition of forskolin-stimulated cyclic AMP production. The cell-lines which gave the highest levels of [<sup>3</sup>H]-melatonin binding and greatest melatonin-induced inhibition of cyclic AMP accumulation were chosen for further [<sup>3</sup>H]-melatonin radioligand binding and cyclic AMP studies, respectively.

### Preparation of cell membranes

CHO cells stably expressing human mt<sub>1</sub> or  $MT_2$  receptors were maintained in DMEM-F12 medium, supplemented with 10% (v v<sup>-1</sup>) foetal calf serum, 2 mM L-glutamine, 1 mg ml<sup>-1</sup> G418 and 400 µg ml<sup>-1</sup> hygromycin. Cells were grown at 37°C in 5% CO<sub>2</sub> in air. Membranes were prepared by harvesting CHO cells using Hanks balanced salt solution, containing EDTA (5 mM). The suspension was homogenized and centrifuged at 4500 × g for 35 min. The pellet was re-homogenized, resuspended in (mM): Tris-HCl buffer 50, containing MgCl<sub>2</sub> 2, EDTA 1, 0.1% ascorbic acid, pH 7.4, and aliquots (~2 mg protein ml<sup>-1</sup>) stored at -80°C until use.

### Radioligand binding assays

The assay was performed in 96-well plates in a final assay volume of 500 µl. Drugs and [<sup>3</sup>H]-melatonin (0.3 nM) or 2-[<sup>125</sup>I]-iodomelatonin (50 pM) were incubated with membranes (50 µg protein ml<sup>-1</sup>) for 120 min at 37°C in a buffer comprising (mM): Tris-HCl 50, MgCl<sub>2</sub> 2, EDTA 1, 0.1% ascorbic acid, pH 7.4. At the concentration of protein used, less than 10% of added radioligand was bound to membrane (data not shown). Non-specific binding was defined with melatonin (1 µM). For association studies, membranes (50 µg ml<sup>-1</sup>) were added to [<sup>3</sup>H]-melatonin (0.3 nM) and incubated for increasing periods of time (2–180 min). For dissociation studies, membranes were incubated with [<sup>3</sup>H]-melatonin (0.3 nM) for 2 h prior to the addition of cold melatonin (1 µM) to initiate dissociation, and then incubated for increasing periods of time (0–240 min). Bound radioactivity was separated by rapid filtration through GF/B filter paper using a Brandel or Wallac cell harvester and filters were washed with 4 × 1 ml Tris-HCl (50 mM; pH 7.4). Radioactivity was measured by liquid scintillation spectrometry.

### cyclic AMP assays

Cells were cultured as described above. The assay was performed in 96-well plates in a final assay volume of 200 µl. Confluent CHO-mt<sub>1</sub> or CHO-MT<sub>2</sub> cells were incubated at 37°C with DMEM-F12, containing 300 µM isobutylmethylxanthine (IBMX) to inhibit phosphodiesterase activity. Following 60 min incubation, agonist (0.01 pM–100 µM) was added. Sixty minutes later, forskolin (30 µM) was added and cells were incubated for a further 15 min. For antagonist experiments, melatonin was co-incubated with luzindole (0.1–100 µM) for 60 min prior to addition of forskolin. The reaction was terminated by removal of media and addition of ice-cold ethanol (100 µl) for 30 min at 4°C. Ethanol samples were evaporated to dryness and cyclic AMP concentrations determined by [<sup>125</sup>I]-cyclic AMP scintillation proximity assay (Amersham).

### Data analysis

Saturation binding experiments were analysed by fitting the sum of a hyperbola and a straight line to the total binding data to generate pK<sub>D</sub> and B<sub>max</sub> values. Competition binding data were analysed by fitting with a four-parameter logistic equation. pIC<sub>50</sub> values were converted to pK<sub>i</sub> values using the equation of Cheng & Prusoff (1973). Association kinetic data were analysed by fitting total binding data to the equation  $B = C + B_{max} \cdot (1 - \exp^{-k \cdot t})$ , where B = binding at time t, C = non-specific binding and k = observed association rate constant. Dissociation kinetic data were analysed by fitting total binding data to the equation  $B = C + B_{max} \cdot \exp^{-k \cdot t}$ , where k = dissociation rate constant.

Cyclic AMP data were fitted with a four parameter logistic equation to determine pIC<sub>50</sub> values and Hill coefficients. Drug responses were expressed as percentage inhibition of forskolin-stimulated cyclic AMP. The potency ratio was defined as the ratio of the EC<sub>50</sub> of the drug relative to that of melatonin determined in the same experiment. The pA<sub>2</sub> values for luzindole were determined using a modified form of the Schild equation (Black *et al.*, 1985; Lew & Angus, 1995).

All data are expressed as arithmetic mean ± s.e.mean. Statistical comparisons were made using two-tailed Student's *t*-test. For luzindole studies, the upper and lower asymptotes

and Hill slopes of the melatonin concentration-effect curves were compared using one-way analysis of variance (ANOVA) followed by Dunnett's test for significance. Where data are presented as the mean of separate experiments, curves were generated from the mean of the individual curve fitting parameter estimates.

### Drugs

GR196429, S20098, 5-MCA-NAT (GR135531), GR128107 and luzindole were synthesized by Medicinal Chemistry, Glaxo Wellcome. Melatonin, 6-hydroxymelatonin, 6-chloromelatonin, N-acetyl-5-HT, 5-HT, forskolin and IBMX were supplied by Sigma. 2-iodomelatonin, 4-phenyl-2-propionamidotetraline (4-P-PDOT) and 8-methoxy-2-propionamidotetraline (8-M-PDOT) were purchased from RBI. [ $^3$ H]-melatonin (85 Ci mol $^{-1}$ ) and 2-[ $^{125}$ I]-iodomelatonin ( $\sim 2000$  Ci mol $^{-1}$ ) were supplied by Amersham.


## Results

### Kinetics and saturation of [ $^3$ H]-melatonin binding to $mt_1$ and $MT_2$ receptors

At 37°C the specific binding of [ $^3$ H]-melatonin (0.4 nM) to  $mt_1$  receptors reached equilibrium within 15–30 min (Figure 1A;  $t_{1/2}$   $3.9 \pm 1.2$  min; observed association rate constant ( $k_{obs}$ )  $3.89 \times 10^8$  M $^{-1}$ .min $^{-1}$ ;  $n=3$ ) and remained stable for more than

360 min. Binding of [ $^3$ H]-melatonin (0.4 nM) to  $MT_2$  receptors occurred more slowly, taking 60–90 min to reach equilibrium (Figure 1B;  $t_{1/2}$   $14 \pm 2$  min;  $k_{obs}$   $9.18 \times 10^7$  M $^{-1}$ .min $^{-1}$ ;  $n=3$ ). For both receptors, an incubation time of 120 min was used routinely. Specific [ $^3$ H]-melatonin binding (0.4 nM) to  $mt_1$  receptors dissociated rapidly to non-specific binding levels (Figure 1C), with a dissociation rate constant ( $k_{-1}$ ) of  $0.056 \pm 0.003$  min $^{-1}$  and half-time of  $12.5 \pm 0.8$  min ( $n=3$ ). In contrast, binding of [ $^3$ H]-melatonin (0.4 nM) to  $MT_2$  receptors dissociated more slowly (Figure 1D), with a dissociation rate constant ( $k_{-1}$ ) of  $0.011 \pm 0.002$  min $^{-1}$  and half-time of  $64 \pm 8$  min ( $n=3$ ). Whilst only  $3.6 \pm 0.6\%$  ( $n=3$ ) of [ $^3$ H]-melatonin remained specifically bound to  $mt_1$  receptors after 240 min,  $22 \pm 2\%$  ( $n=3$ ) of the specific [ $^3$ H]-melatonin remained bound to  $MT_2$  receptors, which was significantly different from non-specific binding ( $P < 0.05$ ). From the kinetic data,  $pK_D$  values of  $9.78 \pm 0.16$  and  $9.92 \pm 0.03$  ( $n=3$ ) for  $mt_1$  and  $MT_2$  receptors, respectively, were determined.

Saturation binding studies indicated that [ $^3$ H]-melatonin bound in a specific, concentration-dependent and saturable manner to both  $mt_1$  and  $MT_2$  receptors (Figure 2A,B), with  $pK_D$  values of  $9.89 \pm 0.13$  and  $9.56 \pm 0.03$ , respectively ( $n=3$ ).  $B_{max}$  values were  $1.20 \pm 0.10$  and  $0.82 \pm 0.06$  pmol mg $^{-1}$  protein, respectively. Hill coefficients were  $1.02 \pm 0.13$  and  $1.06 \pm 0.06$ , respectively, suggesting interactions with single populations of binding sites. In the cell lines selected for cyclic AMP assays, [ $^3$ H]-melatonin saturation binding studies yielded  $B_{max}$  values of  $0.39 \pm 0.07$  and  $0.23 \pm 0.06$  pmoles mg $^{-1}$  protein, and  $pK_D$  values of



**Figure 1** Time-courses of association (A, B) and dissociation (C, D) of [ $^3$ H]-melatonin binding to the human recombinant  $mt_1$  (A, C) and  $MT_2$  (B, D) receptors. Data are the mean of duplicate points and representative of three similar experiments. For association experiments, upper asymptote (mean  $\pm$  s.e.mean) =  $0.37 \pm 0.05$  pmol mg $^{-1}$  ( $mt_1$ ) and  $0.39 \pm 0.01$  pmol mg $^{-1}$  ( $MT_2$ ). For dissociation experiments, binding at time  $t_0$  (mean  $\pm$  s.e.mean) =  $0.38 \pm 0.03$  pmol mg $^{-1}$  ( $mt_1$ ) and  $0.40 \pm 0.03$  pmol mg $^{-1}$  ( $MT_2$ ).

$9.43 \pm 0.06$  and  $9.04 \pm 0.18$  at  $mt_1$  and  $MT_2$  receptors respectively ( $n = 3$ ).

#### Comparison of [ $^3H$ ]-melatonin and 2-[ $^{125}I$ ]-iodomelatonin binding to $mt_1$ receptors

There was excellent agreement between the abilities of a range of melatonin analogues to compete for [ $^3H$ ]-melatonin and 2-[ $^{125}I$ ]-iodomelatonin binding to  $mt_1$  receptors (Table 1). Linear regression analysis yielded a correlation coefficient of 0.99 and a slope of  $1.01 \pm 0.05$  ( $n = 8$ ).


#### Pharmacological characterization of [ $^3H$ ]-melatonin binding to $mt_1$ and $MT_2$ receptors

The abilities of fourteen melatonin analogues to compete for [ $^3H$ ]-melatonin binding to  $mt_1$  and  $MT_2$  receptors were determined (Table 1; Figure 3A,B). All Hill coefficients were not significantly different from unity, with the exception of 2-

iodomelatonin at  $mt_1$  receptors ( $1.24 \pm 0.07$ ,  $P < 0.05$ ). Most melatonin receptor agonists bound with similar affinities to both receptor subtypes (linear regression analysis yielded a correlation coefficient of 0.87 and a slope of  $0.71 \pm 0.12$ ). Greatest selectivity was exhibited by 2-iodomelatonin, which had 5 fold higher affinity for  $mt_1$  receptors, and 6-chloromelatonin, which had 5 fold higher affinity for  $MT_2$  receptors. A number of putative partial and silent receptor antagonists had substantially higher affinities for  $MT_2$  than  $mt_1$  receptors. Greatest selectivity was demonstrated by 4-P-PDOT, which had a 61 fold higher affinity for  $MT_2$  compared to  $mt_1$  receptors.

#### Inhibition of forskolin-stimulated cyclic AMP in CHO- $mt_1$ and CHO- $MT_2$ cells by melatonin

Melatonin produced a potent, monophasic and concentration-dependent inhibition of forskolin-stimulated cyclic AMP in both CHO- $mt_1$  and CHO- $MT_2$  cells, with  $pEC_{50}$  values of



**Figure 2** [ $^3H$ ]-melatonin saturation binding to human recombinant  $mt_1$  (A) and  $MT_2$  ( $MT_2$ ) receptors. Data are mean of duplicate points and representative of three separate experiments.

**Table 1** Affinities of melatonin analogues to compete for [ $^3H$ ]-melatonin and 2-[ $^{125}I$ ]-iodomelatonin binding to human recombinant  $mt_1$  and  $MT_2$  receptors

Compound	$mt_1$ $pK_i$		$MT_2$ $pK_i$		$mt_1K_i/MT_2K_i$
	2-[ $^{125}I$ ]-iodomelatonin	[ $^3H$ ]-melatonin	[ $^3H$ ]-melatonin		
2-Iodomelatonin	$10.57 \pm 0.09$	$10.55 \pm 0.11$	$9.87 \pm 0.10$		0.2
S20098	$9.82 \pm 0.09$	$10.09 \pm 0.09$	$10.23 \pm 0.09$		1.4
Melatonin	$9.67 \pm 0.06$	$9.93 \pm 0.06$	$9.65 \pm 0.05$		0.5
GR196429	$9.39 \pm 0.11$	$9.69 \pm 0.09$	$9.66 \pm 0.07$		0.9
6-Hydroxymelatonin	$8.94 \pm 0.07$	$9.23 \pm 0.05$	$8.79 \pm 0.05$		0.4
6-Chloromelatonin	$8.90 \pm 0.07$	$9.10 \pm 0.05$	$9.77 \pm 0.05$		4.8
8-M-PDOT	ND	$8.23 \pm 0.07$	$8.95 \pm 0.02$		5.2
5-methoxyluzindole	ND	$7.97 \pm 0.03$	$9.20 \pm 0.05$		16.9
GR128107	ND	$7.71 \pm 0.08$	$9.07 \pm 0.01$		22.8
N-acetyl-5-HT	$7.13 \pm 0.35$	$7.08 \pm 0.04$	$7.00 \pm 0.14$		0.8
Luzindole	$6.21 \pm 0.20$	$6.61 \pm 0.04$	$7.63 \pm 0.08$		10.7
4-P-PDOT	ND	$6.59 \pm 0.04$	$8.37 \pm 0.19$		60.7
5-MCA-NAT	ND	$5.85 \pm 0.05$	$5.98 \pm 0.05$		1.4
5-HT	ND	$< 5.6$	$< 5.6$		

Values are mean  $pK_i \pm$  s.e. mean from 3–12 separate experiments. All slope values were not significantly different from unity, with the exception of 2-iodomelatonin at  $mt_1$  receptors ( $1.24 \pm 0.07$ ,  $P < 0.05$ ). ND = not determined.

$9.53 \pm 0.16$  and  $9.74 \pm 0.05$  and maximum inhibitions of  $83 \pm 4$  and  $64 \pm 3\%$ , respectively (Table 2; Figure 4A,B). Melatonin had no effect on basal levels of cyclic AMP in control or pertussis toxin treated cells, or on forskolin-stimulated cyclic AMP levels in mock transfected CHO cells (data not shown). Overnight incubation with pertussis toxin ( $100 \text{ ng ml}^{-1}$ ) completely abolished responses to melatonin in both CHO- $mt_1$  and CHO- $MT_2$  cells (Figure 4A,B). In CHO- $mt_1$  cells, pertussis toxin incubation revealed a small stimulatory response to melatonin in the presence of forskolin, with a  $pEC_{50}$  value of  $8.37 \pm 0.17$  and maximum response of  $26 \pm 3\%$  ( $0.1 \mu\text{M}$ ) of forskolin-stimulated cyclic AMP (Figure 4A). This stimulatory response of melatonin was not observed in pertussis toxin-treated CHO- $MT_2$  cells (Figure 4B).


#### Pharmacological characterization of functional responses in CHO- $mt_1$ and CHO- $MT_2$ cells

The inhibitory effect of melatonin was mimicked by a range of indolic and non-indolic melatonin analogues (Figure 5A,B; Table 2). At the  $mt_1$  receptor, the rank order of potency was 2-iodomelatonin > melatonin = S20098 > 6-chloromelatonin = GR196429 > 6-hydroxymelatonin > N-acetyl-5-HT, whilst at

the  $MT_2$  receptor the rank order was S20098 = 2-iodomelatonin > melatonin = 6-chloromelatonin > GR196429 > 6-hydroxymelatonin > N-acetyl-5-HT. At both  $mt_1$  and  $MT_2$  receptors all compounds tested were full agonists with respect to melatonin, with the exception of 6-hydroxymelatonin at the  $MT_2$  receptor, which was a partial agonist with an intrinsic activity of  $0.82 \pm 0.05$  ( $P < 0.05$  vs melatonin,  $n = 4$ ). 2-iodomelatonin showed the greatest degree of  $mt_1$  selectivity relative to melatonin, being 30 fold more potent than melatonin to inhibit forskolin-stimulated cyclic AMP in CHO- $mt_1$  cells, whilst only 2 fold more potent than melatonin in CHO- $MT_2$  cells. 6-chloromelatonin showed the greatest degree of  $MT_2$  receptor selectivity relative to melatonin. In CHO- $mt_1$  cells, 6-chloromelatonin was 25 fold less potent than melatonin, whilst in CHO- $MT_2$  cells it was equipotent with melatonin. All other agonists had similar potency ratios, with respect to melatonin, at both  $mt_1$  and  $MT_2$  receptors.

#### Antagonist studies in CHO- $mt_1$ and CHO- $MT_2$ cells

The putative melatonin receptor antagonist luzindole ( $0.1$ – $100 \mu\text{M}$ ) produced a concentration-dependent rightward shift




**Figure 3** Competition for [ $^3\text{H}$ ]-melatonin binding at human recombinant  $mt_1$  (A) and  $MT_2$  (B) receptors. Data are mean of duplicate points and representative of at least three separate experiments. Binding in the absence of competitor (mean  $\pm$  s.e.mean) =  $0.42 \pm 0.02 \text{ pmol mg}^{-1}$  ( $mt_1$ ) and  $0.35 \pm 0.02 \text{ pmol mg}^{-1}$  ( $MT_2$ ).


**Table 2** Potencies of melatonin receptor agonists to inhibit forskolin stimulated cyclic AMP accumulation in CHO- $mt_1$  and CHO- $MT_2$  cells

Compound	$pEC_{50}$	$mt_1$ $E_{max}$	PR	$pEC_{50}$	$MT_2$ PR	$E_{max}$	PR $mt_1/PR MT_2$
2-Iodomelatonin	$11.0 \pm 0.1$	$82 \pm 3$	0.03	$10.1 \pm 0.2$	0.50	$76 \pm 3$	0.07
Melatonin	$9.5 \pm 0.2$	$81 \pm 4$	1.00	$9.7 \pm 0.1$	1.00	$65 \pm 3$	1.0
S20098	$9.4 \pm 0.2$	$82 \pm 4$	1.42	$10.3 \pm 0.1$	0.30	$72 \pm 2$	4.7
6-Chloromelatonin	$8.1 \pm 0.1$	$85 \pm 4$	24.6	$9.8 \pm 0.2$	1.02	$60 \pm 9$	24.0
GR196429	$8.0 \pm 0.1$	$87 \pm 2$	36	$9.1 \pm 0.1$	4.61	$68 \pm 3$	7.79
6-Hydroxymelatonin	$7.2 \pm 0.1$	$76 \pm 3$	205	$7.4 \pm 0.1$	231	$50 \pm 5^*$	0.88
N-Acetyl-5-HT	$5.7 \pm 0.1$	$84 \pm 5$	7150	$6.6 \pm 0.1$	1170	$65 \pm 2$	6.1

Values are mean  $pEC_{50} \pm$  s.e.mean of 3–11 separate experiments. PR, potency ratio with respect to melatonin;  $E_{max}$ , maximum response to agonist expressed as percentage inhibition of forskolin stimulated cyclic AMP; \*significantly different from melatonin ( $P < 0.05$ ,  $n = 4$ ).



**Figure 4** The effect of pertussis toxin ( $100 \text{ ng ml}^{-1}$ ) on melatonin-induced inhibition of forskolin-stimulated cyclic AMP production in CHO- $mt_1$  (A) and CHO- $MT_2$  (B) cells. Data are mean  $\pm$  s.e. mean of three separate experiments. Mean basal cyclic AMP levels were  $0.30 \text{ pmol well}^{-1}$  (range  $0.18\text{--}0.35$ ); forskolin-stimulated cyclic AMP levels were  $27 \text{ pmol well}^{-1}$  (range  $14\text{--}40$ ). Similar values were obtained in subsequent experiments.



**Figure 5** Agonist concentration-effect curves for inhibition of forskolin-stimulated cyclic AMP in CHO- $mt_1$  (A) and CHO- $MT_2$  cells. The data are mean of 3–11 separate experiments.

of the melatonin concentration-effect curves in both CHO- $mt_1$  and CHO- $MT_2$  cells (Figure 6A,B). Non-linear regression analyses yielded  $pA_2$  values of  $5.75 \pm 0.10$  and  $7.64 \pm 0.11$  for  $mt_1$  and  $MT_2$  respectively ( $n = 3\text{--}4$ ). Schild slope parameters for luzindole were  $1.00 \pm 0.10$  and  $0.94 \pm 0.07$  for  $mt_1$  and  $MT_2$  respectively, which were not significantly different to unity ( $P > 0.05$ ). In CHO- $mt_1$  cells luzindole had no significant effect ( $P > 0.05$ ) on Hill slopes or maximum responses to melatonin (data not shown). However, luzindole caused a concentration-dependent enhancement of the lower asymptote of the melatonin concentration-effect curve of  $95 \pm 20\%$  ( $P < 0.05$ )


at  $30 \mu\text{M}$ , and  $134 \pm 36\%$  ( $P < 0.01$ ) at  $100 \mu\text{M}$ , of forskolin-stimulated cyclic AMP (Figure 6A). Luzindole alone produced a small and variable enhancement of forskolin-stimulated cyclic AMP, which was not of the same magnitude as that seen in the presence of melatonin (Figure 6A and 7A). In contrast, in CHO- $MT_2$  cells, luzindole was without effect on forskolin-stimulated cyclic AMP, and had no effect on the maximum response to melatonin (Figure 6B and 7B). However, luzindole reduced the Hill slopes for the melatonin concentration-effect curves in a concentration-dependent manner (e.g.  $n_H = 1.65 \pm 0.33$  and  $0.89 \pm 0.19$ , control and  $10 \mu\text{M}$  luzindole

respectively;  $P < 0.05$ ). For these reasons,  $pA_2$  rather than  $pK_B$  values have been quoted.


In CHO- $mt_1$  cells the amidotetraline 4-P-PDOT (10  $\mu M$ ) had no effect on forskolin-stimulated cyclic AMP levels, either alone (Figure 7A), or in the presence of melatonin (data not shown). In contrast, in CHO- $MT_2$  cells, 4-P-PDOT was an agonist, producing a concentration-dependent inhibition of forskolin stimulated cyclic AMP, with a  $pEC_{50}$  value of  $8.72 \pm 0.29$  and intrinsic activity of  $0.86 \pm 0.15$  ( $n = 3$ ; Figure 7B). This degree of agonism precluded the determination of a  $pA_2$  value for 4-P-PDOT at  $MT_2$  receptors.

## Discussion

This study has described the pharmacology of human recombinant melatonin  $mt_1$  and  $MT_2$  receptors stably expressed in CHO cells, using a number of indolic and non-indolic analogues of melatonin. Human recombinant  $mt_1$  and  $MT_2$  receptors were expressed in identical cell lines using the same expression vectors and achieved similar receptor expression levels. This has allowed the pharmacology of each receptor to be compared in systems with identical cellular backgrounds.



**Figure 6** Antagonism by luzindole of melatonin-induced inhibition of forskolin-stimulated cyclic AMP production in CHO- $mt_1$  (A) and CHO- $MT_2$  (B) cells. Data are mean  $\pm$  s.e. mean of 3–4 separate experiments. Inset: Schild plots depicting antagonism of melatonin responses by luzindole. The line through the data was plotted using the parameters obtained from the analysis described in the Methods.



**Figure 7** The effect of melatonin, luzindole and 4-P-PDOT on forskolin-stimulated cyclic AMP production in CHO- $mt_1$  (A) and CHO- $MT_2$  (B) cells. Data are mean  $\pm$  s.e. mean of 3–4 separate experiments.

The affinities of melatonin analogues at the  $mt_1$  receptor measured using [ $^3H$ ]-melatonin are in good agreement with affinities obtained using 2-[ $^{125}I$ ]-iodomelatonin. This is the first report of the use of [ $^3H$ ]-melatonin for the pharmacological characterization of melatonin receptor subtypes and validates the use of this radioligand as an alternative to 2-[ $^{125}I$ ]-iodomelatonin. Similarly, Kennaway *et al.* (1994) showed comparable pharmacological profiles for the binding of these two radioligands to chicken brain. In addition to its safety benefits, [ $^3H$ ]-melatonin is chemically identical to the endogenous hormone and is therefore the ligand of choice for the generation of structure-activity relationships. Indeed, Kennaway *et al.* (1994) concluded that 2-[ $^{125}I$ ]-iodomelatonin may not be an appropriate ligand for melatonin binding studies because of its disproportionately slow dissociation rate.

Studies of association binding kinetics at  $mt_1$  receptors showed that [ $^3H$ ]-melatonin associated rapidly and reached equilibrium after 30 min. At the  $MT_2$  receptor, [ $^3H$ ]-melatonin associated more slowly and reached equilibrium after approximately 60 min. Interestingly, [ $^3H$ ]-melatonin dissociated from the  $MT_2$  receptor over a much longer period of time, and failed to reach non-specific binding levels, with 22% of specifically bound radioligand still present after 240 min. Our data suggests that, over the time frame of the experiment, a component of the [ $^3H$ ]-melatonin binding to the  $MT_2$  receptor may be irreversible. This result may be explained using a model of multiple receptor states. Long term exposure of receptor with radioligand may 'lock' a proportion of occupied receptor into a super-high affinity state which may appear irreversible over the time frame of the experiment. This behaviour of slowly dissociating radioligands has been reported for a number of 7-transmembrane, G-protein coupled receptors and has been shown to be sensitive to the presence of guanine nucleotides, suggesting the presence of high affinity, agonist occupied, G-protein linked receptors (Cohen *et al.*, 1996). Interestingly, whilst the  $pK_D$  of 9.78 for [ $^3H$ ]-melatonin at the  $mt_1$  receptor determined from kinetic studies was in good agreement with the value of 9.89 estimated from equilibrium saturation binding, at the  $MT_2$  receptor the  $pK_D$  of 9.92 determined from kinetic studies was nearly half a log unit higher than the  $pK_D$  of 9.56 estimated from equilibrium saturation binding. The irreversible component of the binding of [ $^3H$ ]-melatonin to the  $MT_2$  receptor may explain this discrepancy.

The affinities of a range of melatonin analogues at  $mt_1$  and  $MT_2$  receptors are in broad agreement with those obtained by other groups (Dubocovich *et al.*, 1997; Reppert *et al.*, 1995). Most ligands were either non-selective or showed some degree of  $MT_2$  selectivity. Ligands which exhibited the greatest degree of selectivity were 2-iodomelatonin, having 5 fold higher affinity at the  $mt_1$  receptor, and the amidotetraline derivative, 4-P-PDOT, with 60 fold higher affinity for the  $MT_2$  receptor. Similarly, Dubocovich and co-workers (1997) have reported that 4-P-PDOT had 300 fold higher affinity at the  $MT_2$  receptor.

Functional studies demonstrated that melatonin receptor activation resulted in the inhibition of forskolin-stimulated cyclic AMP production. The abrogation of the response to melatonin following pre-treatment with pertussis toxin indicates that this response is mediated *via*  $G_i/G_o$  G-proteins, confirming previous reports using both recombinant human receptor systems (Reppert *et al.*, 1994; 1995) and sheep pars tuberalis (Morgan *et al.*, 1990). In CHO- $mt_1$  cells, pertussis toxin treatment revealed a small enhancement by melatonin of forskolin-stimulated cyclic AMP, suggesting that the  $mt_1$  receptor can promiscuously couple to G-proteins other than

$G_i/G_o$ , presumably  $G_s$ . Promiscuous coupling of melatonin receptors has been previously reported for cloned *Xenopus* receptors expressed in human embryonic kidney 293 cells (Yung *et al.*, 1995) and endogenous receptors in sheep pars tuberalis (Morgan *et al.*, 1990; 1995). Interestingly, the stimulatory response to melatonin following pertussis toxin treatment was not observed in CHO- $MT_2$  cells, indicating a difference in the second-messenger coupling characteristics of these receptors.

In both CHO- $mt_1$  and CHO- $MT_2$  cells, luzindole antagonised melatonin responses in a concentration-dependent manner, with approximately 80 fold higher affinity at the  $MT_2$  receptor. This is the first direct comparison of the affinity of luzindole for  $mt_1$  and  $MT_2$  receptors measured in a functional assay and confirms previous reports that luzindole has selectivity for  $MT_2$  receptors (Dubocovich *et al.*, 1997). However, the  $pA_2$  value of 5.75 for luzindole at the  $mt_1$  receptor is somewhat lower than the values of 7.2 and 7.3 determined by Beresford and co-workers (1998b) and Witt-Enderby & Dubocovich (1996) respectively. It is not clear the extent to which the  $pA_2$  value for luzindole at the  $mt_1$  receptor determined in this study is compromised by its complex action in this system. In CHO- $mt_1$ , but not in CHO- $MT_2$  cells, luzindole alone produced a small increase in forskolin-stimulated cyclic AMP, suggesting that luzindole may act as an inverse agonist at the  $mt_1$  receptor. In support of this hypothesis, Dubocovich & Masana (1998) have shown that the affinity of luzindole for the  $mt_1$  receptor increased in the presence of GTP, a property which is displayed by inverse agonists. Interestingly, in the presence of low concentrations of melatonin, luzindole enhanced forskolin-stimulated cyclic AMP, indicating that the inverse agonist response can be augmented in the presence of melatonin. Alternatively, as our experiments with pertussis toxin suggest that the  $mt_1$  receptor can promiscuously couple to non  $G_i/G_o$  G-proteins, luzindole may potentiate the signalling of melatonin through this alternative G-protein pathway. Further studies are clearly required to elucidate the exact nature of the effects of luzindole and its interaction with melatonin at the  $mt_1$  receptor.

In CHO- $mt_1$  cells, the putative  $MT_2$  receptor selective antagonist 4-P-PDOT, at concentrations up to 10  $\mu M$ , had no effect on forskolin-stimulated cyclic AMP levels alone, and did not affect melatonin concentration-effect curves. In contrast, in CHO  $MT_2$  cells, 4-P-PDOT was a partial agonist with respect to melatonin, producing a concentration-dependent inhibition of forskolin-stimulated cyclic AMP accumulation with a  $pEC_{50}$  of 8.7 and intrinsic activity of 0.87. The intrinsic activity of 4-P-PDOT at the  $MT_2$  receptor precluded the determination of a  $pA_2$  value for this compound. Whilst this result supports the claim that 4-P-PDOT is a  $MT_2$  receptor selective compound (Dubocovich *et al.*, 1997), 4-P-PDOT did not behave as a neutral antagonist in our assay. In agreement, Nonno and co-workers (1999) recently reported that 4-P-PDOT was a partial agonist (intrinsic activity of 0.37) at the  $MT_2$  receptor in a [ $^{35}S$ ]-GTP $\gamma$ S binding assay, a functional measure of receptor activation. In contrast, 4-P-PDOT has previously been shown to be a silent, simple competitive antagonist with high affinity at the presynaptic  $MT_2$  receptor mediating the inhibition of electrically-evoked dopamine release in rabbit retina (Dubocovich *et al.*, 1997). The lack of agonism in the rabbit retina may be explained by the lower level of receptor expression in this tissue (8.5 fmol  $mg^{-1}$  protein; Dubocovich, 1995), compared with 0.2 pmol  $mg^{-1}$  protein in our CHO- $MT_2$  cells and 0.5 pmol  $mg^{-1}$  in the NIH3T3 cells used by Nonno *et al.* (1999). It is well established that a reduction in receptor number may reduce the sensitivity of a tissue to low efficacy



agonists (Furchgott, 1966). Therefore, the rabbit retina may not reveal the agonist activity of low efficacy compounds. Interestingly, it has recently been demonstrated that 4-P-PDOT is a silent antagonist of melatonin responses in an animal model of circadian rhythms (Dubocovich *et al.*, 1998b).

While antagonist affinity estimates provide a theoretically robust method of classifying and characterizing G-protein coupled receptors, functional data from agonist experiments can also be used to characterize receptors (Leff, 1987). Estimates of agonist potency alone are of little value in receptor classification, as agonist potency is dependent on both receptor density and transduction efficiency (Kenakin & Beek, 1980). However, the potency of an agonist, when expressed relative to the potency of a reference agonist, provides a robust parameter for receptor classification, since this value will be tissue-independent for full agonists. Therefore, agonists that yield different potency ratios with respect to melatonin can be used to discriminate between  $mt_1$  and  $MT_2$  receptors. 2-iodomelatonin was 30 fold more potent than melatonin in CHO- $mt_1$  cells, but had a similar potency to melatonin in CHO- $MT_2$  cells. Conversely, 6-chloromelatonin was 25 fold less potent than melatonin in CHO- $mt_1$  cells, whilst being equipotent with melatonin in CHO- $MT_2$  cells. At  $MT_2$

receptors, melatonin, 2-iodomelatonin and 6-chloromelatonin are essentially equipotent, whilst at the  $mt_1$  receptor these agonists have the rank order of potency of 2-iodomelatonin > melatonin > 6-chloromelatonin. Thus, in addition to antagonist data, these three agonists may be of use in classifying melatonin receptors.

In conclusion, we have expressed human recombinant melatonin receptors in CHO cells using identical expression systems and have characterized these receptors using [ $^3$ H]-melatonin binding assays and measurement of forskolin stimulated cyclic AMP accumulation. Luzindole is an antagonist at melatonin receptors, with higher affinity at  $MT_2$  receptors. In addition to the use of antagonists, the rank order of potency of melatonin, 2-iodomelatonin, and 6-chloromelatonin can be used to classify melatonin receptors.

We thank Dr Stephen Foord and Jason Brown for providing  $mt_1$  and  $MT_2$  receptor cDNAs, Kerri Cartwright for expert technical assistance and Dr David Hall for advice during the preparation of the manuscript.

## References

- ARENDETT, J. (1995). Melatonin and the mammalian pineal gland. London: Chapman and Hall.
- BERESFORD, I.J.M., BROWNING, C., STARKEY, S.J., BROWN, J., FOORD, S.M., COUGHLAN, J., NORTH, P.C., DUBOCOVICH, M.L. & HAGAN, R.M. (1998a). GR196429: A non-indolic agonist at high-affinity melatonin receptors. *J. Pharmacol. Exp. Therap.*, **285**, 1239–1245.
- BERESFORD, I.J.M., HARVEY, F.J., HALL, D.A. & GILES, H. (1998b). Pharmacological characterisation of melatonin  $mt_1$  receptor mediated stimulation of [ $^{35}$ S]-GTP $\gamma$ S binding. *Biochem. Pharmacol.*, **56**, 1167–1174.
- BLACK, J.W., LEFF, P. & SHANKLEY, N.P. (1985). Further analysis of anomalous  $pK_B$  values for histamine  $H_2$  receptor antagonists on the mouse isolated stomach assay. *Br. J. Pharmacol.*, **86**, 581–587.
- BROWNING, C., BERESFORD, I.J.M. & GILES, H. (1998). Differential pharmacological characterisation of human recombinant melatonin  $mt_1$  and  $MT_2$  receptors. *Br. J. Pharmacol.*, **124**, 34P.
- BROWNING, C., BROWN, J.D., BERESFORD, I.J.M. & GILES, H. (1997). Pharmacological characterisation of [ $^3$ H]-melatonin binding to human melatonin  $Mel_{1A}$  and  $Mel_{1B}$  receptors. *Br. J. Pharmacol.*, **120**, 361P.
- CHENG, Y. & PRUSOFF, W.H. (1973). Relationship between the inhibition constant ( $k_i$ ) and the concentration of inhibitor which causes 50 per cent inhibition ( $IC_{50}$ ) of an enzymatic reaction. *Biochem. Pharmacol.*, **22**, 3099–3108.
- COHEN, F.R., LAZARENO, S. & BIRDSALL, N.J.M. (1996). The effects of saponin on the binding and functional properties of the human adenosine  $A_1$  receptor. *Br. J. Pharmacol.*, **117**, 1521–1529.
- DUBOCOVICH, M.L. (1988). Luzindole (N-0744): A novel melatonin receptor antagonist. *J. Pharmacol. Exp. Therap.*, **246**, 902–910.
- DUBOCOVICH, M.L. (1995). Melatonin receptors: Are there multiple subtypes? *Trends Pharmacol. Sci.*, **16**, 50–56.
- DUBOCOVICH, M.L. & MASANA, M.I. (1998). The efficacy of melatonin receptor analogues is dependent on the level of human melatonin receptor subtype expression. In: *Biological Clocks, Mechanisms and Applications*. ed. Touitou, Y. pp. 289–293. Amsterdam: Elsevier Science B.V.
- DUBOCOVICH, M.L., CARDINALI, D.P., GUARDIOLA-LEMAITRE, B., HAGAN, R.M., KRAUSE, D.N., SUGDEN, D., VANHOUTTE, P.M. & YOCCA, F.D. (1998a). Melatonin receptors. In: *The IUPHAR Compendium of Receptor Characterisation and Classification*. pp. 187–193. London: IUPHAR Media.
- DUBOCOVICH, M.L., MASANA, M.I. & IACOB, S. (1997). Melatonin receptor antagonists that differentiate between the human  $Mel_{1A}$  and  $Mel_{1B}$  recombinant subtypes are used to assess the pharmacological profile of the rabbit retina  $ML1$  presynaptic heteroreceptor. *Naunyn-Schmied. Arch. Pharmacol.*, **355**, 365–375.
- DUBOCOVICH, M.L., YUN, K., ALGHOUL, W.M., BENLOUCIF, S. & MASANA, M.I. (1998b). Selective  $MT_2$  receptor antagonists block melatonin mediated phase advances of circadian rhythms. *FASEB J.*, **12**, 1211–1220.
- EBISAWA, T., KARNE, S., LERNER, M.R. & REPPERT, S.M. (1994). Expression cloning of a high-affinity melatonin receptor from *Xenopus* dermal melanophores. *Proc. Natl. Acad. Sci. (U.S.A.)*, **91**, 6133–6137.
- FURCHGOTT, R.F. (1966). The use of  $\beta$ -haloalkylamines in the differentiation of receptors and in the determination of dissociation constants of receptor-agonist complexes. *Adv. Drug Res.*, **3**, 21–55.
- KENAKIN, T.P. & BEEK, D. (1980). Is prenalterol (H133/80) really a selective beta 1 adrenoceptor agonist? Tissue selectivity resulting from differences in stimulus-response relationships. *J. Pharmacol. Exp. Therap.*, **213**, 406–413.
- KENNAWAY, D.J., HUGEL, H.M. & ROWE, S.A. (1994). Characterisation of the chicken brain melatonin-binding protein using iodinated and tritiated ligands. *J. Pineal Res.*, **17**, 137–148.
- LEFF, P. (1987). Can operational models of agonism provide a framework for classifying hormone receptors? In: *Perspectives on Receptor Classification*. ed. Black, J.W., Jenkinson, D.H. & Gerskowitch, V.P. pp. 157–168. New York: Alan R. Liss Inc.
- LEW, M.J. & ANGUS, J.A. (1995). Analysis of competitive agonist-antagonist interactions by non-linear regression. *Trends Pharmacol. Sci.*, **16**, 328–337.
- MORGAN, P.J., BARRETT, P., HAZLERIGG, D., MILLIGAN, G., LAWSON, W., MACLEAN, A. & DAVIDSON, G. (1995). Melatonin receptors couple through a cholera toxin-sensitive mechanism to inhibit cyclic AMP in the ovine pituitary. *J. Neuroendocrinol.*, **7**, 361–369.
- MORGAN, P.J., BARRETT, P., HOWELL, H.E. & HELLIWELL, R. (1994). Melatonin receptors: localisation, molecular pharmacology and physiological significance. *Neurochem. Int.*, **24**, 101–146.

- MORGAN, P.J., DAVIDSON, G., LAWSON, W. & BARRETT, P. (1990). Both pertussis toxin-sensitive and insensitive G-proteins link melatonin receptors to inhibition of adenylate cyclase in the ovine pars tuberalis. *J. Neuroendocrinol.*, **2**, 773–776.
- NONNO, R., PANNACCI, M., LUCINI, V., ANGELONI, D., FRASCHINI, F. & STANKOV, B.M. (1999). Ligand efficacy and potency at recombinant human  $MT_2$  melatonin receptors: evidence for agonist activity of some  $mt_1$ -antagonists. *Br. J. Pharmacol.*, **127**, 1288–1294.
- REPPERT, S.M., GODSON, C., MAHLE, C.D., WEAVER, D.R., SLAUGENHAUPT, S.A. & GUSELLA, J.F. (1995). Molecular characterisation of a second melatonin receptor expressed in human retina and brain: The  $Mel_{1b}$  melatonin receptor. *Proc. Natl. Acad. Sci. (U.S.A.)*, **92**, 8734–8738.
- REPPERT, S.M., WEAVER, D.R. & EBISAWA, T. (1994). Cloning and characterisation of a mammalian melatonin receptor that mediates reproductive and circadian responses. *Neuron*, **13**, 1177–1185.
- WEAVER, D.R. & REPPERT, S.M. (1996). The  $Mel_{1a}$  melatonin receptor gene is expressed in human suprachiasmatic nuclei. *Neuroreport*, **8**, 109–112.
- WITT-ENDERBY, P.A. & DUBOCOVICH, M.T. (1996). Characterisation and regulation of the human  $ML_{1A}$  melatonin receptor stably expressed in Chinese hamster ovary cells. *Mol. Pharmacol.*, **50**, 166–174.
- YOUS, Y., ANDRIEUX, J., HOWELL, M.E., MORGAN, P.J., RENARD, P., PFEIFFER, B., LESIEUR, D. & GUARDIOLA-LEMAITRE, B. (1992). Novel naphthalenic ligands with high affinity for the melatonin receptor. *J. Med. Chem.*, **35**, 1484–1486.
- YUNG, L.Y., TSIM, S. & WONG, Y.H. (1995). Stimulation of cAMP accumulation by the cloned *Xenopus* melatonin receptor through  $G_i$  and  $G_z$  proteins. *FEBS Letts.*, **372**, 99–102.

(Received August 11, 1999

Revised October 18, 1999

Accepted December 2, 1999)