

OSTROWSKI TYPE INEQUALITIES FOR FUNCTIONS WHOSE MODULUS OF THE DERIVATIVES ARE CONVEX AND APPLICATIONS

N.S. BARNETT, P. CERONE, S.S. DRAGOMIR, M.R. PINHEIRO, AND A. SOFO

ABSTRACT. Some inequalities of the Ostrowski type for functions whose modulus of derivatives are convex and applications for special means and to the f and HH -divergences in Information Theory are given.

1. INTRODUCTION

The following Ostrowski type inequalities for absolutely continuous functions are known (see [2], [3] and [4]).

Theorem 1. *Let $f : [a, b] \rightarrow \mathbb{R}$ be absolutely continuous on $[a, b]$. Then for all $x \in [a, b]$ we have*

$$(1.1) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \begin{cases} \left[\frac{1}{4} + \left(\frac{x - \frac{a+b}{2}}{b-a} \right)^2 \right] (b-a) \|f'\|_\infty & \text{if } f' \in L_\infty[a, b]; \\ \frac{1}{(p+1)^{\frac{1}{p}}} \left[\left(\frac{x-a}{b-a} \right)^{p+1} + \left(\frac{b-x}{b-a} \right)^{p+1} \right] (b-a)^{\frac{1}{p}} \|f'\|_q & \text{if } f' \in L_q[a, b], p > 1, \frac{1}{p} + \frac{1}{q} = 1; \\ \left[\frac{1}{2} + \left| \frac{x - \frac{a+b}{2}}{b-a} \right| \right] \|f'\|_1 & \end{cases}$$

where $\|\cdot\|_r$ ($r \in [1, \infty]$) are the usual Lebesgue norms on $L_r[a, b]$, i.e.,

$$\|g\|_\infty := \operatorname{ess\,sup}_{t \in [a, b]} |g(t)|.$$

The constants $\frac{1}{4}$, $\frac{1}{(p+1)^{\frac{1}{p}}}$ and $\frac{1}{2}$ are sharp in the sense that they cannot be replaced by smaller constants.

The above inequalities may also be obtained from Fink's result in [5] on choosing $n = 1$ and performing some appropriate computations.

Date: January 21, 2001.

1991 Mathematics Subject Classification. 26D15, 94A05.

Key words and phrases. Ostrowski's Inequality, Special Means, Divergence Measures in Information Theory.

The main aim of this paper is to point out some similar results in the case when the modulus of the derivative f' is a convex function on (a, b) . Applications for special means and f and HH -divergence in Information Theory are also provided.

2. THE RESULTS

We start with the following lemma which is of intrinsic interest (see also [1]).

Lemma 1. *Let $f : [a, b] \rightarrow \mathbb{R}$ be an absolutely continuous function on $[a, b]$, then, for any $x \in [a, b]$,*

$$(2.1) \quad f(x) = \frac{1}{b-a} \int_a^b f(t) dt + \frac{1}{b-a} \int_a^b (x-t) \left[\int_0^1 f'[(1-\lambda)x + \lambda t] d\lambda \right] dt$$

Proof. For any $x, t \in [a, b]$, $x \neq t$, one has

$$\frac{f(x) - f(t)}{x-t} = \frac{1}{x-t} \int_t^x f'(u) du = \int_0^1 f'[(1-\lambda)x + \lambda t] d\lambda$$

showing that

$$(2.2) \quad f(x) = f(t) + (x-t) \int_0^1 f'[(1-\lambda)x + \lambda t] d\lambda \text{ for any } x, t \in [a, b].$$

Integrating (2.2) over t on $[a, b]$ and dividing the result by $(b-a)$, gives the desired identity (2.1). ■

Using the above lemma the following result can be pointed out improving Ostrowski's inequality.

Theorem 2. *Let $f : [a, b] \rightarrow \mathbb{R}$ be an absolutely continuous function on $[a, b]$ so that $|f'|$ is convex on (a, b) . If $f' \in L_\infty[a, b]$, then for any $x \in [a, b]$,*

$$(2.3) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{2} \left[\frac{1}{4} + \left(\frac{x - \frac{a+b}{2}}{b-a} \right)^2 \right] (b-a) [|f'(x)| + \|f'\|_\infty].$$

The constant $\frac{1}{2}$ is sharp in the sense that it cannot be replaced by a smaller constant.

Proof. Using (2.1) and taking the modulus, we have

$$\begin{aligned} \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| &= \frac{1}{b-a} \left| \int_a^b \int_0^1 (x-t) f'[(1-\lambda)x + \lambda t] d\lambda dt \right| \\ &\leq \frac{1}{b-a} \int_a^b \int_0^1 |x-t| |f'[(1-\lambda)x + \lambda t]| d\lambda dt \end{aligned}$$

$$\begin{aligned}
&\leq \frac{1}{b-a} \int_a^b \int_0^1 |x-t| [(1-\lambda)|f'(x)| + \lambda|f'(t)|] d\lambda dt \\
&\quad (\text{by convexity of } |f'|) \\
&= \frac{1}{b-a} \int_a^b |x-t| \left[|f'(x)| \int_0^1 (1-\lambda) d\lambda + |f'(t)| \int_0^1 \lambda d\lambda \right] dt \\
&= \frac{1}{b-a} \int_a^b |x-t| \left[\frac{|f'(x)| + |f'(t)|}{2} \right] dt := M(x) \\
&\leq \frac{1}{2} \frac{1}{b-a} \operatorname{ess. sup}_{t \in [a,b]} [|f'(x)| + |f'(t)|] \int_a^b |x-t| dt \\
&= \frac{1}{2} \left[\frac{(x-a)^2 + (b-x)^2}{2(b-a)} \right] [|f'(x)| + \|f'\|_\infty] \\
&= \frac{1}{2} \left[\frac{1}{4} + \left(\frac{x - \frac{a+b}{2}}{b-a} \right)^2 \right] (b-a) [|f'(x)| + \|f'\|_\infty],
\end{aligned}$$

and the inequality (2.3) is proved.

Assume that (2.3) holds with a constant $C > 0$, that is,

$$\begin{aligned}
(2.4) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\
\leq C \left[\frac{1}{4} + \left(\frac{x - \frac{a+b}{2}}{b-a} \right)^2 \right] (b-a) [|f'(x)| + \|f'\|_\infty]
\end{aligned}$$

for any $x \in [a, b]$ with f as in the hypothesis of the theorem.

Consider the function

$$f_0 : [a, b] \rightarrow \mathbb{R}, f_0(t) = k \left| t - \frac{a+b}{2} \right|, \quad k > 0, t \in [a, b].$$

Since $|f_0'(t)| = k$, for any $t \in [a, b]$ and

$$\frac{1}{b-a} \int_a^b f_0(t) dt = \frac{k}{4} (b-a), \quad \|f_0'\|_\infty = k$$

then choosing $f = f_0$ and $x = \frac{a+b}{2}$ in (2.4), we get

$$\frac{k}{4} (b-a) \leq \frac{Ck(b-a)}{2}$$

giving $C \geq \frac{1}{2}$, which proves the sharpness of the constant $\frac{1}{2}$. ■

The following particular case is interesting.

Corollary 1. *With the assumptions of Theorem 3, we have the inequality*

$$(2.5) \quad \left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{8} (b-a) \left[\left| f'\left(\frac{a+b}{2}\right) \right| + \|f'\|_\infty \right]$$

and the constant $\frac{1}{8}$ is the best possible.

The following result in terms of the p -norms also holds:

Theorem 3. *Let $f : [a, b] \rightarrow \mathbb{R}$ be as in Theorem 3. If $f' \in L_p[a, b], p > 1, \frac{1}{p} + \frac{1}{q} = 1$, then for any $x \in [a, b]$,*

$$(2.6) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{2(q+1)^{\frac{1}{q}}} \left[\left(\frac{b-x}{b-a} \right)^{q+1} + \left(\frac{x-a}{b-a} \right)^{q+1} \right]^{\frac{1}{q}} (b-a)^{\frac{1}{q}} \| |f'(x)| + |f'| \|_p$$

The constant $\frac{1}{2}$ is sharp in the sense that it cannot be replaced by a smaller constant.

Proof. According to the proof of Theorem 2, we have

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{b-a} \int_a^b |x-t| \left[\frac{|f'(x)| + |f'(t)|}{2} \right] dt := M(x).$$

Using Hölder's integral inequality for $p > 1, \frac{1}{p} + \frac{1}{q} = 1$, we get that

$$\begin{aligned} M(x) &\leq \frac{1}{2(b-a)} \left(\int_a^b |x-t|^q dt \right)^{\frac{1}{q}} \left(\int_a^b (|f'(x)| + |f'(t)|)^p dt \right)^{\frac{1}{p}} \\ &= \frac{1}{2(b-a)} \left[\frac{(b-x)^{q+1} + (x-a)^{q+1}}{q+1} \right]^{\frac{1}{q}} \| |f'(x)| + |f'| \|_p \end{aligned}$$

and the inequality (2.6) is proved.

Reconsider the function utilised in Theorem 2,

$$f_0 : [a, b] \rightarrow \mathbb{R}, f_0(t) = k \left| t - \frac{a+b}{2} \right|, k > 0, t \in [a, b]$$

which has $|f'_0(t)| (= k)$ convex in $[a, b]$. If we assume that (2.6) holds with a constant $D > 0$ instead of $\frac{1}{2}$, so that

$$\begin{aligned} \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| &\leq \frac{D}{(q+1)^{\frac{1}{q}}} \left[\left(\frac{b-x}{b-a} \right)^{q+1} + \left(\frac{x-a}{b-a} \right)^{q+1} \right]^{\frac{1}{q}} (b-a)^{\frac{1}{q}} \| |f'(x)| + |f'| \|_p, \end{aligned}$$

then taking $f = f_0$ over $x = \frac{a+b}{2}$, we get,

$$\frac{k}{4} (b-a) \leq \frac{D}{(q+1)^{\frac{1}{q}}} \left(\frac{1}{2^q} \right)^{\frac{1}{q}} (b-a)^{\frac{1}{q}} k (b-a)^{\frac{1}{p}}, q > 1, p > 1, \frac{1}{p} + \frac{1}{q} = 1$$

giving, on simplification,

$$D \geq \frac{1}{2} (q+1)^{\frac{1}{q}}, q > 1.$$

Taking the limit as $q \rightarrow \infty$ and since,

$$\lim_{q \rightarrow \infty} (q+1)^{\frac{1}{q}} = \exp \left\{ \lim_{q \rightarrow \infty} \left[\frac{\ln(1+q)}{q} \right] \right\} = \exp 0 = 1,$$

we deduce that $D \geq \frac{1}{2}$, which proves the sharpness of the constant. ■

A particular case is the following mid-point inequality:

Corollary 2. *With the assumptions of Theorem 3, we have,*

$$(2.7) \quad \left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ \leq \frac{1}{4(q-a)^{\frac{1}{q}}} (b-a)^{\frac{1}{q}} \left(\int_a^b \left[\left| f'\left(\frac{a+b}{2}\right) \right| + |f'(t)| \right]^p dt \right)^{\frac{1}{p}} \quad \left(p > 1, \frac{1}{p} + \frac{1}{q} = 1 \right)$$

The constant $\frac{1}{4}$ is sharp in the previous sense.

Finally, the case involving the 1-norm is embodied in the following theorem:

Theorem 4. *Let $f : [a, b] \rightarrow \mathbb{R}$ be as in Theorem 2. If $f' \in L_1[a, b]$, then, for any $x \in [a, b]$,*

$$(2.8) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{2} \left[\frac{1}{2} + \left| \frac{x - \frac{a+b}{2}}{b-a} \right| \right] [(b-a) |f'(x)| + \|f'\|_1].$$

Proof. We have, from the proof of Theorem 2, that

$$\begin{aligned} M(x) &\leq \sup_{t \in [a, b]} |x-t| \frac{1}{b-a} \int_a^b \left[\frac{|f'(x)| + |f'(t)|}{2} \right] dt \\ &= \frac{1}{2(b-a)} \max(x-a, b-x) \left[(b-a) |f'(x)| + \int_a^b |f'(t)| dt \right] \\ &= \frac{1}{2} \left[\frac{1}{2} + \left| \frac{x - \frac{a+b}{2}}{b-a} \right| \right] [(b-a) |f'(x)| + \|f'\|_1] \end{aligned}$$

and the inequality (2.8) is proved. ■

In particular, we have the mid-point inequality:

Corollary 3. *Assume that f is as in Theorem 4. Then*

$$(2.9) \quad \left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{4} \left[(b-a) \left| f'\left(\frac{a+b}{2}\right) \right| + \int_a^b |f'(t)| dt \right].$$

Another way to estimate the difference

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right|$$

is presented in the following theorem.

Theorem 5. *Let $f : [a, b] \rightarrow \mathbb{R}$ be an absolutely continuous function on $[a, b]$ so that $|f'|$ is convex on (a, b) . Then, for any $x \in [a, b]$,*

$$(2.10) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ \leq \frac{1}{2} \left\{ \left[\frac{1}{4} + \left(\frac{x - \frac{a+b}{2}}{b-a} \right)^2 \right] |f'(x)| (b-a) \right. \\ \left. + \frac{1}{(q+1)^{\frac{1}{q}}} \left[\left(\frac{b-x}{b-a} \right)^{q+1} + \left(\frac{x-a}{b-a} \right)^{q+1} \right]^{\frac{1}{q}} (b-a)^{\frac{1}{q}} \|f'\|_p \right\},$$

where $p > 1, \frac{1}{p} + \frac{1}{q} = 1$.

Proof. With the notation of Theorem 2, we have,

$$\begin{aligned} M(x) &= \frac{1}{2(b-a)} \left[|f'(x)| \int_a^b |x-t| dt + \int_a^b |x-t| |f'(t)| dt \right] \\ &= \frac{1}{2(b-a)} \left[|f'(x)| \frac{(x-a)^2 + (b-x)^2}{2} + \int_a^b |x-t| |f'(t)| dt \right] \\ &= \frac{1}{2} \left[|f'(x)| \left[\frac{1}{4} + \left(\frac{x - \frac{a+b}{2}}{b-a} \right)^2 \right] (b-a) + \frac{1}{b-a} \int_a^b |x-t| |f'(t)| dt \right]. \end{aligned}$$

Using Hölder's inequality,

$$\begin{aligned} &\frac{1}{b-a} \int_a^b |x-t| |f'(t)| dt \\ &\leq \frac{1}{b-a} \left(\int_a^b |x-t|^q dt \right)^{\frac{1}{q}} \left(\int_a^b |f'(t)|^p dt \right)^{\frac{1}{p}} \\ &= \frac{1}{b-a} \left[\frac{(b-x)^{q+1} + (x-a)^{q+1}}{q+1} \right]^{\frac{1}{q}} \|f'\|_p \\ &= \frac{1}{(q+1)^{\frac{1}{q}}} \left[\left(\frac{b-x}{b-a} \right)^{q+1} + \left(\frac{x-a}{b-a} \right)^{q+1} \right]^{\frac{1}{q}} (b-a)^{\frac{1}{q}} \|f'\|_p, \end{aligned}$$

and the theorem is proved. ■

The following particular corollary is of interest providing a bound for the mid-point.

Corollary 4. *Let f be as in the previous theorem. Then one has the inequality:*

$$(2.11) \quad \left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \\ \leq \frac{1}{4} \left\{ \frac{1}{2} \left| f'\left(\frac{a+b}{2}\right) \right| (b-a) + \frac{1}{(q+1)^{\frac{1}{q}}} (b-a)^{\frac{1}{q}} \|f'\|_p \right\}$$

3. APPLICATIONS FOR SPECIAL MEANS

In the applications below we consider the following definitions of some special means:

– Arithmetic mean,

$$A = A(a, b) = \frac{a+b}{2}; \quad a, b > 0.$$

– Geometric mean,

$$G = G(a, b) = \sqrt{ab}; \quad a, b > 0.$$

– Logarithmic mean,

$$L = L(a, b) = \begin{cases} \frac{b-a}{\ln b - \ln a}, & a \neq b > 0, \\ a, & a = b. \end{cases}$$

– p -Logarithmic mean,

$$L_p(a, b) = \begin{cases} a & \text{if } a = b \\ \left[\frac{b^{p+1} - a^{p+1}}{(p+1)(b-a)} \right]^{\frac{1}{p}} & \text{if } a \neq b \end{cases} \quad \text{for } p \in \mathbb{R} \setminus \{0, -1\}.$$

– Identric mean,

$$I = I(a, b) = \begin{cases} a & \text{if } a = b \\ \frac{1}{e} \left(\frac{b^b}{a^a} \right)^{\frac{1}{b-a}} & \text{if } a \neq b \end{cases}.$$

The well known fact that $G < L < I < A$ will be used in the following.

1. Consider the function f with domain $[a, b] \subset (0, \infty)$, $f(x) = x^p$ and $p \in \mathbb{R}$, $p \geq 2$, which is absolutely continuous, and whose modulus of the first derivative is a convex function.

1.1 If we use this function in Corollary 1, we get that

$$\left| \left(\frac{a+b}{2} \right)^p - \frac{1}{b-a} \int_a^b t^p dt \right| \leq \frac{1}{8} (b-a) \left[\left| p \left(\frac{a+b}{2} \right)^{p-1} \right| + pb^{p-1} \right]$$

so that

$$|A^p(a, b) - L_p^p(a, b)| \leq \frac{p}{8} (b-a) [A^{p-1}(a, b) + b^{p-1}]$$

or equivalently

$$0 \leq L_p^p(a, b) - A^p(a, b) \leq \frac{p}{8} (b-a) [A^{p-1}(a, b) + b^{p-1}].$$

1.2 For the same function, we get from Corollary 3 that

$$\begin{aligned} |A^p(a, b) - L_p^p(a, b)| &\leq \frac{1}{4} \left[(b-a) \left| p \left(\frac{a+b}{2} \right)^{p-1} \right| + \int_a^b |pt^{p-1}| dt \right] \\ &= \frac{p}{4} \left[(b-a) A^{p-1}(a, b) + \frac{b^p - a^p}{p} \right] \\ &= \frac{p}{4} (b-a) \left[A^{p-1}(a, b) + L_{p-1}^{p-1}(a, b) \right]. \end{aligned}$$

That is,

$$0 \leq L_p^p(a, b) - A^p(a, b) \leq \frac{p}{4}(b-a) \left[A^{p-1}(a, b) + L_{p-1}^{p-1}(a, b) \right].$$

2. Now, consider the function f with domain $[a, b] \subset (0, \infty)$, $f(x) = \ln(x)$. The function is absolutely continuous, and the modulus of the first derivative is convex.

2.1 From Corollary 1, we obtain,

$$\begin{aligned} 0 &\leq \left| \ln \left(\frac{a+b}{2} \right) - \frac{1}{b-a} \int_a^b \ln(t) dt \right| = \left| \ln \left(\frac{a+b}{2} \right) - \ln I(a, b) \right| \\ &= \ln \frac{A(a, b)}{I(a, b)} \\ &\leq \frac{1}{8}(b-a) \left[\left| \frac{2}{a+b} \right| + \frac{1}{a} \right] \\ &= \frac{1}{8}(b-a) \left[A^{-1}(a, b) + \frac{1}{a} \right] \end{aligned}$$

and so

$$0 \leq \ln \frac{A(a, b)}{I(a, b)} \leq \frac{b-a}{8} [A^{-1}(a, b) + a^{-1}]$$

or, equivalently,

$$1 \leq \frac{A(a, b)}{I(a, b)} \leq \exp \left[\frac{b-a}{8} [A^{-1}(a, b) + a^{-1}] \right].$$

2.2 From Corollary 3, we get that

$$\begin{aligned} 0 &\leq |\ln A(a, b) - \ln I(a, b)| = \ln \frac{A(a, b)}{I(a, b)} \\ &\leq \frac{1}{4} \left[(b-a) \left| \frac{2}{a+b} \right| + \int_a^b \left| \frac{1}{t} \right| dt \right]. \end{aligned}$$

That is,

$$0 \leq \ln \frac{A(a, b)}{I(a, b)} \leq \frac{b-a}{4} A^{-1}(a, b) + \frac{1}{4} \ln \frac{b}{a}$$

or, equivalently,

$$1 \leq \frac{A(a, b)}{I(a, b)} \leq \left(\frac{b}{a} \right)^{\frac{1}{4}} \exp \frac{b-a}{4} [A^{-1}(a, b)].$$

2.3 Taking $f(x) = \ln x$ in Corollary 4, gives

$$1 \leq \frac{A(a, b)}{I(a, b)} \leq \frac{b-a}{4} \left[\frac{A^{-1}(a, b)}{2} + \frac{1}{(q+1)^{\frac{1}{q}}} L_{-p}^{-1}(a, b) \right].$$

3. Now, consider the function $f(x) = \frac{1}{x}$ which has domain $[a, b] \subset (0, \infty)$. This function is absolutely continuous and the modulus of the first derivative is convex.

3.1 From Corollary 1, we have,

$$\left| \frac{2}{a+b} - L^{-1}(a, b) \right| \leq \frac{1}{8}(b-a) \left[\left| \frac{1}{\left(\frac{a+b}{2} \right)^2} \right| + \frac{1}{a^2} \right]$$

giving

$$|A^{-1}(a, b) - L^{-1}(a, b)| \leq \frac{1}{8}(b-a) [A^{-2}(a, b) + a^{-2}]$$

or equivalently

$$0 \leq L^{-1}(a, b) - A^{-1}(a, b) \leq \frac{1}{8}(b-a) [A^{-2}(a, b) + a^{-2}], \quad (\text{since } A(a, b) \geq L(a, b))$$

which may further be represented as

$$0 \leq A(a, b) - L(a, b) \leq \frac{1}{8}(b-a)A(a, b)L(a, b) [A^{-2}(a, b) + a^{-2}].$$

3.2 From Corollary 3, we get,

$$|A^{-1}(a, b) - L^{-1}(a, b)| \leq \frac{1}{4}(b-a) [A^{-2}(a, b) + G^{-2}(a, b)]$$

or equivalently

$$0 \leq L^{-1}(a, b) - A^{-1}(a, b) \leq \frac{1}{4}(b-a) [A^{-2}(a, b) + G^{-2}(a, b)]$$

or still further

$$0 \leq A(a, b) - L(a, b) \leq \frac{1}{4}(b-a)A(a, b)L(a, b) [A^{-2}(a, b) + G^{-2}(a, b)].$$

3.3 Taking $f(x) = \frac{1}{x}$ in Corollary 4, produces

$$|A^{-1}(a, b) - L^{-1}(a, b)| \leq \frac{b-a}{4} \left[\frac{A^{-2}(a, b)}{2} + \frac{1}{(q+1)^{\frac{1}{q}}} L_{-2p}^{-2}(a, b) \right]$$

or

$$0 \leq L^{-1}(a, b) - A^{-1}(a, b) \leq \frac{b-a}{4} \left[\frac{A^{-2}(a, b)}{2} + \frac{1}{(q+1)^{\frac{1}{q}}} L_{-2p}^{-2}(a, b) \right]$$

which may be further expressed as

$$0 \leq A(a, b) - L(a, b) \leq \frac{b-a}{4}A(a, b)L(a, b) \left[\frac{A^{-2}(a, b)}{2} + \frac{1}{(q+1)^{\frac{1}{q}}} L_{-2p}^{-2}(a, b) \right].$$

4. APPLICATIONS FOR f AND HH -DIVERGENCE MEASURES IN INFORMATION THEORY

Assume that a set χ and the σ -finite measure $\mu : \chi \rightarrow \bar{\mathbb{R}}$ are given. Consider the set of all probability densities on μ to be

$$(4.1) \quad \Omega := \left\{ p|p : \chi \rightarrow \mathbb{R}, p(x) \geq 0, \int_{\chi} p(x) d\mu(x) = 1 \right\}.$$

The f -divergence on Ω is defined as follows

$$(4.2) \quad D_f(p, q) := \int_{\chi} p(x) f \left[\frac{q(x)}{p(x)} \right] d\mu(x), \quad p, q \in \Omega,$$

where f is convex on $(0, \infty)$. It is also assumed that $f(u)$ is zero and strictly convex at $u = 1$.

By appropriately defining this convex function, various divergences such as the Kullback-Leibler divergence D_{KL} , variation distance D_v , Hellinger distance D_H ,

χ^2 -divergence D_{χ^2} , Jeffrey's distance D_J , triangular discrimination D_Δ , etc. may be obtained. They are defined as follows:

$$(4.3) \quad D_v(p, q) := \int_{\chi} |p(x) - q(x)| d\mu(x), \quad p, q \in \Omega;$$

$$(4.4) \quad D_H(p, q) := \int_{\chi} \left| \sqrt{p(x)} - \sqrt{q(x)} \right| d\mu(x), \quad p, q \in \Omega;$$

$$(4.5) \quad D_{\chi^2}(p, q) := \int_{\chi} p(x) \left[\left(\frac{q(x)}{p(x)} \right)^2 - 1 \right] d\mu(x), \quad p, q \in \Omega;$$

$$(4.6) \quad D_J(p, q) := \int_{\chi} [p(x) - q(x)] \ln \left[\frac{p(x)}{q(x)} \right] d\mu(x), \quad p, q \in \Omega;$$

$$(4.7) \quad D_\Delta(p, q) := \int_{\chi} \frac{[p(x) - q(x)]^2}{p(x) + q(x)} d\mu(x), \quad p, q \in \Omega.$$

In [6], Shioya and Da-te introduced the generalised Ling-Wong f -divergence $D_f(p, \frac{1}{2}p + \frac{1}{2}q)$ and the Hermite-Hadamard (HH)-divergence

$$(4.8) \quad D_{HH}^f(p, q) := \int_{\chi} \frac{p^2(x)}{q(x) - p(x)} \left(\int_1^{\frac{q(x)}{p(x)}} f(t) dt \right) d\mu(x), \quad p, q \in \Omega.$$

They proved, by the use of the Hermite-Hadamard inequality for convex functions,

$$(4.9) \quad D_f \left(p, \frac{1}{2}p + \frac{1}{2}q \right) \leq D_{HH}^f(p, q) \leq \frac{1}{2} D_f(p, q),$$

provided that f is convex and normalised, i.e., $f(1) = 0$.

We will illustrate the approach to developing bounds and expressions involving various divergence measures from the inequalities developed in Section 2.

We will use the inequality (2.5), namely

$$(4.10) \quad \left| f \left(\frac{a+b}{2} \right) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \frac{1}{8} |b-a| \left[\left| f' \left(\frac{a+b}{2} \right) \right| + \|f'\|_{\infty} \right],$$

where $a, b \in \overset{\circ}{I}$, $a \neq b$ and $f : \overset{\circ}{I} \subset \mathbb{R} \rightarrow \mathbb{R}$ is a differentiable function on the interior of I with $|f'| : \overset{\circ}{I} \rightarrow \mathbb{R}$ convex on $\overset{\circ}{I}$, to prove the following result.

Theorem 6. *Let r, R be such that $0 \leq r \leq 1 \leq R \leq \infty$ and $p, q \in \Omega$ with*

$$(4.11) \quad r \leq \frac{q(x)}{p(x)} \leq R, \quad \text{for a.e. } x \in \chi.$$

If $f : [0, \infty) \rightarrow \mathbb{R}$ is differentiable on $(0, \infty)$ and $|f'|$ is convex on $[r, R]$ then,

$$(4.12) \quad \begin{aligned} & \left| D_f \left(p, \frac{1}{2}p + \frac{1}{2}q \right) - D_{HH}^f(p, q) \right| \\ & \leq \frac{1}{8} \left[\|f'\|_{[r, R], \infty} D_v(p, q) + D_{f^*}(p, q) \right], \end{aligned}$$

where $f^(x) = |x-1| |f'(\frac{x+1}{2})|$, $x \in [r, R]$ and $\|h\|_{[a, b], \infty} := \text{ess sup}_{t \in [a, b]} |h(t)|$.*

Proof. If in (4.10) we choose $a = 1$, $b = \frac{q(x)}{p(x)}$, $x \in \chi$, then

$$(4.13) \quad \left| f\left(\frac{p(x)+q(x)}{2p(x)}\right) - \frac{p(x)}{q(x)-p(x)} \left(\int_1^{\frac{q(x)}{p(x)}} f(t) dt \right) \right| \\ \leq \frac{1}{8} \cdot \frac{|q(x)-p(x)|}{p(x)} \left[\left| f'\left(\frac{p(x)+q(x)}{2p(x)}\right) \right| + \|f'\|_{[r,R],\infty} \right].$$

Multiplying (4.13) with $p(x) \geq 0$ and integrating on χ , we deduce the desired inequality (4.12). ■

Another approach is embodied in the following theorem.

Theorem 7. *Let r, R be as in Theorem 6. If $f : [0, \infty) \rightarrow \mathbb{R}$ is twice differentiable on $(0, \infty)$ and $|f''|$ is convex on $[r, R]$, then,*

$$(4.14) \quad |D_f(p, q) - f(1) - D_{f^\#}(p, q)| \leq \frac{1}{8} \left[\|f'\|_{[r,R],\infty} D_{\chi^2}(p, q) + D_{f^\dagger}(p, q) \right],$$

where $f^\#(x) := (x-1)f'(\frac{1+x}{2})$, and $f^\dagger(x) := (x-1)^2 |f''(\frac{1+x}{2})|$, $x \in [0, \infty)$.

Proof. Applying the inequality (4.10) for $a = 1$, $b = u$ and choosing instead of f , its derivative f' , one may state the inequality

$$\left| f(u) - f(1) - (u-1)f'\left(\frac{u+1}{2}\right) \right| \\ \leq \frac{1}{8} (u-1)^2 \left[\left| f''\left(\frac{u+1}{2}\right) \right| + \|f'\|_{[r,R],\infty} \right].$$

If in this inequality we choose $u = \frac{q(x)}{p(x)}$, $x \in \chi$, then we get

$$(4.15) \quad \left| f\left(\frac{q(x)}{p(x)}\right) - f(1) - \left(\frac{q(x)}{p(x)} - 1\right) f'\left(\frac{p(x)+q(x)}{2p(x)}\right) \right| \\ \leq \frac{1}{8} \frac{(p(x)-q(x))^2}{p^2(x)} \left[\left| f''\left(\frac{p(x)+q(x)}{2p(x)}\right) \right| + \|f'\|_{[r,R],\infty} \right].$$

Multiplying (4.15) by $p(x) \geq 0$, $x \in \chi$ and then integrating on χ , we deduce the desired inequality (4.14). ■

REFERENCES

- [1] S.S. DRAGOMIR, An Ostrowski type inequality for isotonic linear functionals, submitted.
- [2] S.S. DRAGOMIR and S. WANG, A new inequality of Ostrowski's type in L_p -norm and applications to some special means and to some numerical quadrature rules, *Tamkang J. of Math.*, **28** (1997), 239-244.
- [3] S.S. DRAGOMIR and S. WANG, A new inequality of Ostrowski's type in L_p -norm, *Indian J. Math.*, **40**(3) (1998), 293-304.
- [4] S.S. DRAGOMIR and S. WANG, Applications of Ostrowski's inequality to the estimation of error bounds for some special means and some numerical quadrature rules, *Appl. Math. Lett.*, **11** (1998), 105-109.
- [5] A.M. FINK, Bounds on the derivation of a function from its averages, *Czech. Math. Journal.*, **42** (1992), No. 117, 289-310.
- [6] H. SHIOYA and T. DA-TE, A generalization of Lin divergence and the derivation of a new information divergence, *Elec. Comm. in Japan*, **78**(7) (1995), 37-40.

SCHOOL OF COMMUNICATIONS AND INFORMATICS, VICTORIA UNIVERSITY OF TECHNOLOGY, PO
BOX 14428, MCMC 8001, VIC, AUSTRALIA.

E-mail address: `neil@matilda.vu.edu.au`

URL: `http://sci.vu.edu.au/staff/neilb.html`

E-mail address: `pc@matilda.vu.edu.au`

URL: `http://rgmia.vu.edu.au/cerone`

E-mail address: `sever@matilda.vu.edu.au`

URL: `http://rgmia.vu.edu.au/SSDragomirWeb.html`

E-mail address: `marcia@matilda.vu.edu.au`

URL: `http://www.geocities.com/msorfiap`

E-mail address: `sofo@matilda.vu.edu.au`

URL: `http://cams.vu.edu.au/staff/anthonys.html`