

Journal: Regional Environmental Change

Article: Runoff from glacier ice and seasonal snow in High Asia: separating melt water sources in river flow

Authors: Richard L. Armstrong^{1,2}, Karl Rittger^{1,2}, Mary J. Brodzik^{1,2}, Adina Racoviteanu^{1,4}, Andrew P. Barrett^{1,2}, Siri-Jodha Singh Khalsa^{1,2}, Bruce Raup^{1,2}, Alice F. Hill^{1,2}, Alia Khan^{1,2}, Alana M. Wilson^{1,2}, Rijan Bhakta Kayastha⁴, Florence Fetterer^{1,2}, Betsy Armstrong^{1,2}














Affiliations: ¹National Snow and Ice Data Center, ²University of Colorado, ³Kathmandu University, ⁴Aberystwyth University, ⁵Institute of Arctic and Alpine Research

Corresponding author: Richard L. Armstrong, rlax@nsidc.org, 303-494-7040














[Figure found on the following 4 pages]

Fig. S3 (a-d): Comparison of uncorrected downscaled European Reanalysis (ERA)-Interim temperatures with observed temperatures at 13 meteorological sites in the Upper Indus basin for the period 2001 to 2010 for a) winter (DJF), b) spring (MAM), c) summer (JJA) and d) autumn (SON). The mean seasonal cycle has been removed. Correlation coefficient and root mean square error (RMSE) is shown for each site.














(a)



(b)



(c)



(d)

