

Journal: Regional Environmental Change

Article: Runoff from glacier ice and seasonal snow in High Asia: separating melt water sources in river flow

Authors: Richard L. Armstrong^{1,2}, Karl Rittger^{1,2}, Mary J. Brodzik^{1,2}, Adina Racoviteanu^{1,4}, Andrew P. Barrett^{1,2}, Siri-Jodha Singh Khalsa^{1,2}, Bruce Raup^{1,2}, Alice F. Hill^{1,2}, Alia Khan^{1,2}, Alana M. Wilson^{1,2}, Rijan Bhakta Kayastha⁴, Florence Fetterer^{1,2}, Betsy Armstrong^{1,2}

Affiliations: ¹National Snow and Ice Data Center, ²University of Colorado, ³Kathmandu University, ⁴Aberystwyth University, ⁵Institute of Arctic and Alpine Research

Corresponding author: Richard L. Armstrong, [rlax@nsidc.org](mailto:r lax@nsidc.org), 303-494-7040

Table S3: Melt model calibration basin degree-day factors (DDFs) for best overall models, years, and calibration statistics. Minimum and maximum snow (ice) DDFs used as extrema for seasonally-adjusted sine curve for melt model snow (ice) DDFs.

Basin	Major Basin	Min Snow DDF (mm °C ⁻¹ day ⁻¹)	Max Snow DDF (mm °C ⁻¹ day ⁻¹)	Min Ice DDF (mm °C ⁻¹ day ⁻¹)	Max Ice DDF (mm °C ⁻¹ day ⁻¹)	Calibration Years	Annual Volumetric Difference (%)	Monthly RMSE (km ³)
Naryn	SY	4.61	5.49	53.30	57.16	2003, 2005, 2007	0.4	0.25
Vakhsh	AM	2.38	4.86	55.86	56.94	2004, 2005, 2007	2.6	1.27
Hunza	IN	10.41	10.48	37.94	42.30	2001, 2002, 2003	0.1	0.39
Karnali	GA	17.42	17.60	57.32	57.41	2001, 2003, 2004	3.8	1.25
SaptaKosi	BR	32.98	33.09	35.52	49.53	2003, 2004, 2005	0.1	2.82