

Supplementary information 2 (Table) | Mammalian Bicarbonate Transporters.

Gene/common name	Tissue distribution/epithelial localization	Transport mode	Murine null phenotype	Refs
SLC4a1/AE1, Band 3	Erythrocyte, kidney, heart/ <i>kAE1 Basolateral</i>	$\text{Cl}^-/\text{HCO}_3^-$ exchange/electroneutral	Hemolytic anemia in two independent mouse AE1 ^{-/-} lines, also seen in spontaneous bovine mutant.	1-3 4
SLC4a2/AE2	Widespread/ <i>Basolateral</i>	$\text{Cl}^-/\text{HCO}_3^-$ exchange/electroneutral	Loss of stomach acid secretion, failed dentition; altered immune function.	5, 6 7
SLC4a3/AE3	Brain, heart, retina, pituitary, adrenal gland; Non-epithelial	$\text{Cl}^-/\text{HCO}_3^-$ exchange /electroneutral	Inner retinal defects, similar to human vitreoretinal degeneration syndromes; sensitivity to chemical-induced seizures.	8-10
SLC4a4/NBCe1	Pancreas, kidney, heart, cornea, prostate, colon, stomach, thyroid, brain/ <i>Basolateral</i>	$\text{Na}^+/\text{HCO}_3^-$ co-transport/electrogenic	Metabolic acidosis, runting, splenomegaly, altered dentition, intestinal obstruction; death before weaning.	11 12-17
SLC4a5/NBCe2	Brain (especially prefrontal cortex), epididymis, cardiac muscle, smooth muscle, kidney, choroid plexus/ <i>Apical</i>	$\text{Na}^+/\text{HCO}_3^-$ co-transport/electrogenic	No mouse model.	18-21
SLC4a7/NBCn1	Heart, kidney, skeletal muscle, smooth muscle, submandibular gland, pancreas, stomach/ <i>Basolateral</i>	$\text{Na}^+/\text{HCO}_3^-$ co-transport/electroneutral	Blindness and auditory defect.	22 23-25
SLC4a9	Kidney, testis, pancreas, widespread/ <i>Apical</i>	$\text{Na}^+/\text{HCO}_3^-$ co-transport/electroneutral	No mouse model.	26-29
SLC4a8/NDCBE	Prefrontal cortex of brain, testis, cardiac myocytes, oocytes	Na^+ -dependent $\text{Cl}^-/\text{HCO}_3^-$ exchange and (electroneutral NBC: splice variant)/electroneutral	No mouse model.	30-32
SLC4a10/NBCn2	Cardiac myocytes, neurons, kidney, uterus, adrenal cortex, choroid plexus/ <i>Basolateral</i>	$\text{Na}^+/\text{HCO}_3^-$ co-transport or Na^+ -dependent $\text{Cl}^-/\text{HCO}_3^-$ exchange/electroneutral	Epileptic seizures, reduced brain ventricle volume 2° to choroid plexus defect.	33-36
SLC26a3/DRA	Colon, ileum, cardiac myocytes eccrine sweat gland/ <i>Apical</i>	$\text{Cl}^-/\text{HCO}_3^-$ exchange/ electroneutral	High chloride diarrhea.	37-39
SLC26a4/Pendrin	Thyroid, inner ear, thyroid, kidney, prostate/ <i>Apical</i>	$\text{Cl}^-/\text{HCO}_3^-$ exchange/ electrogenic	Deafness with vestibular dysfunction; inner ear endolymph dilatation.	40 41-43
SLC26a6/PAT-1	Kidney, heart, bronchial epithelium, pancreas, prostate, thymus, intestine/ <i>Apical</i>	$\text{Cl}^-/\text{HCO}_3^-$ exchange/electrogenic	Calcium oxalate urolithiasis, hyperoxaluria, high plasma oxalate.	44 45-47
SLC26a7	Thyroid, kidney, stomach, retina, olfactory epithelium/ <i>Basolateral</i>	$\text{Cl}^-/\text{HCO}_3^-$ exchange/electrogenic	No mouse model.	48-50
SLC26a9	Salivary gland, heart, brain, stomach, trachea, kidney/ <i>Apical</i>	$\text{Cl}^-/\text{HCO}_3^-$ exchange/electrogenic	No mouse model.	51, 52

- Southgate, C.D., Chishti, A.H., Mitchell, B., Yi, S.J. & Palek, J. Targeted disruption of the murine erythroid band 3 gene results in spherocytosis and severe haemolytic anaemia despite a normal membrane skeleton. *Nat. Genet.* **14**, 227-230 (1996).
- Peters, L.L. *et al.* Anion exchanger 1 (band 3) is required to prevent erythrocyte membrane surface loss but not to form the membrane skeleton. *Cell* **86**, 917-927 (1996).
- Inaba, M. *et al.* Defective anion transport and marked spherocytosis with membrane instability caused by hereditary total deficiency of red cell band 3 in cattle due to a nonsense mutation. *J. Clin. Invest.* **97**, 1804-1817 (1996).
- Kopito, R.R. & Lodish, H.F. Primary structure and transmembrane orientation of the murine anion exchange protein. *Nature* **316**, 234-238 (1985).
- Gawenis, L.R. *et al.* Mice with a targeted disruption of the AE2 $\text{Cl}^-/\text{HCO}_3^-$ exchanger are achlorhydric. *J Biol. Chem.* **279**, 30531-30539 (2004).
- Salas, J.T. *et al.* Ae2a,b-deficient mice develop antimitochondrial antibodies and other features resembling primary biliary cirrhosis. *Gastroenterology* **134**, 1482-1493 (2008).
- Alper, S.L., Kopito, R.R., Libresco, S.M. & Lodish, H.F. Cloning and characterization of a murine Band 3-related cDNA from kidney and a lymphoid cell line. *J. Biol. Chem.* **263**, 17092-17099 (1988).
- Hentschke, M. *et al.* Mice with a targeted disruption of the $\text{Cl}^-/\text{HCO}_3^-$ exchanger AE3 display a reduced seizure threshold. *Mol Cell Biol.* **26**, 182-191 (2006).
- Alvarez, B.V. *et al.* Blindness results from deficiency in AE3 chloride/bicarbonate exchanger. *PLoS ONE* **2**, e839 (2007).
- Kudrycki, K.E., Newman, P.R. & Shull, G.E. cDNA cloning and tissue distribution of mRNAs for two proteins that are related to the Band 3 $\text{Cl}^-/\text{HCO}_3^-$ exchanger. *J. Biol. Chem.* **265**, 462-471 (1990).
- Gawenis, L.R. *et al.* Colonic anion secretory defects and metabolic acidosis in mice lacking the NBC1 $\text{Na}^+/\text{HCO}_3^-$ cotransporter. *J. Biol. Chem.* **282**, 9042-9052 (2007).
- Burnham, C.E., Amlal, H., Wang, Z., Shull, G.E. & Soleimani, M. Cloning and functional expression of a human kidney $\text{Na}^+/\text{HCO}_3^-$ cotransporter. *J. Biol. Chem.* **272**, 19111-19114 (1997).
- Romero, M.F., Hediger, M.A., Boulpaep, E.L. & Boron, W.F. Expression cloning and characterization of a renal electrogenic $\text{Na}^+/\text{HCO}_3^-$ cotransporter. *Nature* **387**, 409-413 (1997).
- Abuladze, N. *et al.* Molecular cloning, chromosomal localization, tissue distribution, and functional expression of the human pancreatic sodium bicarbonate cotransporter. *J. Biol. Chem.* **273**, 17689-17695 (1998).
- Choi, I., Romero, M.F., Khandoudi, N., Bril, A. & Boron, W.F. Cloning and characterization of a renal $\text{Na}^+/\text{HCO}_3^-$ cotransporter. *J. Biol. Chem.* **274**, 21231-21237 (1999).

SUPPLEMENTARY INFORMATION

- W.F. Cloning and characterization of a human electrogenic Na^+ - HCO_3^- cotransporter isoform (hhNBC). *Am. J. Physiol.* **276**, C576-584 (1999).
16. Bevensee, M.O., Schmitt, B.M., Choi, I., Romero, M.F. & Boron, W.F. An electrogenic Na^+ - HCO_3^- cotransporter (NBC) with a novel COOH-terminus, cloned from rat brain. *Am. J. Physiol.* **278**, C1200-C1211 (2000).
 17. Sun, X.C. & Bonanno, J.A. Identification and cloning of the Na/HCO_3^- cotransporter (NBC) in human corneal endothelium. *Exp Eye Res* **77**, 287-295 (2003).
 18. Pushkin, A. *et al.* Cloning, characterization and chromosomal assignment of NBC4, a new member of the sodium bicarbonate cotransporter family. *Biochim Biophys Acta* **1493**, 215-218. (2000).
 19. Sassani, P. *et al.* Functional characterization of NBC4: a new electrogenic sodium-bicarbonate cotransporter. *Am J Physiol Cell Physiol* **282**, C408-416 (2002).
 20. Virkki, L.V., Wilson, D.A., Vaughan-Jones, R.D. & Boron, W.F. Functional characterization of human NBC4 as an electrogenic Na^+ - HCO_3^- cotransporter (NBCe2). *Am J Physiol Cell Physiol* **282**, C1278-1289 (2002).
 21. Xu, J. *et al.* Expression of the Na^+ - HCO_3^- cotransporter NBC4 in rat kidney and characterization of a novel NBC4 variant. *Am J Physiol Renal Physiol* **284**, F41-50 (2003).
 22. Bok, D. *et al.* Blindness and auditory impairment caused by loss of the sodium bicarbonate cotransporter NBC3. *Nat Genet* **34**, 313-319 (2003).
 23. Ishibashi, K., Sasaki, S. & Marumo, F. Molecular cloning of a new sodium bicarbonate cotransporter cDNA from human retina. *Biochem. Biophys. Res. Commun.* **246**, 535-538 (1998).
 24. Pushkin, A. *et al.* Cloning, tissue distribution, genomic organization, and functional characterization of NBC3, a new member of the sodium bicarbonate cotransporter family. *J. Biol. Chem.* **274**, 16569-16575 (1999).
 25. Choi, I., Aalkjaer, C., Boulaep, E.L. & Boron, W.F. An electroneutral sodium/bicarbonate cotransporter NBCn1 and associated sodium channel. *Nature* **405**, 571-575 (2000).
 26. Tsuganezawa, H. *et al.* A new member of the HCO_3^- transporter superfamily is an apical anion exchanger of beta-intercalated cells in the kidney. *J Biol Chem* **276**, 8180-8189 (2001).
 27. Parker, M.D., Ourmazdi, E.P. & Tanner, M.J. Human BTR1, a New Bicarbonate Transporter Superfamily Member and Human AE4 from Kidney. *Biochem Biophys Res Commun* **282**, 1103-1109. (2001).
 28. Romero, M.F., Fulton, C.M. & Boron, W.F. The SLC4 family of HCO_3^- transporters. *Pflugers Arch* **14**, 14 (2004).
 29. Parker, M.D., Boron, W.F. & Tanner, M.J. Characterization of human "AE4" as an electroneutral sodium bicarbonate cotransporter. *FASEB J* **16**, A796 (2002).
 30. Romero, M.F. *et al.* Cloning and characterization of a Na^+ driven anion exchanger (NDAE1): a new bicarbonate transporter. *J. Biol. Chem.* **275**, 24552-24559 (2000).
 31. Grichtchenko, II *et al.* Cloning, characterization, and chromosomal mapping of a human electroneutral Na^+ -driven $\text{Cl}^-/\text{HCO}_3^-$ exchanger. *J. Biol. Chem.* **276**, 8358-8363 (2001).
 32. Wang, Z. *et al.* Mouse $\text{Na}^+ : \text{HCO}_3^-$ cotransporter isoform NBC-3 (kNBC-3): cloning, expression, and renal distribution. *Kidney Int* **59**, 1405-1414 (2001).
 33. Jacobs, S. *et al.* Mice with targeted Slc4a10 gene disruption have small brain ventricles and show reduced neuronal excitability. *Proc Natl Acad Sci U S A* **105**, 311-316 (2008).
 34. Wang, C.Z., Yano, H., Nagashima, K. & Seino, S. The Na^+ -driven $\text{Cl}^-/\text{HCO}_3^-$ exchanger: Cloning, tissue distribution, and functional characterization. *J. Biol. Chem.* **275**, 35486-35490 (2000).
 35. Praetorius, J., Nejsum, L.N. & Nielsen, S. A SCL4A10 gene product maps selectively to the basolateral plasma membrane of choroid plexus epithelial cells. *Am J Physiol Cell Physiol* **286**, C601-610 (2004).
 36. Parker, M.D. *et al.* Characterization of human SLC4A10 as an electroneutral Na/HCO_3 cotransporter (NBCn2) with Cl^- self-exchange activity. *J. Biol. Chem.* **283**, 12777-12788 (2008).
 37. Schweinfest, C.W. *et al.* slc26a3 (dra)-deficient mice display chloride-losing diarrhea, enhanced colonic proliferation, and distinct up-regulation of ion transporters in the colon. *J. Biol. Chem.* **281**, 37962-37971 (2006).
 38. Melvin, J.E., Park, K., Richardson, L., Schultheis, P.J. & Shull, G.E. Mouse down-regulated in adenoma (DRA) is an intestinal $\text{Cl}^-/\text{HCO}_3^-$ exchanger and is up-regulated in colon of mice lacking the NHE3 Na^+/H^+ exchanger. *J. Biol. Chem.* **274**, 22855-22861 (1999).
 39. Barmeyer, C. *et al.* Characteristics of rat downregulated in adenoma (rDRA) expressed in HEK 293 cells. *Pflugers Arch* **454**, 441-450 (2007).
 40. Everett, L.A. *et al.* Targeted disruption of mouse Pds provides insight about the inner-ear defects encountered in Pendred syndrome. *Hum Mol Genet* **10**, 153-161 (2001).
 41. Scott, D.A., Wang, R., Kreman, T.M., Sheffield, V.C. & Karnishki, L.P. The Pendred syndrome gene encodes a chloride-iodide transport protein. *Nat Genet* **21**, 440-443 (1999).
 42. Soleimani, M. *et al.* Pendrin: an apical $\text{Cl}/\text{OH}/\text{HCO}_3^-$ exchanger in the kidney cortex. *Am J Physiol Renal Physiol* **280**, F356-364. (2001).
 43. Shcheynikov, N. *et al.* Slc26a4 functions as an electroneutral $\text{Cl}^-/\text{HCO}_3^-$ exchanger: Role of Slc26a4 and Slc26a6 in I^- and HCO_3^- secretion and in regulation of CFTR in the parotid duct. *J. Physiol. In Press* (2008).
 44. Jiang, Z. *et al.* Calcium oxalate urolithiasis in mice lacking anion transporter Slc26a6. *Nat Genet* **38**, 474-478 (2006).
 45. Waldegger, S. *et al.* Cloning and characterization of SLC26A6, a novel member of the solute carrier 26 gene family. *Genomics* **72**, 43-50 (2001).
 46. Wang, Z., Petrovic, S., Mann, E. & Soleimani, M. Identification of an apical Cl/HCO_3^- exchanger in the small intestine. *Am J Physiol Gastrointest Liver Physiol* **282**, G573-579 (2002).
 47. Xie, Q., Welch, R., Mercado, A., Romero, M.F. & Mount, D.B. Molecular characterization of the murine Slc26a6 anion exchanger: functional comparison with Slc26a1. *Am J Physiol Renal Physiol* **283**, F826-838 (2002).
 48. Lohi, H. *et al.* Functional characterization of three novel tissue-specific anion exchangers: SLC26A7, A8 and A9. *J. Biol. Chem.* **277**, 14246-14254 (2002).
 49. Vincent, J.B., Jullien, D., Kossida, S., Amalric, F. & Girard, J.P. Molecular Cloning of SLC26A7, a Novel Member of the SLC26 Sulfate/Anion Transporter Family, from High Endothelial Venules and Kidney. *Genomics* **79**, 249-256 (2002).
 50. Petrovic, S. *et al.* Identification of a basolateral $\text{Cl}^-/\text{HCO}_3^-$ exchanger specific to gastric parietal cells. *Am J Physiol Gastrointest Liver Physiol* **284**, G1093-1103 (2003).
 51. Xu, J. *et al.* SLC26A9 is expressed in gastric surface epithelial cells, mediates $\text{Cl}^-/\text{HCO}_3^-$ exchange, and is inhibited by NH_4^+ . *Am J Physiol Cell Physiol* **289**, C493-505 (2005).
 52. Dorwart, M.R., Shcheynikov, N., Wang, Y., Stippec, S. & Muallim, S. SLC26A9 is a Cl channel regulated by the WNK kinases. *J. Physiol.* **584**, 333-345 (2007).