

Title: Differential effect of HLA class-I *versus* class-II transgenes on human T and B cell reconstitution and function in NRG mice

Authors: Sai Majji, Wathsala Wijayalath, Soumya Shashikumar, Luis Pow-Sang, Eileen Villasante, Teodor D. Brumeanu and Sofia Casares

Supplementary Table S1.HLA genotype of cord bloods used for reconstitution of mice

Cord blood*	HLA-A	HLA-DRB1	Number of infused mice			
			DRAGA	A2	DRAG	NRG
Donor #1	02:01 33:01	04:0101 15:01	11	6	9	2
Donor #2	02:01 01:01	04:0101 04:04	11	3	13	0
Donor #3	02:01 29:01	04:0101 03:01	9	9	13	2
Donor #4	02:01 29:01	04:0101 13:0501	8	5	0	0
Donor #5	02:01 03:01	04:0101 08:01	0	2	0	0
Donor #6	02:01 01:01	04:0101 11:0102	4	5	0	0
Donor #7	02:01 01:03	04:0101 15:0101	4	0	8	3

DRAGA, NOD.HLA-A2.HLA-DR4.RagKO.IL2R γ cKO; DRAG, NOD.HLA-DR4.RagKO.IL2R γ cKO; A2, NOD.HLA-A2.RagKO.IL2R γ cKO, NRG, NOD.RagKO.IL2R γ cKO.