


Figure S1

Absolute bias of the frequentist estimators as a function of the number of primary studies



Note. Absolute bias of the frequentist estimators as a function of the number of primary studies, the amount of heterogeneity and the shape of the random-effects distribution. CA = Cochran estimator; MBH = Malzahn-Böhning-Holling estimator; SJ(CA) = Sidik-Jonkman estimator with prior CA estimation; MPM = median-unbiased Mandel-Paule estimator; SJ = Sidik-Jonkman estimator; MP = Mandel-Paule estimator; CA2 = two-step Cochran estimator; DL2 = two-step DerSimonian-Laird estimator; DLm = multistep DerSimonian-Laird estimator; HS(ss) = Hunter-Schmidt estimator weighted by sample size; ML = maximum likelihood estimator; REML = restricted maximum likelihood estimator; LCHr = Lin-Chu-Hodges r estimator; LCHm = Lin-Chu-Hodges m estimator; GENQM = median-unbiased generalized Q statistic estimator; DLp = positive DerSimonian-Laird estimator; DL = DerSimonian-Laird estimator; HS(k) = Hunter-Schmidt estimator corrected by small sample size; DLb = nonparametric bootstrap DerSimonian-Laird estimator; HS(iv) = Hunter-Schmidt estimator weighted by inversed variance; HM = Hartung-Makambi estimator.