

Gluino Particle, The Cousin Of The Higgs Boson

A team of scientists currently working at the Large Hadron Collider at the European Organization for Nuclear Research (CERN) announced that it has possibly discovered the existence of a particle integral to nature in a statement on Tuesday, Dec. 15, and again on Dec.16. [9]

In 2012, a proposed observation of the Higgs boson was reported at the Large Hadron Collider in CERN. The observation has puzzled the physics community, as the mass of the observed particle, 125 GeV, looks lighter than the expected energy scale, about 1 TeV. [8]

'In the new run, because of the highest-ever energies available at the LHC, we might finally create dark matter in the laboratory,' says Daniela. 'If dark matter is the lightest SUSY particle than we might discover many other SUSY particles, since SUSY predicts that every Standard Model particle has a SUSY counterpart.' [7]

The problem is that there are several things the Standard Model is unable to explain, for example the dark matter that makes up a large part of the universe. Many particle physicists are therefore working on the development of new, more comprehensive models. [6]

They might seem quite different, but both the Higgs boson and dark matter particles may have some similarities. The Higgs boson is thought to be the particle that gives matter its mass. And in the same vein, dark matter is thought to account for much of the 'missing mass' in galaxies in the universe. It may be that these mass-giving particles have more in common than was thought. [5]

The magnetic induction creates a negative electric field, causing an electromagnetic inertia responsible for the relativistic mass change; it is the mysterious Higgs Field giving mass to the particles. The Planck Distribution Law of the electromagnetic oscillators explains the electron/proton mass rate by the diffraction patterns. The accelerating charges explain not only the Maxwell Equations and the Special Relativity, but the Heisenberg Uncertainty Relation, the wave particle duality and the electron's spin also, building the bridge between the Classical and Relativistic Quantum Theories. The self maintained electric potential of the accelerating charges equivalent with the General Relativity space-time curvature, and since it is true on the quantum level also, gives the base of the Quantum Gravity.

Contents

Preface.....	2
Popular questions about the Higgs Field:	2
How can we answer these questions?	3
Physicists Have Discovered Evidence Of A Gluino Particle, The Cousin Of The Higgs Boson.....	3
Recent study predicts that Higgs particles are much heavier than earlier observation	4
Exploring the Higgs boson's dark side.....	5
The Higgs particle can disintegrate into particles of dark matter, according to new model	6
Will the Large Hadron Collider find dark matter?.....	7
The Classical Relativistic effect	9
The Relativistic Quantum Mechanics	9
The Heisenberg Uncertainty Relation.....	9
The General Relativity - Electromagnetic inertia and mass.....	9
Electromagnetic Induction	9
Relativistic change of mass.....	9
The frequency dependence of mass	10
Electron – Proton mass rate	10
Higgs Field	10
The Higgs boson	10
What is the Spin?	11
The Weak Interaction	11
Higgs mechanism	11
Gravity from the point of view of quantum physics	12
The Gravitational force	12
The Graviton	12
Conclusions	13
References	13

Author: George Rajna

Preface

Popular questions about the Higgs Field:

1.) If the Higgs field is responsible for imbuing particles with mass, and mass is responsible for

gravity, is it possible that the Higgs field will provide the missing link between general relativity and quantum mechanics i.e. could the Higgs field be the basis of a quantum theory of gravity?

2.) Can the theoretical Higgs Field be used as the “cause” of relativistic momentum or relativistic kinetic energy of a moving body?

3.) Does Einstein's General Relativity need to be adjusted for the Higgs field?

4.) Since the Higgs field gives most particles mass, and permeates all space, then GR needs the Higgs field to be a theory of space?

5.) So where GR is highly curved, the Higgs field is also curved? And does a highly curved Higgs field affect the way particles acquire mass? For that matter, a curved space-time would also curve electromagnetic field?

How can we answer these questions?

There is an explanation of the magnetic effect caused by the electric current from the observed effects of the accelerating electrons, causing naturally the experienced changes of the electric field potential along the electric wire. The charge distribution is lowering in the reference frame of the accelerating charges linearly: $ds/dt = at$ (time coordinate), but in the reference frame of the current it is parabolic: $s = a/2 t^2$ (geometric coordinate). The accelerating electrons explain not only the Maxwell Equations and the Special Relativity, but the Heisenberg Uncertainty Relation, the wave particle duality and the electron's spin also, building the bridge between the Classical and Quantum Theories. [1]

One origin of the Quantum Physics is the Planck Distribution Law of the electromagnetic oscillators, giving equal intensity for 2 different wavelengths on any temperature. Any of these two wavelengths will give equal intensity diffraction patterns, building different asymmetric constructions, for example proton - electron structures (atoms), molecules, etc. Since the particles are centers of diffraction patterns they also have particle – wave duality as the electromagnetic waves have. The Electroweak Interaction shows that the Weak Interaction is basically electromagnetic in nature. The arrow of time shows the entropy grows by changing the temperature dependent diffraction patterns of the electromagnetic oscillators. [2]

Physicists Have Discovered Evidence Of A Gluino Particle, The Cousin Of The Higgs Boson

The two teams working in concert, named Atlas and CMS, presented their findings on the particle from the Large Hadron Collider's second run (LHC Run 2). The results were based on what the scientists observed during the particle collisions. The previously-hypothesized particle, named the gluino, is theoretically the supersymmetric partner of the gluon (or glue particle, which is comprised entirely of nuclear force). This would mean that the gluino could be pair-produced by colliders like the LHC, and would more or less be described as a heavier version of the Higgs boson, a particle that essentially helps us understand why other particles contain mass and was identified at the LHC at CERN in 2012.

Despite the promising findings, the scientists stressed that there was not enough data to pinpoint anything definitive about the existence of the possible — and possibly important — particle.

"I don't think there is anyone around who thinks this is conclusive," said Kyle Cranmer, one of the members of Atlas, in an interview with the New York Times.

"But it would be huge if true," added the NYU-based physicist.




Image of a particle collision.

(Photo : CERN) Image of a particle collision.

The scientists first noticed data that would point to the existence of the gluino with extra gamma rays found within the collider. The gamma rays, which were congruent with the charge power of about 750 billion electron volts, were theorized to be the product of radioactive decay in a particle. This behavior would be symptomatic of a gluino.

"We are barely coming to terms with the power and the glory," another CERN team member, Maria Spiropulu, told the New York Times via text message, referring to the collider's expansive capabilities, among them the "ability to operate at 13 trillion electron volts."

"We are now entering the era of taking a shot in the dark!" the physicist concluded. [9]

Recent study predicts that Higgs particles are much heavier than earlier observation

Researchers at Aalto University in Finland now propose that there is more than one Higgs boson, and they are much heavier than the 2012 observation. The results were recently published in Nature Communications.

"Our recent ultra-low temperature experiments on superfluid helium (^3He) suggest an explanation why the Higgs boson observed at CERN appears to be too light.

By using the superfluid helium analogy, we have predicted that there should be other Higgs bosons, which are much heavier (about 1 TeV) than previously observed," says Professor (emeritus) Grigory E. Volovik.

Prof. Volovik holds a position in the Low Temperature Laboratory at Aalto University and in Landau Institute, Moscow. He has received the international Simon Prize in 2004 for distinguished work in theoretical low temperature physics, and the Lars Onsager Prize in 2014 for outstanding research in theoretical statistical physics.

At the same time, the new CERN experiments have shown evidence of the second Higgs in just the suggested region (at 0.75 TeV). This evidence has immediately been commented and discussed in a large number of papers submitted to arXiv, an e-print service widely utilised by the physics community to distribute manuscripts of their unpublished work. [8]

Exploring the Higgs boson's dark side

Last month, after two years of preparation, the LHC began smashing its proton beams together at 13 Trillion electron Volts (TeV), close to double the energy achieved during its first run.

'We do not know what we will find next and that makes the new run even more exciting,' Daniela Bortoletto of Oxford University's Department of Physics, a member of the team running the LHC's ATLAS experiment, tells me. 'We hope to finally find some cracks in the Standard Model as there are many questions about our universe that it does not answer.'

One of the big questions concerns dark matter, the invisible 'stuff' that astrophysicists estimate makes up over 80% of the mass of the Universe. As yet nobody has identified particles of dark matter although physicists think it could be the lightest supersymmetric (SUSY) particle.

'In the new run, because of the highest-ever energies available at the LHC, we might finally create dark matter in the laboratory,' says Daniela. 'If dark matter is the lightest SUSY particle than we might discover many other SUSY particles, since SUSY predicts that every Standard Model particle has a SUSY counterpart.'

Then there's the puzzle of antimatter: in the early Universe matter and antimatter were created in equal quantities but now matter dominates the Universe.

'We still do not know what caused the emergence of this asymmetry,' Daniela explains. 'We have finally discovered the Higgs boson: this special particle, a particle that does not carry any spin, might decay to dark matter particles and may even explain why the Universe is matter dominated.'

Discovering the Higgs boson was a huge achievement but now the race is on to understand it: a prospect that Daniela is particularly excited about.

'This particle is truly fascinating,' she says. 'Spin explains the behaviour of elementary particles: matter particles like the electron have spin $1/2$ while force particles

like the photon, which is responsible for the electromagnetic interaction, have spin 1. Spin $1/2$ particles obey the Pauli principle that forbids electrons to be in the same quantum state.

'The Higgs is the first spin 0 particle, or as particle physicists would say the first 'scalar particle' we've found, so the Higgs is neither matter nor force.'

Because of its nature the Higgs could have an impact on cosmic inflation and the energy of a vacuum as well as explaining the mass of elementary particles.

Daniela tells me: 'Because of the Higgs the electron has mass, atoms can be formed, and we exist. But why do elementary particles have such different masses?

The data of run 2 will enable us to study, with higher precision, the decays of the Higgs boson and directly measure the coupling of the Higgs to quarks. It will also enable us to search for other particles similar to the Higgs and determine if the Higgs decays to dark matter.'

Daniela is one of 13 academics at Oxford working on ATLAS supported by a team of postdoctoral fellows, postgraduate students and engineering, technical, and computing teams. The Oxford group plays a lead role in operating the Semiconductor Tracker (SCT), most of which was assembled in an Oxford lab. This provides information on the trajectories of the particles produced when the LHC's beams collide, which was crucial to the discovery of the Higgs boson.

Whilst the next few years will see the Oxford group busy with research that exploits the LHC's new high-energy run, the team are also looking ahead to 2025 when the intensity or 'luminosity' of the beams will be increased.

The LHC is filled with 1,380 bunches of protons each containing almost a billion protons and colliding 40 million times per second. This means that every time two bunches of protons cross they generate not one collision but many, an effect called 'pile-up'.

'After this luminosity upgrade the LHC will operate at collision rates five to ten times higher than it does at present,' Daniela explains. 'In run 1 of the LHC we had a maximum of 37 pile-up collisions per crossing but with the upgrade to the High Luminosity LHC, or 'HL-LHC', this will increase to an average of 140 pile-up events in each bunch crossing.'

With the HL-LHC generating many more collisions, the international Oxford-led team are designing and prototyping parts of a new semiconductor tracker that will be needed to help reconstruct particles from the complex web of decay trails they leave inside the machine.

As the LHC ramps up both its energy and luminosity it promises to give scientists working on experiments such as ATLAS answers to some of the biggest questions in physics. One thing is certain: this new physics will also lead to a whole set of new questions about the matter that makes up us and the Universe around us. [7]

The Higgs particle can disintegrate into particles of dark matter, according to new model

The problem is that there are several things the Standard Model is unable to explain, for example the dark matter that makes up a large part of the universe. Many particle physicists are therefore working on the development of new, more comprehensive models.

One of them is Christoffer Petersson, who carries out research in theoretical particle physics at Chalmers University of Technology in Sweden and the Université Libre in Belgium. Together with two research colleagues he has proposed a particle model based on what is known as supersymmetry.

This model contains more elementary particles than the Standard Model, including dark matter particles. In addition, the model gives the Higgs particle different properties than the Standard Model predicts. The model proposes that the Higgs particle can disintegrate into a photon (a particle of light) and particles of dark matter. However, these properties are quite difficult to discover – you have to look for them specifically to have a chance of finding them.

But Christoffer Petersson is fortunate – his model has met with a response at CERN. Two independent experimental stations – Atlas and CMS – at the Large Hadron Collider are now looking for the very properties of the Higgs particle his model predicts. If the properties are there, it is a clear indication that the model fits.

“It’s a dream for a theorist in particle physics. LHC is the only place where the model can be tested. It’s even nicer that two independent experiments are going to do it,” says Christoffer Petersson.

In the first studies the volume of data was unfortunately too small for it to be possible to either confirm or reject Petersson’s model.

A Higgs particle has been created in an LHC detector and has then disintegrated into four muons (the four red lines). According to Christoffer Petersson’s model the Higgs particle can also disintegrate into a photon and particles of dark matter. Picture: CERN


A Higgs particle has been created in an LHC detector and has then disintegrated into four muons (the four red lines). According to Christoffer Petersson’s model the Higgs particle can also disintegrate into a photon and particles of dark matter.

“But we are already in full swing with new analyses in which we are testing his model in other ways and with more data. We congratulate Christoffer Petersson for having done an important job,” says Zeynep Demiragli at the CMS experiment at CERN.

After being closed down for a time for an upgrade, LHC will start up again in the spring of 2015. With higher energies in the accelerator, the experiments will finally gather sufficient data to evaluate Petersson’s model properly. He is on tenterhooks awaiting the results. [6]

Will the Large Hadron Collider find dark matter?

Atom smasher could soon solve one the universe's greatest mysteries, claims scientist. Dr Monica Dunford worked at Cern in Switzerland up until 2013. She was directly involved in the detection of the Higgs boson in 2012. Speaking to Mail Online she said of the time leading up to its discovery: 'I don't think there will be another time like that in my career for sure'. But she said that the Large Hadron Collider could find dark matter. And if it did it would be a 'bigger discovery' than the Higgs boson. In March 2015 the LHC will be restarted at double its previous power.



'One of the things I'm most interested in is creating and discovering dark matter,' Dr Dunford said. 'We know from measurements of cosmology that 25 per cent of the universe is dark matter and we have absolutely no idea what that is.' An illustration of dark matter in the universe is shown.

When physicists study the dynamics of galaxies and the movement of stars, they are confronted with a mystery.

If they only take visible matter into account, their equations simply don't add up: the elements that can be observed are not sufficient to explain the rotation of objects and the existing gravitational forces. There is something missing.

From this they deduced that there must be an invisible kind of matter that does not interact with light, but does, as a whole, interact by means of the gravitational force. Called 'dark matter', this substance appears to make up at least 80 per cent of the universe. Finding the Higgs boson was one of the primary goals of the LHC - but perhaps the LHC's most important moment is yet to come.

'One of the things I'm most interested in is creating and discovering dark matter,' Dr Dunford said. We know from measurements of cosmology that 25 per cent of the universe is dark matter and we have absolutely no idea what that is. For comparison, what we do know, electrons and protons, only count for four per cent. You have this huge chunk of a pie and no idea what it consists of.

One thing we could possibly produce would be a dark matter candidate via its decay products. Being able to produce it at the LHC would be a huge connection between our astronomical measurements and what we can produce in the laboratory. [5]

The Classical Relativistic effect

The moving charges are self maintain the electromagnetic field locally, causing their movement and this is the result of their acceleration under the force of this field.

In the classical physics the charges will distributed along the electric current so that the electric potential lowering along the current, by linearly increasing the way they take every next time period because this accelerated motion. [1]

The Relativistic Quantum Mechanics

The same thing happens on the atomic scale giving a dp impulse difference and a dx way difference between the different part of the not point like particles.

Commonly accepted idea that the relativistic effect on the particle physics it is the fermions' spin - another unresolved problem in the classical concepts. If the electric charges can move only with accelerated motions in the self maintaining electromagnetic field, once upon a time they would reach the velocity of the electromagnetic field. The resolution of this problem is the spinning particle, constantly accelerating and not reaching the velocity of light because the acceleration is radial.

The Heisenberg Uncertainty Relation

I think that we have a simple bridge between the classical and quantum mechanics by understanding the Heisenberg Uncertainty Relations. It makes clear that the particles are not point like but have a dx and dp uncertainty.

The General Relativity - Electromagnetic inertia and mass

Electromagnetic Induction

Since the magnetic induction creates a negative electric field as a result of the changing acceleration, it works as an electromagnetic inertia, causing an electromagnetic mass. [1]

Relativistic change of mass

The increasing mass of the electric charges the result of the increasing inductive electric force acting against the accelerating force. The decreasing mass of the decreasing acceleration is the result of the inductive electric force acting against the decreasing force. This is the relativistic mass change explanation, especially importantly explaining the mass reduction in case of velocity decrease.

The frequency dependence of mass

Since $E = h\nu$ and $E = mc^2$, $m = h\nu / c^2$ that is the m depends only on the ν frequency. It means that the mass of the proton and electron are electromagnetic and the result of the electromagnetic induction, caused by the changing acceleration of the spinning and moving charge! It could be that the m_0 inertial mass is the result of the spin, since this is the only accelerating motion of the electric charge. Since the accelerating motion has different frequency for the electron in the atom and the proton, they masses are different, also as the wavelengths on both sides of the diffraction pattern, giving equal intensity of radiation. [2]

Electron – Proton mass rate

There is an asymmetry between the mass of the electric charges, for example proton and electron, can understood by the asymmetrical Planck Distribution Law. This temperature dependent energy distribution is asymmetric around the maximum intensity, where the annihilation of matter and antimatter is a high probability event. In the maximum intensity no diffraction patterns with equal intensity that is no fermions only bosons. The asymmetric sides are creating different frequencies of electromagnetic radiations being in the same intensity level and compensating each other. One of these compensating ratios is the electron – proton mass ratio. The lower energy side has no compensating intensity level, it is the dark energy and the corresponding matter is the dark matter. The Planck distribution law explains the different frequencies of the proton and electron, giving equal intensity to different lambda wavelengths! Also since the particles are diffraction patterns they have some closeness to each other – can be seen as a gravitational force.

In Quantum Field Theory (QFT), particles are described by excitations of a quantum field that satisfies the appropriate quantum mechanical field equations.

The excitations of the quantum field mean diffraction patterns in my theory. [2]

Higgs Field

The Higgs mechanism is a result of something called a field that extends throughout space, even where no particles are present. This notion is probably most familiar to you from a magnetic field. You feel a force between a magnet and your refrigerator even when “nothing” is there. A field can fill “empty” space. The Higgs field extends throughout space. Elementary particles acquire their masses by interacting with this field. It is kind of like space is charged and particles get mass through their interactions with this charge.

The Higgs boson is not directly responsible for mass. The Higgs field is. The boson is a particle that tells us our understanding of this mechanism is correct. It also is a big clue as to where that field came from in the first place. Its discovery tells us that what we expected to be true was indeed correct, and it gives us clues as to what else might underlie the Standard Model. [4]

The Higgs boson

By March 2013, the particle had been proven to behave, interact and decay in many of the expected ways predicted by the Standard Model, and was also tentatively confirmed to have + parity and zero spin, two fundamental criteria of a Higgs boson, making it also the first known scalar particle to be discovered in nature, although a number of other properties were not fully proven and some partial results do not yet precisely match those expected; in some cases data is also still awaited or being analyzed.

In my opinion, the best explanation of the Higgs mechanism for a lay audience is the one invented by David Miller. You can find it here: <http://www.strings.ph.qmul.ac.uk/~jmc/epp/higgs3.html> .

The field must come first. The boson is an excitation of the field. So no field, no excitation. On the other hand in quantum field theory it is difficult to separate the field and the excitations.

The Higgs field is what gives particles their mass.

There is a video that gives an idea as to the Higgs field and the boson. It is here:

<http://www.youtube.com/watch?v=RIg1Vh7uPyw> . Note that this analogy isn't as good as the Miller one, but as is usually the case, if you look at all the analogies you'll get the best understanding of the situation.

What is the Spin?

So we know already that the new particle has spin zero or spin two and we could tell which one if we could detect the polarizations of the photons produced. Unfortunately this is difficult and neither ATLAS nor CMS are able to measure polarizations. The only direct and sure way to confirm that the particle is indeed a scalar is to plot the angular distribution of the photons in the rest frame of the centre of mass. A spin zero particles like the Higgs carries no directional information away from the original collision so the distribution will be even in all directions. This test will be possible when a much larger number of events have been observed. In the mean time we can settle for less certain indirect indicators.

The Weak Interaction

Since the Higgs boson is necessary to the W and Z bosons, the dipole change of the Weak interaction and the change in the magnetic effect caused gravitation must be conducted. The Wien law is also important to explain the Weak interaction, since it describes the T_{\max} change and the diffraction patterns change. [2]

Higgs mechanism

The magnetic induction creates a negative electric field, causing an electromagnetic inertia. Probably it is the mysterious Higgs field giving mass to the charged particles? We can think about the photon as an electron-positron pair, they have mass. The neutral particles are built from negative and positive charges, for example the neutron, decaying to proton and electron. The wave – particle duality makes sure that the particles are oscillating and creating magnetic induction as an inertial mass, explaining also the relativistic mass change. Higher frequency creates stronger magnetic induction, smaller frequency results lesser magnetic induction. It seems to me that the magnetic induction is the secret of the Higgs field.

In particle physics, the Higgs mechanism is a kind of mass generation mechanism, a process that gives mass to elementary particles. According to this theory, particles gain mass by interacting with the Higgs field that permeates all space. More precisely, the Higgs mechanism endows gauge bosons in a gauge theory with mass through absorption of Nambu–Goldstone bosons arising in spontaneous symmetry breaking.

The simplest implementation of the mechanism adds an extra Higgs field to the gauge theory. The spontaneous symmetry breaking of the underlying local symmetry triggers conversion of components of this Higgs field to Goldstone bosons which interact with (at least some of) the other fields in the theory, so as to produce mass terms for (at least some of) the gauge bosons. This mechanism may also leave behind elementary scalar (spin-0) particles, known as Higgs bosons.

In the Standard Model, the phrase "Higgs mechanism" refers specifically to the generation of masses for the W^\pm and Z weak gauge bosons through electroweak symmetry breaking. The Large Hadron Collider at CERN announced results consistent with the Higgs particle on July 4, 2012 but stressed that further testing is needed to confirm the Standard Model.

Gravity from the point of view of quantum physics

The Gravitational force

The gravitational attractive force is basically a magnetic force.

The same electric charges can attract one another by the magnetic force if they are moving parallel in the same direction. Since the electrically neutral matter is composed of negative and positive charges they need 2 photons to mediate this attractive force, one per charges. The Big Bang caused parallel moving of the matter gives this magnetic force, experienced as gravitational force.

Since graviton is a tensor field, it has spin = 2, could be 2 photons with spin = 1 together.

You can think about photons as virtual electron – positron pairs, obtaining the necessary virtual mass for gravity.

The mass as seen before a result of the diffraction, for example the proton – electron mass ratio $M_p = 1840 M_e$. In order to move one of these diffraction maximum (electron or proton) we need to intervene into the diffraction pattern with a force appropriate to the intensity of this diffraction maximum, means its intensity or mass.

The Big Bang caused acceleration created radial currents of the matter, and since the matter is composed of negative and positive charges, these currents are creating magnetic field and attracting forces between the parallel moving electric currents. This is the gravitational force experienced by the matter, and also the mass is result of the electromagnetic forces between the charged particles. The positive and negative charged currents attracts each other or by the magnetic forces or by the much stronger electrostatic forces!?

The gravitational force attracting the matter, causing concentration of the matter in a small space and leaving much space with low matter concentration: dark matter and energy.

There is an asymmetry between the mass of the electric charges, for example proton and electron, can understood by the asymmetrical Planck Distribution Law. This temperature dependent energy distribution is asymmetric around the maximum intensity, where the annihilation of matter and antimatter is a high probability event. The asymmetric sides are creating different frequencies of electromagnetic radiations being in the same intensity level and compensating each other. One of these compensating ratios is the electron – proton mass ratio. The lower energy side has no compensating intensity level, it is the dark energy and the corresponding matter is the dark matter.

The Graviton

In physics, the graviton is a hypothetical elementary particle that mediates the force of gravitation in the framework of quantum field theory. If it exists, the graviton is expected to be massless (because the gravitational force appears to have unlimited range) and must be a spin-2 boson. The spin follows from the fact that the source of gravitation is the stress-energy tensor, a second-rank tensor (compared to electromagnetism's spin-1 photon, the source of which is the four-current, a first-rank

tensor). Additionally, it can be shown that any massless spin-2 field would give rise to a force indistinguishable from gravitation, because a massless spin-2 field must couple to (interact with) the stress-energy tensor in the same way that the gravitational field does. This result suggests that, if a massless spin-2 particle is discovered, it must be the graviton, so that the only experimental verification needed for the graviton may simply be the discovery of a massless spin-2 particle. [3]

Conclusions

“If the model is found to fit, it would completely change our understanding of the fundamental building blocks of nature. If not, just the fact that they are willing to test my model at CERN is great,” he says. [6]

On whether it would be the LHC’s most important discovery to date, she said: ‘Personally yes. It would be a bigger discovery than the Higgs boson. ‘For the Higgs we had a very good concrete theoretical prediction; for dark matter we really have no idea what it would be.’

She added: ‘There is no particle that we know of today that can explain dark matter, let alone what dark energy might be. So if we could directly produce dark matter particles at the LHC this would be a huge step forward in our understanding of the composition of the universe!’ [5]

The electric currents causing self maintaining electric potential is the source of the special and general relativistic effects. The Higgs Field is the result of the electromagnetic induction. The Graviton is two photons together.

References

[1] The Magnetic field of the Electric current and the Magnetic induction

Author: George Rajna

Publication date: 15/09/2013

Website: Academia.edu

[http://academia.edu/3833335/The Magnetic field of the Electric current](http://academia.edu/3833335/The_Magnetic_field_of_the_Electric_current)

Last accessed: 28/02/2014

[2] 3 Dimensional String Theory

Author: George Rajna

Publication date: 13/06/2013

Website: Academia.edu

[http://academia.edu/3834454/3 Dimensional String Theory](http://academia.edu/3834454/3_Dimensional_String_Theory)

Last accessed: 28/02/2014

[3] Graviton Production By Two Photon and Electron-Photon Processes In Kaluza-Klein Theories With Large Extra Dimensions

Authors: [David Atwood](#), [Shaouly Bar-Shalom](#), [Amarjit Soni](#)

Journal reference: Phys.Rev. D61 (2000) 116011

Publication date: 21/09/1999

Website: Arxiv.org

<http://arxiv.org/abs/hep-ph/9909392>

Last accessed: 28/02/2014

[4] Explaining the Higgs – Interview with Lisa Randall, Professor of Science in Harvard’s Department of Physics

By Sarah Sweeney, Harvard Staff Writer

Publication date: 13/01/2014

Website: Harvard Gazette

<http://news.harvard.edu/gazette/story/2014/01/explaining-the-higgs/>

Last accessed: 28/02/2014

[5] <http://www.dailymail.co.uk/sciencetech/article-2891468/Will-Large-Hadron-Collider-dark-matter-Atom-smasher-soon-solve-one-universe-s-greatest-mysteries-claims-scientist.html>

[6] The Higgs particle can disintegrate into particles of dark matter, according to new model

<http://www.chalmers.se/en/departments/fp/news/Pages/The-Higgs-particle-can-disintegrate-into-particles-of-dark-matter,-according-to-new-model.aspx>

[7] Exploring the Higgs boson's dark side

<http://phys.org/news/2015-07-exploring-higgs-boson-dark-side.html>

[8] Recent study predicts that Higgs particles are much heavier than earlier observation

<http://phys.org/news/2016-01-higgs-particles-heavier-earlier.html>

[9] Physicists Have Discovered Evidence Of A Gluino Particle, The Cousin Of The Higgs Boson

<http://www.techtimes.com/articles/117035/20151216/hadron-collider-finds-new-particle-gluino.htm>