

## 1. ABSTRACT

This paper develops the Dark Matter by Gravitation theory, DMbG theory hereafter, in clusters of galaxies in the current cosmologic model  $\Lambda$ CDM of the Universe.

Originally this theory was developed by the author for galaxies, especially using MW and M31 rotation curves. An important results got by the DMbG theory is that the total mass associated to a galactic halo depend on the square root of radius, being its dominion unbounded. Apparently, this result would be absurd because of divergence of the total mass.

As the DE is negligible at galactic scale it is needed to extend the theory to clusters in order to study the capacity of DE to counterbalance to DM. Thanks this property, the DMbG theory finds unexpected theoretical results. In this work it is defined, the total mass as baryonic matter plus DM and the gravitating mass as the addition of the total mass plus the negative mass associated to dark energy.

In clusters it is defined the zero gravity radius ( $R_{ZG}$  hereafter) as the radius needed by the dark energy to counterbalance the total mass. It have been found, that the ratio  $R_{ZG}/R_{VIRIAL} \approx 7.3$  and its Total mass associated at  $R_{ZG}$  is  $\approx 2.7 \cdot M_{VIRIAL}$ . In addition it has been calculated that the sphere with the extended halo radius  $R_E = 1.85 \cdot R_{ZG}$  has a ratio DM density versus DE density equal to  $3/7$  and its total mass associated at  $R_E$  is  $\approx 3.67 \cdot M_{VIRIAL}$ . This works postulates that the factor 3.67 may equilibrate perfectly the strong imbalance between the Local mater density parameter (0.08) versus the current Global matter density one (0.3). Currently this fact is a big conundrum in cosmology, see chapter 7.

Also it has been found that the zero velocity radius,  $R_{ZV}$  hereafter, i.e. the cluster border because of the Hubble flow, is  $\approx 0.6 \cdot R_{ZG}$  and its gravitating mass is  $\approx 1.5 M_{VIR}$

By derivation of gravitating mass function it is calculated that at  $0.49 \cdot R_{ZG}$  this function reaches its maximum whose value is  $\approx 1.57 \cdot M_{VIR}$

Throughout the paper some of these results have been validated with recent data published for the Virgo cluster. As Virgo is the nearest big cluster, it is the perfect benchmark to validate any new theory about DM and DE.

These new theoretical findings offer to scientific community a wide number of tests to validate or reject the theory. The validation of DMbG theory would mean to know the nature of DM that at the present it is an important challenge for the astrophysics science.

1. ABSTRACT .....	1
2. INTRODUCTION.....	2
3. VIRIAL MASS AND VIRIAL RADIUS IN CLUSTER OF GALAXIES .....	4
3.1 CHECKING THE VIRIAL MASS APROXIMATION ON A SAMPLE OF CLUSTERS AND GROUP OF G. 4	
4. VIRIAL THEOREM AS A METHOD TO GET THE DIRECT MASS FORMULA IN CLUSTERS.....	5
4.1 PARAMETER $a^2$ FORMULA DEPENDING ON VIRIAL RADIUS AND VIRIAL MASS .....	5
4.2 PARAMETER $a^2$ FORMULA DEPENDING ON VIRIAL MASS ONLY .....	5
5. DARK MATTER IS COUNTER BALANCED BY DARK ENERGY AT ZERO GRAVITY RADIUS .....	6
5.1 ZERO GRAVITY RADIUS DEPENDING ON PARAMETER $a^2$ FORMULA .....	6
5.2 ZERO GRAVITY RADIUS FORMULA DEPENDING ON VIRIAL MASS .....	7
5.3 ZERO GRAVITY RADIUS VERSUS VIRIAL RADIUS.....	7
5.4 TOTAL MASS ASSOCIATED TO A CLUSTER OF GALAXIES .....	8
5.4.1 TOTAL MASS ASSOCIATED TO THE SPHERE WITH ZERO GRAVITY RADIUS .....	8
5.4.2 TOTAL MASS AT ZERO GRAVITY RADIUS USING THE VIRIAL MASS .....	8
5.5 TOTAL DARK ENERGY AT ZERO GRAVITY RADIUS .....	9
5.6 GRAVITATING MASS FUNCTION.....	9
5.7 CALCULUS FOR THE MAXIMUM OF GRAVITATING MASS .....	10
5.8 DENSITY OF THE TOTAL MASS INTO THE HALO CLUSTER THEOREM.....	10
5.8.1 DARK ENERGY DENSITY INTO THE HALO CLUSTER .....	11
5.9 EXTENDED HALO WHERE THE RATIO TOTAL MASS VERSUS DARK ENERGY IS EQUAL TO 3 / 7.....	11
6. ZERO VELOCITY RADIUS BECAUSE OF THE HUBBLE FLOW.....	12
6.1 GRAVITATIONAL POTENTIAL INTO THE HALO CLUSTER .....	12
6.2 EQUATION FOR ZERO VELOCITY RADIUS .....	12
6.3 ZERO VELOCITY RADIUS FOR VIRGO .....	12
6.4 ZERO VELOCITY RADIUS THEOREM.....	13
6.5 GRAVITATING MASS AT ZERO VELOCITY RADIUS.....	13
7. VALIDATION OF THE THEORY WITH RESULTS PUBLISHED ABOUT VIRGO CLUSTER .....	13
7.1 GRAVITATING MASS ASSOCIATED UP TO THE ESTIMATED ZERO VELOCITY RADIUS .....	14
7.2 GRAVITATING MASS ASSOCIATED UP TO THE TWICE OF VIRIAL RADIUS.....	14
7.3 SOLVING THE CONUNDRUM: LOCAL DENSITY MATTER VERSUS GLOBAL DENSITY MATTER..	15
8. CONCLUDING REMARKS.....	16
9. BIBLIOGRAPHYC REFERENCES .....	16

## 2. INTRODUCTION

The bases of this paper are developed in [1] Abarca, M. 2023, so it is highly recommended to read it to understand the meaning of this paper. The dark matter by gravitation theory, DMbG theory hereafter, is an original theory developed since 2013 through more than 20 papers, although in [1] Abarca, M. 2023 is published the best version as physical as

mathematically. Therefore is not possible to understand this paper if reader have not at least a general knowledge about the DMbG theory.

The hypothesis of DMbG theory is that the DM is generated by the own gravitational field. In order to study purely the DM phenomenon it is needed to consider a radius dominion where it is supposed that baryonic matter is negligible. For example, for MW galaxy the radius must be bigger than 30 kpc and bigger than 40 kpc for M31 galaxy. For galactic clusters, the radius must be bigger than its virial radius.

This hypothesis has two main consequences: the first one is that the law of dark matter generation, in the halo region, has to be the same for all the galaxies and clusters; the second one is that the DM haloes are unlimited so the total dark matter goes up without limit.

The DMbG theory has been developed assuming the hypothesis that DM is a quantum gravitational effect. However, it is possible to remain into the Newtonian framework to develop the theory. In my opinion there are two factors to manage the DM phenomenon with a quite simple theory.

The first one, that it is developed into the halo region, where baryonic matter is negligible. The second one, that the mechanics movements of celestial bodies are very slow regarding velocity of light, which is supposed to be the speed of gravitational bosons. It is known that community of physics is researching a quantum gravitation theory since many years ago, but does not exist yet; however my works in this area support strongly that DM is a quantum gravitation phenomenon.

Use a more simple theory instead the general theory is a typical procedure in physics.

For example the Kirchhoff's laws are the consequence of Maxwell theory for direct current and remain valid for alternating current, introducing complex impedances, on condition that signals must have low frequency.

In [1] Abarca, M. 2023, in the framework of DMbG, is demonstrated mathematically that the total mass (baryonic plus DM) enclosed by a sphere with a specific radius is given by the Direct mass into the galactic halo and that the direct mass formula goes up proportionally to the root square of radius, formula (4.1).

It is well known that DE may be modelled as a constant density of negative mass in the whole space, see [ 5] Chernin,A.D. et all.2013, therefore the total amount of DE grows up with the cubic power of the sphere radius, so it is clear that DE is able to counterbalance the total mass of the clusters, which grows up more slowly. Precisely, the main goal of this paper is to study the relation between both phenomenons in clusters. Namely in cluster haloes.

This paper explores the mutual counterbalance between DM and DE in the framework of DMbG theory and the result got have been fructiferous, with a dozen of new formulas never published before.

In the following paragraphs will be introduced the paper structure:

The newness of the important results got in this paper are due to the possibility to approximate the virial radius to  $R_{200}$  and the virial mass to  $M_{200}$ , the chapter 3 is dedicated to validate this approximation using recent data published for some important clusters such as Virgo or some others.

The chapter 4 is dedicated to extend the direct mass formula to clusters. The direct mass formula (4.1) has only one parameter  $a^2$  whose units are  $m^{5/2}/s^2$ . Using the approximation  $R_{200}$  as virial radius and  $M_{200}$  as virial mass into the direct mass formula it is got the formula  $a^2 = (G \cdot M_{VIR})^{5/6} \cdot (10 \cdot H)^{1/3}$  which is on the basis for some important results got in the following chapters.

The chapter 5 contains three main concepts:

Firstly, it is defined the zero gravity radius,  $R_{ZG}$ , as the sphere radius where the total mass is counterbalanced by the DE. It is found that the ratio  $R_{ZG} / R_{VIRIAL} \approx 7.3$  is universal.

Secondly, it is defined the gravitating mass as the addition of total mass plus the dark energy and the gravitating mass function using a dimensionless parameter  $f = \text{Radius} / R_{ZG}$ . It is found that for any cluster at  $\approx 0.5 \cdot R_{ZG}$  is reached the maximum of gravitating mass and its value is  $M_G (< R_M) \approx 1.57 \cdot M_{VIR}$

Finally it is defined the concept of extended halo ( $R_E$ ) as the spherical region where the ratio  $\frac{Density_{TM}^{SPHERE}(<R_E)}{\rho_{DE}} = 3/7$  i.e. the local ratio of such densities is equal to the current global ratio one, and it is found that  $R_E \approx 1.85 R_{ZG}$

In the sixth chapter it is defined the zero velocity radius,  $R_{ZV}$ , as the sphere radius where the escape velocity is zero because of the Hubble flow. It is demonstrated that the ratio  $R_{ZV} / R_{ZG} \approx 0.602$  is universal.

In the seventh chapter, it is validated the gravitating mass formula into the Virgo cluster for a couple of radius. Namely at 7.3 Mpc and at 3.4 Mpc. The calculus made with the formula of gravitating mass is compared successfully with recent results published in 2020. Also it is validated the theoretical result of  $R_{ZV} / R_{ZG} \approx 0.602$  with result of measures published.

Two of the most important results got in this work are the formula  $M_{TOTAL}(< R_{ZG}) = U \cdot M_{VIR}$  being  $U \approx 2.7$  and the formula  $M_{TOTAL}(< R_E) \approx 3.67 \cdot M_{VIR}$  associated to zero gravity radius and the extended radius respectively.

Thanks these formulas this work suggest the possibility to solve the current discrepancy between the local parameter of matter density,  $\Omega_m^{local} = 0.08$ , see [6] Karachentsev et al. 2014, and the current global one  $\Omega_m^{global} = 0.3$ . This discrepancy is an open problem for the current cosmology.

Finally the chapter 8 is devoted to the concluding remarks.

The reader can consult the paper [8] Abarca, M, 2024, which is an extension of this paper, 24 pages, where the theory is validated through the Local Group and the Coma cluster data published as well.

### 3. VIRIAL MASS AND VIRIAL RADIUS IN CLUSTER OF GALAXIES

In cluster, it is a good estimation about virial radius and virial mass to consider  $R_{vir} = R_{200}$  and  $M_{vir} = M_{200}$ . Where  $R_{200}$  is the radius of a sphere whose mean density is 200 times bigger than the critic density of Universe

$$\rho_c = \frac{3H^2}{8\pi G} \quad (3.1) \text{ and } M_{200} \text{ is the total mass enclosed by the radius } R_{200}.$$

Considering the spherical volume formula, it is right to get the following relation between both concepts

$$R_{VIR}^3 \approx R_{200}^3 = \frac{G \cdot M_{200}}{100 \cdot H^2} \quad (3.2) \text{ or } M_{VIR} \approx M_{200} = \frac{100 H^2 R_{200}^3}{G} \quad (3.3)$$

#### 3.1 CHECKING THE VIRIAL MASS APROXIMATION ON A SAMPLE OF CLUSTERS AND GROUP OF G.

Data [4] R.Ragusa et al.2022		Table 1		
Group of galaxies G. Or Clusters C.	Virial Radius	Virial Mass	Mass calculated	Relative diff for M
Name	Mpc	$\times 10^{13} M_{\odot}$	$\times 10^{13} M_{\odot}$	%
Antlia C.	1,28	26,3	2,39E+01	-9,21E+00
NGC596/584 G.	0,5	1,55	1,42E+00	-8,18E+00
NGC 3268 G.	0,9	8,99	8,30E+00	-7,67E+00
NGC 4365 Virgo SubG.	0,32	0,4	3,73E-01	-6,73E+00
NGC 4636 Virgo SubG.	0,63	3,02	2,85E+00	-5,73E+00
NGC 4697 Virgo Sub G.	1,29	26,9	2,44E+01	-9,14E+00
NGC 5846 G.	1,1	16,6	1,52E+01	-8,71E+00
NGC 6868 G.	0,6	2,69	2,46E+00	-8,57E+00

Data beside in green have been taken from [4] R. Ragusa et al. 2022 and using the formula  $M_{200} = \frac{100 H^2 R_{200}^3}{G}$  it is calculated its mass associated for each radius. The yellow column shows the relative difference for masses, always under 10 %.

As the Virgo cluster is the nearest between the big clusters it is crucial to check the approximation for virial mass and radius with its data.

According [7] Olga Kashibadze, I. Karachentsev 2020, see pag 9,  $R_{vir} = 1.7$  Mpc and  $M_{vir} = (6.3 \pm 0.9) \cdot 10^{14} M_{\odot}$ . Using formula (3.3) with  $R_{200} = 1.7$  Mpc it is got  $M_{200} = 5.59 \cdot 10^{14} M_{\odot}$  which match with mass published if it is considered the range of errors.

<b>Table 2</b>	Virial Radius	Virial mass	Calculated $M_{200}$	Mass Relative diff.
Cluster	Mpc	$\times 10^{14} M_{\odot}$	$\times 10^{14} M_{\odot}$	%
Virgo [13]Kashibadze 2020	1.7	6.3±0.9	5.59	11

In conclusion  $R_{200}$  and  $M_{200}$  are a very good estimation for Virial radius and Virial mass for group of galaxies and cluster of galaxies, when they are in dynamical equilibrium.

#### 4. VIRIAL THEOREM AS A METHOD TO GET THE DIRECT MASS FORMULA IN CLUSTERS

In chapter 9, of paper [1]Abarca,M.2023 was demonstrated that direct formula  $M_{TOTAL}(< r) = \frac{a^2 \cdot \sqrt{r}}{G}$  (4.1) is the most suitable formula to calculate the total mass (baryonic and DM) depending on radius in the galactic halo.

##### 4.1 PARAMETER $a^2$ FORMULA DEPENDING ON VIRIAL RADIUS AND VIRIAL MASS

Due to the fact that the Direct mass formula has one parameter only, is enough to know the mass associated to a specific radius to be able to calculate parameter  $a^2$ . That is the situation when it is known the virial mass and the virial radius for a cluster of galaxies.

If it is considered that the virial radius is the border of halo cluster where galaxies are in dynamical equilibrium and at the same time is negligible the amount of Baryonic matter outside the sphere with such radius, then according DMbG theory is possible to do an equation between  $M_{VIRIAL}(< R_{VIRIAL}) = M_{DIRECT}(< R_{VIRIAL})$  (4.2.1) i.e.

$$M_{VIRIAL} \equiv M_{VIRIAL}(< R_{VIRIAL}) = \frac{a^2 \cdot \sqrt{R_{VIRIAL}}}{G} \quad (4.2.2) \text{ and clearing up } a^2 = \frac{G \cdot M_{VIRIAL}}{\sqrt{R_{VIRIAL}}} \quad (4.3), \text{ this formula will be}$$

called parameter  $a^2$  ( $M_{VIR}$ ,  $R_{VIR}$ ) because depend on both measures.

##### 4.2 PARAMETER $a^2$ FORMULA DEPENDING ON VIRIAL MASS ONLY

In chapter 3 was got this formula  $R_{VIR}^3 = \frac{G \cdot M_{VIR}}{100 \cdot H^2}$  (3.2) as a good approximation between virial mass and virial radius .

So using that formula and by substitution of virial radius in  $a^2 = \frac{G \cdot M_{VIRIAL}}{\sqrt{R_{VIRIAL}}}$  (4.3) it is right to get parameter  $a^2$

depending on  $M_{VIR}$  only  $a^2 = (G \cdot M_{VIR})^{5/6} \cdot (10 \cdot H)^{1/3}$  (4.4) This formula will be called parameter  $a^2$  ( $M_{VIR}$ ) as depend on  $M_{VIR}$  only.

With the virial data for Virgo cluster will be calculated its parameter  $a^2$  with the formula (4.3) i.e.  $a^2$  ( $M_{VIR}$ ,  $R_{VIR}$ ) and with formula (4.4) i.e.  $a^2$  ( $M_{VIR}$ )

The last formula is an approximation of the previous formula as it is supposed that  $R_{VIR} = R_{200}$ . Below are calculated both formulas and fortunately its relative difference is negligible.

<b>Table 3</b>	Virial Radius	Virial mass	Parameter $a^2 (M_{VIR}, R_{VIR})$	Parameter $a^2 (M_{VIR})$	Relative diff.
Clusters	Mpc	$\cdot 10^{14} M_{\odot}$	I.S. units $m^{5/2} / s^2$	I.S. units $m^{5/2} / s^2$	%
Virgo	1.7	6.3±0.9	3.6527E23	3.581E23	2

Green data come from [7] Olga Kashibadze.2020

## 5. DARK MATTER IS COUNTER BALANCED BY DARK ENERGY AT ZERO GRAVITY RADIUS

The basic concepts about DE on the current cosmology can be studied in [5] Chernin,A.D.

As currently there is a tension regarding the experimental value of Hubble constant, in this paper will be used  $H = 70$  Km/s/Mpc and  $\Omega_{DE} = 0.7$  as the fraction of Universal density of DE.

### 5.1 ZERO GRAVITY RADIUS DEPENDING ON PARAMETER $a^2$ FORMULA

According [ 5] Chernin,A.D. in the current cosmologic model  $\Lambda$ CDM , dark energy has an effect equivalent to antigravity i.e. the mass associated to dark energy is negative and the dark energy have a constant density for all the

Universe equal to  $\rho_{DE} = \rho_C \cdot \Omega_{DE} = -6.444 \cdot 10^{-27} \text{ kg/m}^3$  being  $\Omega_{DE} = 0.7$  and  $\rho_C = \frac{3H^2}{8\pi G} = 9.205E-27 \text{ kg/m}^3$

the critic density of the Universe.

As DE density is constant, the total DE mass is proportional to Radius with power 3, whereas DM mass grows with radius power 0.5 so it is right to get a radius where DM is counter balanced by DE.

According [ 5] Chernin, A.D.The mass associated to DE is  $M_{DE}(< R) = -\frac{\rho_{DE} 8\pi R^3}{3}$ , (5.1) , or equivalently

$M_{DE}(< R) = -\rho_{DE} \frac{8\pi R^3}{3} = \frac{-H^2 \cdot \Omega_{DE}}{G} \cdot R^3$  (5.2). Notice that the author multiplies by two the volume of a sphere by reasons explained in his work.

[ 5] Chernin defines gravitating mass  $M_G(< R) = M_{DE}(< R) + M_{TOTAL}(< R)$  (5.3), where  $M_{TOTAL}$  is baryonic plus dark matter mass, and defines  $R_{ZG}$  , Radius at zero Gravity as the radius where  $M_{DE}(< R_{ZG}) + M_{TOTAL}(< R_{ZG}) = 0$  . i.e. where the gravitating mass is zero.

According DMbG theory this definition leads to equation  $M_{TOTAL}(< R_{ZG}) = \rho_{DE} \frac{8\pi R_{ZG}^3}{3}$ , (5.4).

Using (4.1) formula  $M_{TOTAL}(< R_{ZG}) = \frac{a^2 \cdot \sqrt{R_{ZG}}}{G}$  the expression (5.4) leads to  $\frac{a^2 \cdot \sqrt{R_{ZG}}}{G} = \rho_{DE} \frac{8\pi \cdot R_{ZG}^3}{3}$ , (5.5),

where it is possible to clear up  $R_{ZG} = \left[ \frac{3a^2}{8\pi G \rho_{DE}} \right]^{2/5}$  (5.6) and as  $\rho_{DE} = \frac{3 \cdot H^2}{8\pi G} \Omega_{DE}$  (5.7) then by substitution

$$R_{ZG} = \left[ \frac{a^2}{H^2 \cdot \Omega_{DE}} \right]^{2/5} \quad (5.8) \text{ This formula will be called } R_{ZG} \text{ (parameter } a^2 \text{).}$$

As the radius  $R_{ZG}$  is the distance to cluster centre where is zero the gravitating mass, it is right to consider  $R_{ZG}$  as the halo radius and its sphere defined as the halo cluster.

**5.2 ZERO GRAVITY RADIUS FORMULA DEPENDING ON VIRIAL MASS**

In previous chapter was got the value for  $a^2 = (G \cdot M_{VIR})^{5/6} \cdot (10 \cdot H)^{1/3}$  ,(4.4), depending on  $M_{VIR}$  , so by

substitution in (5.8) it is right to get  $R_{ZG} = \frac{(G \cdot M_{VIR})^{1/3} \cdot \sqrt[15]{100}}{H^{2/3} \Omega_{DE}^{2/5}}$  (5.9)

Below are calculated  $R_{ZG}$  by two ways: Formulas (5.8) and (5.9) . Both calculus are mathematically equivalents. See in table below how the grey values match perfectly.

<b>Table 4</b>	Virial mass	Parameter $a^2 (M_{VIR})$	$R_{ZG}$ (parameter $a^2$ )	$R_{ZG} (M_{VIR})$	Relative diff.
Clusters	$\cdot 10^{14} M_{\odot}$	I.S. units $m^{5/2} / s^2$	Mpc	Mpc	%
Virgo	6.3±0.9	3.581E23	12.871	12.871	0

Green data come from [7] Olga Kashibadze.2020. With this important cluster of galaxies, it has been illustrated how the total mass, calculated by  $M_{TOTAL}(< r) = \frac{a^2 \cdot \sqrt{r}}{G}$  , is counter balanced by dark energy at mega parsecs scale, and precisely this Radius at zero gravity determines the region size where the cluster has gravitational influence.

**5.3 ZERO GRAVITY RADIUS VERSUS VIRIAL RADIUS**

From formula (3.2)  $R_{VIR}^3 \approx R_{200}^3 = \frac{G \cdot M_{200}}{100 \cdot H^2} \approx \frac{G \cdot M_{VIR}}{100 \cdot H^2}$  it is right to get  $R_{VIR} = \left(\frac{G \cdot M_{VIR}}{100 \cdot H^2}\right)^{1/3}$  . In previous epigraph

was got  $R_{ZG} = \frac{(G \cdot M_{VIR})^{1/3} \cdot \sqrt[15]{100}}{H^{2/3} \Omega_{DE}^{2/5}}$  (see 5.9). So it is right to get the ratio  $\frac{R_{ZG}}{R_{VIR}} = \left(\frac{100}{\Omega_{DE}}\right)^{2/5} \approx 7.277$  (5.10) which is

universal as it does not depend on virial mass associated to a specific cluster.

<b>Table 5</b>	Virial Radius	Virial Mass	Zero Gravity R.	Ratio
Celestial Body	Mpc	1E13 $M_{\odot}$	Mpc	$R_{ZG} / R_{VIR}$
Antlia cluster	1,28	26,3	9,62E+00	7,52E+00
NGC596/584	0,5	1,55	3,74E+00	7,49E+00
NGC 3268	0,9	8,99	6,73E+00	7,47E+00
NGC 4365	0,32	0,4	2,38E+00	7,45E+00
NGC 4636	0,63	3,02	4,68E+00	7,42E+00
NGC 4697	1,29	26,9	9,69E+00	7,51E+00
NGC 5846	1,1	16,6	8,25E+00	7,50E+00
NGC 6868	0,6	2,69	4,50E+00	7,50E+00

Columns in blue come from [4] R. Ragusa et al.2022

The second and third columns show the virial radius and the virial mass for each cluster.

The column in green shows the  $R_{ZG}$  using formula (5.9)

The column in pink is the ratio of radius.

It is clear that the ratio  $R_{ZG}/R_{VIR}$  got in this sample of celestial bodies match very well with the theoretical ratio formula (5.10)

<b>Table 6</b>	VirialRadius	Zero Grav R	Ratio
Cluster	Mpc	Mpc	$R_{ZG} / R_{VIR}$
Virgo	1.7	12.871	7.57

For Virial radius see table 3, for Zero gravity rad. see table 4

The data of Virgo cluster back the theoretical ratio formula as well.

## 5.4 TOTAL MASS ASSOCIATED TO A CLUSTER OF GALAXIES

### 5.4.1 TOTAL MASS ASSOCIATED TO THE SPHERE WITH ZERO GRAVITY RADIUS

In epigraph 5.1 was shown how this equation  $M_{TOTAL}(< R_{ZG}) = \rho_{DE} \cdot \frac{8\pi \cdot R_{ZG}^3}{3}$  (see 5.4) is used to define  $R_{ZG}$ . By simplification it is got  $M_{TOTAL}(< R_{ZG}) = \frac{H^2 \cdot \Omega_{DE}}{G} \cdot R_{ZG}^3 = M \cdot R_{ZG}^3$  (5.11) being  $M = 5.3984 \cdot 10^{-26}$  (I.S.) So that is the total mass formula, using the equation between total mass and dark energy at zero gravity radius.

In table below is shown calculus of total mass for Virgo.

<b>Table 7</b>	Radius ZG	$M_{TOTAL}(< R_{ZG}) = M \cdot R_{ZG}^3$
Clusters	Mpc	$M_{\odot}$
Virgo	12.871	1.699945E15

### 5.4.2 TOTAL MASS AT ZERO GRAVITY RADIUS USING THE VIRIAL MASS

Using rightly the direct mass formula (see 4.1) at  $R_{ZG}$  it is got  $M_{TOTAL}(< R_{ZG}) = \frac{a^2 \cdot \sqrt{R_{ZG}}}{G}$  (5.12)

As in epigraph 4.2 was got  $a^2 = (G \cdot M_{VIR})^{5/6} \cdot (10 \cdot H)^{1/3}$  and in epigraph 5.2 was got  $R_{ZG} = \frac{(G \cdot M_{VIR})^{1/3} \cdot \sqrt[5]{100}}{H^{2/3} \Omega_{DE}^{2/5}}$

by substitution in  $M_{TOTAL}(< R_{ZG}) = \frac{a^2 \cdot \sqrt{R_{ZG}}}{G}$  (see 5.12) it is got  $M_{TOTAL}(< R_{ZG}) = \frac{\sqrt[5]{100}}{\Omega_{DE}^{1/5}} \cdot M_{VIR}$  (5.13) calling

$U = \frac{\sqrt[5]{100}}{\Omega_{DE}^{1/5}}$  (5.14) then  $M_{TOTAL}(< R_{ZG}) = U \cdot M_{VIR}$  (5.15) being  $U \approx 2.6976$ .

So may be stated that according Dark matter by gravitation theory, the total mass (baryonic plus DM) enclosed by the sphere with radius  $R_{ZG}$  is equivalent to 2.7 times the Virial Mass.

Below are compared the masses for Virgo using the formula (5.15) and the previous one (5.11). It is clear that both are mathematically equivalents.

<b>Table 8</b>	Virial mass	$M_{TOTAL}(< R_{ZG}) = U \cdot M_{VIR}$	$M_{TOTAL}(< R_{ZG}) = M \cdot R_{ZG}^3$	Relatif diff.
Clusters	$\cdot 10^{14} M_{\odot}$	$M_{\odot}$	$M_{\odot}$	%
Virgo	6.3±0.9	1.6994E15	1.6994E15	0

Green data come from [7] Olga Kashibadze.2020.

By the direct formula for total mass, may be calculated the total mass associated to a radius, which is a fraction of  $R_{ZG}$ , with the following property: if  $(R = f \cdot R_{ZG})$  then

$$M_{TOTAL}(< R) = M_{TOTAL}(< f \cdot R_{ZG}) = \sqrt{f \cdot R_{ZG}} \cdot \frac{a^2}{G} = \sqrt{f} \cdot U \cdot M_{VIR} \quad (5.16)$$

because  $M_{TOTAL}(< R_{ZG}) = U \cdot M_{VIR}$  (See 5.15)

For example, using the data for Virgo cluster  $R_{ZG} = 12.87$  and  $R_{VIR} = R_{200} = 1.7687$  Mpc, so  $R_{VIR} / R_{ZG} = f = 0.137428$  and by the formula (5.16)  $M_{TOTAL}(< R_{VIR}) = M_{TOTAL}(< f \cdot R_{ZG}) = \sqrt{f} \cdot U \cdot M_{VIR} \approx 0.99999 M_{VIR}$ .



### 5.5 TOTAL DARK ENERGY AT ZERO GRAVITY RADIUS

Using equation (5.2) at  $R_{ZG}$  it is got  $M_{DE}(< R_{ZG}) = -\frac{\rho_{DE} 8\pi R_{ZG}^3}{3} = \frac{-H^2 \cdot \Omega_{DE}}{G} \cdot R_{ZG}^3$  (5.17) that it is just the opposite

value to  $M_{TOTAL}(< R_{ZG}) = \frac{\sqrt[5]{100}}{\Omega_{DE}^{1/5}} \cdot M_{VIR}$  (see 5.13) because the total gravitating mass enclosed into the sphere of zero

gravity radius is zero by definition .  $M_G(< R_{ZG}) = M_{TOTAL}(< R_{ZG}) + M_{DE}(< R_{ZG}) = 0$  (5.18). See epigraph 5.1

Therefore  $M_{DE}(< R_{ZG}) = -\frac{\sqrt[5]{100}}{\Omega_{DE}^{1/5}} \cdot M_{VIR} = -U \cdot M_{VIR}$  (See 5.15), being  $U \approx 2.6976$  and joining both formulas it is got

$$M_{DE}(< R_{ZG}) = -\frac{\rho_{DE} 8\pi R_{ZG}^3}{3} = \frac{-H^2 \cdot \Omega_{DE}}{G} \cdot R_{ZG}^3 = -U \cdot M_{VIR} \quad (5.19)$$

This formula may be used to calculate the mass associated to DE if the radius is a fraction of  $R_{ZG}$  : if  $(R = f \cdot R_{ZG})$

$$\text{then (See 5.1) } M_{DE}(< R) = -\frac{\rho_{DE} 8\pi R^3}{3} = M_{DE}(< f \cdot R_{ZG}) = \frac{-f^3 \cdot H^2 \cdot \Omega_{DE}}{G} \cdot R_{ZG}^3 = -f^3 \cdot U \cdot M_{VIR} \quad (5.20)$$

For example using the Virgo data  $R_{ZG} = 12.871$  Mpc and  $R_{VIR} = R_{200} = 1.7687$  Mpc, so  $R_{VIR} / R_{ZG} = f = 0.137428$  and

$M_{DE}(< R_{VIR}) = M_{DE}(< f \cdot R_{ZG}) = -f^3 \cdot U \cdot M_{VIR} = -0.007 \cdot M_{VIR}$  So at  $R_{VIR}$  the DE is negligible. However at  $R_{ZG}$  is able to cancel the total mass.

### 5.6 GRAVITATING MASS FUNCTION

In the epigraph 5.1 was defined the gravitating mass  $M_G = M_{DE} + M_{TOTAL}$ , where  $M_{TOTAL}$  is baryonic plus dark matter mass and  $M_{DE}$  is the negative mass associated to DE.

The best way to calculate the gravitating mass is using the formulas got in epigraph 5.4 and 5.5 where are calculated the both types of masses associated to a radius, which is a fraction of  $R_{ZG}$ , i.e.  $R = f \cdot R_{ZG}$ , these formulas are:

$$M_{TOTAL}(< f \cdot R_{ZG}) = \sqrt{f} \cdot U \cdot M_{VIR} \quad (\text{See 5.16}) \quad \text{and} \quad M_{DE}(< f \cdot R_{ZG}) = \frac{-f^3 \cdot H^2 \cdot \Omega_{DE}}{G} \cdot R_{ZG}^3 = -f^3 \cdot U \cdot M_{VIR} \quad (\text{See 5.20})$$


and joining both formulas it is got  $M_G(< R) = M_G(< f \cdot R_{ZG}) = [\sqrt{f} - f^3] \cdot U \cdot M_{VIR}$  (5.21) where  $U \approx 2.6976$ .

This way the gravitating mass depends on the dimensionless factor f.

### DOMINION OF GRAVITATING MASS FUNCTION

As the gravitating mass is defined into the halo cluster, its dominion begins at  $R_{VIR}$  so the gravitating mass depending on f begins at  $R_{VIR}/R_{ZG}$ . By (5.10) formula  $R_{ZG}/R_{VIR} = 7.277$  then  $R_{VIR}/R_{ZG} = 1/7.277$  therefore the dominion of the gravitating mass function depending on  $f = R/R_{ZG}$  begins at 0.13732

In this epigraph this function will be studied up to  $f = 1$  i.e. when the radius reaches the zero gravity radius.



Above is represented the ratio *gravitating mass / virial mass* versus the ratio  $R/R_{ZG}$  in its dominion and close to 0.5 there is the function maximum that will be calculated below.

### 5.7 CALCULUS FOR THE MAXIMUM OF GRAVITATING MASS

It is clear that such function will have a maximum, that it is found easily by derivation,  $f = \frac{1}{\sqrt[5]{36}}$ , being  $R_M = \frac{1}{\sqrt[5]{36}} \cdot R_{ZG} \approx 0.48836 \cdot R_{ZG}$ . (5.22)

By substitution of this value  $f = \frac{1}{\sqrt[5]{36}} \approx 0.48836$  into the mass gravitating formula

$M_G(< R_M) = M_G(< f \cdot R_{ZG}) = [\sqrt{f} - f^3] \cdot U \cdot M_{VIR}$  being  $U \approx 2.6976$  it is got

$$M_G^{MAXIMUM}(< R_M) = 1.57 \cdot M_{VIR} \quad (5.23).$$

So may be stated one important result, got by the Dark Matter by Gravitation theory:

***For any cluster of galaxies at a half of  $R_{ZG}$ , it is reached the maximum of gravitating mass which is  $1.57 M_{VIR}$ .***

### 5.8 DENSITY OF THE TOTAL MASS INTO THE HALO CLUSTER THEOREM

Assuming the hypothesis that  $R_{vir} = R_{200}$  and  $M_{vir} = M_{200}$ , i.e. the virial sphere  $Density_{VIR} = \frac{M_{vir}}{Vol_{VIR}} = 200 \rho_c$  being

$$\rho_c = \frac{3H^2}{8\pi G} \text{ the critical density of Universe.}$$

Then into the halo cluster sphere  $Density_{TOTAL MASS}(< R_{ZG}) = 1.4 \cdot \rho_c$  (5.26). Being  $R_{ZG}$  the halo radius.

**Proof.**

$$Density_{TOTAL MASS}^{CLUSTER} = \frac{M_{TOTAL}(< R_{ZG})}{Vol_{CL}(< R_{ZG})} = \frac{M_{VIR} \cdot U}{Vol_{vir} \cdot \left(\frac{R_{ZG}}{R_{VIR}}\right)^3} = \frac{M_{VIR} \cdot U}{Vol_{vir} \cdot U^6} = \frac{200 \cdot \rho_c}{U^5} = \frac{200 \cdot \rho_c \cdot \Omega_{DE}}{100} = 1.4 \cdot \rho_c$$

In that chained equalities has been used two main properties before got: (5.15) and (5.10)

### 5.8.1 DARK ENERGY DENSITY INTO THE HALO CLUSTER

As into the current  $\Lambda$ CDM model the DE density has a constant value,  $\rho_{DE} = 0.7\rho_C$ , it is not necessary to demonstrate such property. Nevertheless, in this epigraph it will be checked such value.

So, according [ 5] Chernin, A.D. The mass associated to DE is  $M_{DE}(< R) = -\frac{\rho_{DE} 8\pi R^3}{3}$ , (5.1).

By other side, according DMbG theory  $M_{DE}(< R_{ZG}) = -\frac{\rho_{DE} 8\pi R_{ZG}^3}{3} = \frac{-H^2 \cdot \Omega_{DE} \cdot R_{ZG}^3}{G} = -U \cdot M_{VIR}$  (5.19)

So, at a specific radius such formula may be rewritten  $M_{DE}(< R) = -2 \cdot \rho_{DE} \cdot Volume$  (5.27) and by similar reasons that previous epigraph may be got that

$$Density_{DAR ENERGY}^{CLUSTER} = \frac{M_{DE}(< R_{ZG})}{2 \cdot Vol_{CL}(< R_{ZG})} = \frac{-U \cdot M_{VIR}}{2 \cdot Vol_{CL}(< R_{ZG})} = -0.7 \cdot \rho_C = \rho_{DE} \quad (5.28)$$

### COROLARIUS

The ratio density of total mass versus density of DE into the halo cluster is

$$\frac{Density_{TM}^{HALO}(< R_{ZG})}{\rho_{DE}} = 2 \quad (5.29)$$

### 5.9 EXTENDED HALO WHERE THE RATIO TOTAL MASS VERSUS DARK ENERGY IS EQUAL TO 3 / 7

At the present, it is accepted that  $\frac{\Omega_M}{\Omega_{DE}} = 3/7$  as a global average in the current Universe, so it is worth to calculate an extension of halo cluster,  $R_E$ , where it is reached such ratio. Below, it will be calculated the radius of a sphere that verify  $\frac{Density_{TM}^{SPHERE}(< R_E)}{\rho_{DE}} = 3/7$  (5.30)

As  $M_{TM}(< R) = Density_{TM}^{SPHERE} \cdot Volume$  and by (5.27)  $M_{DE}(< R) = 2 \cdot \rho_{DE} \cdot Volume$  then by substitution into (5.30) it is got  $\frac{Density_{TM}^{SPHERE}}{\rho_{DE}} = \frac{2 \cdot M_{TM}(< R)}{M_{DE}(< R)} = 3/7$  (5.31)

Using (5.10) and (5.20) it is right to get the second one equality below.

$$\frac{Density_{TM}^{SPHERE}(< R)}{\rho_{DE}} = 2 \cdot \frac{M_T(< R)}{M_{DE}(< R)} = \frac{2 \cdot \sqrt{f}}{f^3} \quad (5.32)$$

Using the condition (5.30) it is got  $\frac{2 \cdot \sqrt{f}}{f^3} = \frac{3}{7}$  whose solution is  $f = \left(\frac{14}{3}\right)^{2/5} \approx 1.85$  (5.33)

Therefore the radius of extended halo searched is  $R_E \approx 1.85 \cdot R_{ZG}$  (5.34)

In order to validate such calculus is enough to check that the Density of total mass with  $R_E$  radius is  $0.3 \cdot \rho_C$

$$Density_{TOTAL MASS}^{EXT. HALO}(< R_E) = \frac{M_{VIR} \cdot U \cdot \sqrt{1.85}}{Vol_{vir} \cdot \left(\frac{R_{ZG}}{R_{VIR}}\right)^3 \cdot 1.85^3} = \frac{200 \cdot \rho_C \cdot \Omega_{DE}}{100 \cdot 1.85^{5/2}} = \frac{1.4}{1.85^{5/2}} \cdot \rho_C = 0.3 \cdot \rho_C \quad (5.35) \text{ as it was expected.}$$

For example for the Virgo cluster  $R_E = 1.85 \cdot 12.9 \text{ Mpc} = 23.9 \text{ Mpc}$

## 6. ZERO VELOCITY RADIUS BECAUSE OF THE HUBBLE FLOW

It is defined the zero velocity radius as the distance to the cluster centre, where the escape velocity from gravitation field is equal to Hubble flow velocity. i.e.  $V_E = V_{HF}$  (6.1).

From classical dynamic it is taken the formula  $\frac{V_E^2}{2} = -V(R)$  (6.2) i.e. the kinetic energy associated to escape velocity compensates the potential energy getting zero as total energy ad infinitum.

It is not possible to use the classical escape velocity formula  $V_E^2 = \frac{2GM}{R}$  (6.3) because of two reasons:

1° The gravitational potential in DMbG theory is different to  $V = -\frac{GM}{R}$

2° The border of the halo cluster is  $R_{ZG}$  where the gravitating mass is zero.

### 6.1 GRAVITATIONAL POTENTIAL INTO THE HALO CLUSTER

As the gravitating mass is zero at  $R_{ZG}$  the line integral for potential goes up to  $R_{ZG}$ , so the formula is

$$V = \int_R^{R_{ZG}} \left[ \frac{-GM_G(<r)}{r^2} \right] \cdot dr \quad (6.4)$$

The gravitating mass  $M_G(<r) = M_G(<f \cdot R_{ZG}) = [\sqrt{f} - f^3] \cdot U \cdot M_{VIR}$  (5.21) will be used to do the integral, doing some little changes.

As  $f = r/R_{ZG}$  and calling  $K = U \cdot M_{VIR}$ , by substitution in (5.21) it is got  $M_G(<r) = K \cdot \left[ \frac{\sqrt{r}}{\sqrt{R_{ZG}}} - \frac{r^3}{R_{ZG}^3} \right]$  (6.5) that by

substitution in (6.4) results  $V(R) = -GK \int_R^{R_{ZG}} \left[ \frac{r^{-3/2}}{\sqrt{R_{ZG}}} - \frac{r}{R_{ZG}^3} \right] \cdot dr$  (6.6) whose result is

$$V(r) = \frac{5GK}{2 \cdot R_{ZG}} - GK \cdot \left[ \frac{2}{\sqrt{r} \cdot \sqrt{R_{ZG}}} + \frac{r^2}{2 \cdot R_{ZG}^3} \right] \quad (6.7)$$

where its dominion ranges from  $R_{VIR}$  up to  $R_{ZG}$ .

Notice that  $V(R_{ZG}) = 0$  and  $V(r)$  is negative inside its dominion.

### 6.2 EQUATION FOR ZERO VELOCITY RADIUS

In this epigraph will be developed the equation  $V_E(r) = V_{HF}(r)$  (6.1) to calculate the  $R_{ZV}$

From (6.2)  $V_E^2 = -2 \cdot V(R)$  So  $-2 \cdot V(R) = V_{HF}^2 = H^2 \cdot R^2$  (6.8) and by substitution of potential formula (6.7) it is got  $\frac{-5GK}{R_{ZG}} + 2GK \cdot \left[ \frac{2}{\sqrt{r} \cdot \sqrt{R_{ZG}}} + \frac{r^2}{2 \cdot R_{ZG}^3} \right] = H^2 \cdot r^2$  (6.9) reorganising that equation it is got:

$$\frac{4GK}{\sqrt{R_{ZG}}} \cdot \frac{1}{\sqrt{r}} + \left[ \frac{GK}{R_{ZG}^3} - H^2 \right] \cdot r^2 = \frac{5GK}{R_{ZG}} \quad (6.10)$$

and multiplying that equation by this factor  $\frac{R_{ZG}}{GK}$  it is got a better expression

$$4\sqrt{R_{ZG}} \cdot \frac{1}{\sqrt{r}} + \left[ \frac{1}{R_{ZG}^2} - \frac{H^2 \cdot R_{ZG}}{GK} \right] \cdot r^2 = 5 \quad (6.11)$$

which is the equation for zero velocity radius.

That equation is not possible to solve with algebraic methods, but is quite easy to solve numerically for specific data.

### 6.3 ZERO VELOCITY RADIUS FOR VIRGO

Green data come from [7] Olga Kashibadze.2020.

Table 9	Virial Radius	Virial mass	$R_{ZG}$
Cluster	Mpc	$\cdot 10^{14} M_{\odot}$	
Virgo	1.7	6.3±0.9	12.87 Mpc

Solving the conundrum of Dark matter and Dark energy in galaxy clusters

$$4\sqrt{R_{ZG}} \cdot \frac{1}{\sqrt{r}} + \left[ \frac{1}{R_{ZG}^2} - \frac{H^2 \cdot R_{ZG}}{GK} \right] \cdot r^2 = 5 \quad (6.11) \text{ As } U \approx 2.7 \text{ and } K = U \cdot M_{VIR} \text{ then } GK = 2.259E+35 \text{ (I.S. units),}$$

$$\frac{H^2 \cdot R_{ZG}}{GK} = 9.047E-48 \text{ m}^{-2}, \frac{1}{R_{ZG}^2} = 6.3397E-48 \text{ m}^{-2} \text{ and } 4\sqrt{R_{ZG}} = 2.5208E+12 \text{ m}^{1/2}$$

The equation (6.11) is easy to be solved numerically. Using  $f = \text{Radius} / R_{ZG}$  it is got that  $f = 0.602$  is a very good approximation for the solution. Therefore the  $R_{ZV} = f \cdot R_{ZG} = 7.75 \text{ Mpc}$

#### 6.4 ZERO VELOCITY RADIUS THEOREM

Considering (6.11) as the equation for the  $R_{ZV}$  then according DMbG theory the ratio  $R_{ZV} / R_{ZG}$  is universal and its value is  $R_{ZV} / R_{ZG} \approx 0.602016$

**Proof**

The expression (6.11) is  $4\sqrt{R_{ZG}} \cdot \frac{1}{\sqrt{r}} + \left[ \frac{1}{R_{ZG}^2} - \frac{H^2 \cdot R_{ZG}}{GK} \right] \cdot r^2 = 5$  where  $K = U \cdot M_{VIR}$  being  $U = \frac{\sqrt[5]{100}}{\Omega_{DE}^{1/5}}$  (5.14) and

$R_{ZG} = \frac{(G \cdot M_{VIR})^{1/3} \cdot \sqrt[15]{100}}{H^{2/3} \Omega_{DE}^{2/5}}$  (5.9) then using the values for  $K$ ,  $U$  and  $R_{ZG}$ , doing some algebraic substitutions and

transformations it is not difficult to get the following equation.  $4\sqrt{R_{ZG}} \cdot \frac{1}{\sqrt{r}} + \frac{1}{R_{ZG}^2} \left[ \frac{\Omega_{DE} - 1}{\Omega_{DE}} \right] \cdot r^2 = 5$  (6.12) and

defining  $f = r / R_{ZG}$  the equation (6.12) becomes

$$4 \cdot \frac{1}{\sqrt{f}} + \left[ \frac{\Omega_{DE} - 1}{\Omega_{DE}} \right] \cdot f^2 = 5 \quad (6.13) \text{ that considering } \Omega_{DE} = 0.7 \text{ becomes:}$$

$$4 \cdot \frac{1}{\sqrt{f}} + \left[ \frac{-3}{7} \right] \cdot f^2 = 5 \quad (6.14) \text{ By elementary algebraic operations it is got this equivalent equation:}$$

$$9 \cdot f^5 + 210 \cdot f^3 + 1225 \cdot f - 784 = 0$$

Thanks Wolfram alpha software, this is its only real solution  $f \approx 0.602016$

By this new formulation of the equation for  $R_{ZV}$  calculus, it has been demonstrated that the ratio  $f = R_{ZV} / R_{ZG}$  is universal and depends on  $\Omega_{DE}$  solely.

#### 6.5 GRAVITATING MASS AT ZERO VELOCITY RADIUS

Using a similar procedure used to calculate the maximum of the gravitating mass, see epigraph 5.7, being  $R_{ZV} = f \cdot R_{ZG}$  where  $f \approx 0.602$  by substitution at formula  $M_G(< R_{ZV}) = M_G(< f \cdot R_{ZG}) = \left[ \sqrt{f} - f^3 \right] \cdot U \cdot M_{VIR}$  (5.21)

where  $U = \frac{100^{1/5}}{\Omega_{DE}^{1/5}} \approx 2.6976$  it is right to get that  $M_G(< R_{ZV}) \approx 1.5 \cdot M_{VIR}$  (6.12)

### 7. VALIDATION OF THE THEORY WITH RESULTS PUBLISHED ABOUT VIRGO CLUSTER

In this chapter some theoretical results got in this paper will be validated with three results published about the Virgo cluster.

The first test is relative to the  $R_{ZV}$  and its associated mass. These results were studied in chapter 6.

The second test is relative to the gravitating mass associated to the twice of virial radius.

In the third test, the most important, it is postulated that DMbG theory is able to multiply by the factor  $U \cdot \sqrt{1.85} = 2.7 \cdot 1.36 = 3.67$  the current parameter of local matter density  $\Omega_m^{local} = 0.08$  reaching 0.294 which match with the value  $\Omega_m^{Global} = 0.3$  accepted currently by the scientific community.

**7.1 GRAVITATING MASS ASSOCIATED UP TO THE ESTIMATED ZERO VELOCITY RADIUS AT 7.3 Mpc**

Clipped text of [7] O. Kashibadze.2020

**7. Concluding remarks**

The analysis of galaxy motions in the outskirts of the Virgo cluster makes it possible to measure the radius of the zero-velocity surface,  $R_0 = 7.0 - 7.3$  Mpc (Karachentsev et

al. 2014, Shaya et al. 2017, Kashibadze et al. 2018), corresponding to the total mass of the Virgo cluster  $M_T = (7.4 \pm 0.9) \times 10^{14} M_\odot$  inside the  $R_0$ . The numerical simulated trajectories of nearby galaxies with accurate distance estimates performed by Shaya et al. (2017) confirmed the obtained estimate of the total mass of the cluster. The virial mass of the cluster, being determined independently at the scale of  $R_g = 1.7$  Mpc from the internal motions, is nearly the same -  $M_{VIR} = (6.3 \pm 0.9) \times 10^{14} M_\odot$ . The agreement of

In the clipped text [7] O. Kashibadze, the authors gives the interval for [ 7, 7.3] Mpc for  $R_{ZV}$ .

According DMbG theory, the zero velocity radius  $R_{ZV} = 7.75$  Mpc, see epigraph 6.3, so the relative difference is only 6%.

This is a very good match between experimental results and theoretical results by DMbG theory.

Also at the concluding remarks [7] O.

Kashibadze, the authors give for the total mass  $M_T (< R_0) = (7.4 \pm 0.9) \cdot 10^{14} M_\odot$ . As this value is got by dynamical measures in fact this value must be considered as gravitational mass.

The theoretical value (6.12) calculated in epigraph 6.4 is  $M_G (< R_{ZV}) \approx 1.5 \cdot M_{VIR} = 9.48E14 M_\odot$  which is 14 % bigger regarding the value given by the authors, if it is considered the upper value of the interval. So both results may be considered compatibles.

In table 15 are summarized and compared the observational results and theoretical results.

Table 10 Virgo cluster	[7] O. Kashibadze	DMbG theory	Relative difference
$R_{ZV}$	[ 7, 7.3] Mpc	7.75 Mpc	6% - Very good
$M_{VIR}$	$(6.3 \pm 0.9) \cdot 10^{14} M_\odot$		
$M_G (< R_{ZV})$	$(7.4 \pm 0.9) \cdot 10^{14} M_\odot$	$1.5 \cdot M_{VIR} = 9.45 \cdot 10^{14} M_\odot$	14% - Compatibles

**7.2 GRAVITATING MASS ASSOCIATED UP TO THE TWICE OF VIRIAL RADIUS**

As  $R_{VIR} = 1.7$  Mpc its twice value is 3.4 Mpc. As  $R_{ZG} = 12.9$  Mpc then  $f = 3.4/12.9 = 0.2635$  and

$M_G (< 2 \cdot R_{VIR}) = 1.335 \cdot M_{VIR} = 8.4 \cdot 10^{14} M_\odot$ . This value match perfectly with the interval of masses given below in the clipped text.

As mentioned above, the Planck Collaboration (2016) performed a detailed study of the Virgo cluster through Sunyaev-Zeldovich effect and found the total mass of warm/hot gas to be $(1.4 - 1.6) \times 10^{14} M_\odot$ . Assuming the cosmic value for the baryon fraction, $f_b = \Omega_b / \Omega_m = 0.1834$ , they found that the total mass of the cluster would be $(7.6 - 8.7) \times 10^{14} M_\odot$ on a scale up to 2 times larger than the virial radius.	Clipped text from page 9 of [7] O. Kashibadze.2020
--	--

The total mass mentioned in Kashibadze paper it is considered in this paper as gravitating mass because in the whole paper of Kashibadze et al. they use always the concept of total mass as a result of dynamical measures so it is more suitable to interpret his total mass as gravitating mass.

Anyway, considering the total mass given by the formula (5.16)  $M_{TOTAL} (< 2R_{VIR}) = 1.386 \cdot M_{VIR} = 8.73 \cdot 10^{14} M_\odot$  that match with the upper value of mass range given by the authors.

### 7.3 SOLVING THE CONUNDRUM: LOCAL DENSITY MATTER VERSUS GLOBAL DENSITY MATTER

Below is the clipped text of a paper published for a team of well known astrophysicist.

As it has been noted by different authors (Vennik 1984, Tully 1987, Crook et al. 2007, Makarov & Karachentsev 2011, Karachentsev 2012), the total virial masses of nearby groups and clusters leads to a mean local density of matter of  $\Omega_m \simeq 0.08$ , that is 1/3 the mean global density  $\Omega_m = 0.24 \pm 0.03$  (Spergel et al. 2007). One possible explanation of the disparity between the local and global density estimates may be that the outskirts of groups and clusters contain significant amounts of dark matter beyond their virial radii, beyond what is anticipated from the integrated light of galaxies within the infall domain. If so, to get agreement between local and global values of  $\Omega_m$ , the total mass of the Virgo cluster (and other clusters) must be 3 times their virial masses. A measure of this missing

Clipped text from introduction of paper [6] Karachentsev I.D., R. Brent Tully, et al. 2014

In page 3 of that paper, they state that at the nearby clusters the mean local density of matter is  $\Omega_m^{local} = 0.08$ , whereas the global mass density in the Universe is  $\Omega_m^{global} = 0.24$ . (data year 2007)

Currently the data updated for scientific community is  $\Omega_m^{global} = 0.3$

The authors suggest that a possible solution for this tension would be that the total mass for cluster haloes must be three times the virial mass. That is justly what is found in this paper studying the DM at cluster scale as universal law in the framework of DMbG theory.

In chapter 5, the formula (5.13)  $M_{TOTAL}(< R_{ZG}) = \frac{\sqrt[5]{100}}{\Omega_{DE}^{1/5}} M_{VIR} = U \cdot M_{VIR}$ , shows that the total mass (baryonic and DM) enclosed into the halo cluster is  $U \approx 2.7$  times the virial mass equal to  $M_{TOTAL}(< R_{VIR}) \approx M_{200}$

However in the epigraph 5.9 has been calculated an extension of the halo cluster up to radius  $R_E = 1.85 \cdot R_{ZG}$  to obtain a ratio  $\frac{\Omega_{TM}^{LOCAL}}{\Omega_{DE}^{LOCAL}} = \frac{\Omega_{TM}^{GLOBAL}}{\Omega_{DE}^{GLOBAL}} = 3/7$ . Now using the formula (5.16) it is right to calculate the total mass enclosed by such sphere  $M_{TOTAL}(< R_E) = M_{TOTAL}(< f \cdot R_{ZG}) = \sqrt{f} \cdot U \cdot M_{VIR}$  being  $f = 1.85$  the factor  $\sqrt{f} \cdot U = 3.67$  then  $M_{TOTAL}(< R_E) = 3.67 \cdot M_{VIR}$  (7.1).

Therefore with such factor the parameter  $\Omega_m^{local} = 0.08$  is increased up to a  $\Omega_m^{local} = 0.08 \cdot 3.67 = 0.2936$  (7.2) because according [6] Karachentsev et al. 2014, the coefficient  $\Omega_m^{local} = 0.08$  is calculated considering the virial masses of clusters into the Local Universe, and as according DMbG such total mass is increased by the factor 3.67 if it is considered an extended halo with radius  $R_E = 1.85 \cdot R_{ZG}$  then the coefficient  $\Omega_m^{local} = 0.08 \cdot 3.67 = 0.293$  match perfectly with  $\Omega_m^{global} = 0.3$

This result enables an experimental test to validate these theoretical findings: If at the present Universe, the average distance between clusters is about its  $R_E$  associated to each one, then the DMbG theory would explain the current  $\Omega_m^{GLOBAL} = 0.3$

In other words, considering that almost the total baryonic matter at the current Universe is enclosed inside the virial radius of the clusters, as it is confirmed by multiples measures, the DMbG is able to justify that  $\Omega_m^{LOCAL} = \Omega_m^{GLOBAL}$  on condition that the average distance between clusters is the extended radius  $R_E$

## 8. CONCLUDING REMARKS

Thanks direct mass  $M_{DIRECT}(<r) = \frac{a^2 \cdot \sqrt{r}}{G}$  (4.1) and using the approximation of virial mass for  $M_{200}$  and the virial radius for  $R_{200}$  it is possible to get the formula  $a^2 = (G \cdot M_{VIR})^{5/6} \cdot (10 \cdot H)^{1/3}$  (4.4), which is on the base of all the important formulas got in this paper.

In chapter 4, the direct mass has been extended to clusters, and it is possible to state that all the new theoretical results obtained in this paper are based on this formula, co working with the well known properties of DE.

- A) The universality of ratio  $R_{ZG} / R_{VIR} \approx 7.3$  and its total mass associated, being  $M_{TOTAL}(<R_{ZG}) \approx 2.7 \cdot M_{VIR}$
- B) For any cluster at  $0.488 \cdot R_{ZG}$  is reached the maximum of gravitating mass and  $M_G(<R_M) \approx 1.57 \cdot M_{VIR}$
- C) The universality of the ratio  $R_{ZV} / R_{ZG} \approx 0.602$ , and its gravitating mass  $M_G(<R_{ZV}) \approx 1.5 M_{VIR}$
- D) The universality of the ratio  $R_E / R_{ZG} \approx 1.85$ , where the ratio total mass density versus DE density is  $3/7$  and its total mass  $M_{TOTAL}(<R_E) = 3.67 \cdot M_{VIR}$

Finally in chapter 7, are introduced some results published about Virgo cluster that back fully the previous findings:

- 1° Regarding the property D) may be consulted [ ] Karachentsev I.D.,R. Brent Tully.2014. to understand more in deep the current tension between the low value for Local mass density parameter  $\Omega_m = 0.08$  and the current global matter density parameter  $\Omega_m = 0.3$ .
- 2° Calculus of the zero velocity radius and its associated gravitating mass of Virgo are compatibles with the result published by [7] Olga Kashibadze et al. 2020, see epigraph 7.1
- 3° Calculus of the mass gravitating at two times the virial radius match fully with the result published by [7] Olga Kashibadze et al. 2020, see epigraph 7.2

These new theoretical findings offer to scientific community a wide number of tests to validate or reject the theory. The validation of DMbG theory would mean to understand that DM is a quantum gravitation effect, giving to scientific community new elements to continue searching a quantum gravitation theory.

## 9. BIBLIOGRAPHYC REFERENCES

- [1] Abarca, M. 2023. viXra:2312.0002 *A dark matter theory by gravitation for galaxies and clusters-v2*
- [2] Sofue, Y.2015. arXiv:1504.05368v1 *Dark halos of M31 and the Milky Way.*
- [3] Sofue, Y.2020.MDPI/Galaxies *Rotation curve of the Milky Way and the dark matter density.*
- [4] R.Ragusa et al.2022 arXiv:2212.06164v1  
*Does the virial mass drive the intra-cluster light? The relationship between the ICL and Mvir from VEGAS*
- [5] Chernin, A.D. et al.2013. arXiv:1303.3800 *Dark energy and the structure of the Coma cluster of galaxies*
- [6] Karachentsev I.D.,R. Brent Tully, et al. 2014. arXiv:1312.6769v2  
*Infall of nearby galaxies into the Virgo cluster as traced with HST*
- [7] Olga Kashibadze, Igor Karachentsev et al.2020. arXiv:2002.12820v1  
*On structure and kinematics of the Virgo cluster of galaxies*
- [8] Abarca, M, 2024. viXra:2404.0131  
*Relation Between Dark Matter and Dark Energy in Galaxy Clusters and Its Successful Implications*
- [9] Abarca, M.2014. viXra:1410.0200 *Dark Matter Model by Quantum Vacuum*