


PRECLINICAL RESEARCH

Hydrogen Sulfide Attenuates Renin Angiotensin and Aldosterone Pathological Signaling to Preserve Kidney Function and Improve Exercise Tolerance in Heart Failure



Zhen Li, BS,^{a,b} Chelsea L. Organ, PhD,^{a,b} Jianming Kang, PhD,^c David J. Polhemus, PhD,^{a,b} Rishi K. Trivedi, PhD,^{a,b} Thomas E. Sharp III, PhD,^a Jack S. Jenkins,^a Ya-xiong Tao, PhD,^d Ming Xian, PhD,^c David J. Lefer, PhD^{a,b}



HIGHLIGHTS

- Pressure overload induced by transverse aortic constriction led to severe heart failure accompanied by renal fibrosis and dysfunction, endothelial dysfunction, and exercise intolerance in mice.
- 3-week but not 10-week-delayed H₂S therapy preserved left ventricular ejection fraction and attenuated cardiac fibrosis.
- H₂S therapy attenuated renal fibrosis and dysfunction, preserved endothelium, and enhanced treadmill exercise capacity.
- H₂S therapy blunted the maladaptive overactivation of the sympathetic nervous system, resulting in reduced renal tissue norepinephrine levels as well as attenuated circulating levels of renin, angiotensin II, and aldosterone.

From the ^aCardiovascular Center of Excellence, Louisiana State University Health Sciences Center, New Orleans, Louisiana; ^bDepartment of Pharmacology and Experimental Therapeutics, Louisiana State University Health Sciences Center, New Orleans, Louisiana; ^cDepartment of Chemistry, Washington State University, Pullman, Washington; and the ^dDepartment of Anatomy, Physiology, and Pharmacology, Auburn University College of Veterinary Medicine, Auburn, Alabama. Supported by National Heart, Lung, and Blood Institute/National Institutes of Health grants 5R01HL092141, 5R01HL093579, and 1U24 HL094373 to Dr. Lefer; and grant 1R01HL116571 to Drs. Xian and Lefer. Funding was also received from the Louisiana State University Medical School Foundation and LSU Medical School Alumni Association. All authors have reported that they have no relationships relevant to the contents of this paper to disclose.

SUMMARY

Cardioprotective effects of H₂S have been well documented. However, the lack of evidence supporting the benefits afforded by delayed H₂S therapy warrants further investigation. Using a murine model of transverse aortic constriction-induced heart failure, this study showed that delayed H₂S therapy protects multiple organs including the heart, kidney, and blood-vessel; reduces oxidative stress; attenuates renal sympathetic and renin-angiotensin-aldosterone system pathological activation; and ultimately improves exercise capacity. These findings provide further insights into H₂S-mediated cardiovascular protection and implicate the benefits of using H₂S-based therapies clinically for the treatment of heart failure.

(*J Am Coll Cardiol Basic Trans Science* 2018;3:796-809) © 2018 The Authors. Published by Elsevier on behalf of the American College of Cardiology Foundation. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

ABBREVIATIONS AND ACRONYMS

BNP = B-type natriuretic peptide

cGMP = cyclic guanosine monophosphate

HF = heart failure

LVEF = left ventricular ejection fraction

RAAS = renin-angiotensin-aldosterone system

SNS = sympathetic nervous system

TAC = transverse aortic constriction

Hear failure (HF) is a heterogeneous disease resulting from a number of common pathological stimuli including, but not limited to, long-standing hypertension, profound inflammation, and overactivation of neurohumoral pathways (1). HF is a major public health problem in the United States, with a prevalence of 7 million people, and it is projected to affect more than 9 million people by 2030 (2). Due to its heterogeneous nature, the consequences of HF are not limited to cardiac pathology. Noncardiac comorbidities such as renal dysfunction and vascular dysfunction frequently accompany HF and further decrease patients' quality of life and clinical outcomes (3,4). For example, HF patients with renal dysfunction have greater chance for readmission, longer hospital stay, and worse prognosis, and vascular injury has been linked to reduced exercise capacity, leading to decreased quality of life (5,6).


Neurohumoral dysregulation is a hallmark feature in many cardiovascular diseases including heart failure (7-10). Specifically, overactivation of the sympathetic nervous system and the renin-angiotensin-aldosterone system (RAAS) exacerbates HF symptoms and contributes to the development of comorbidities (4,11,12). In fact, efficacious first-line medications for HF such as beta-blockers, mineralocorticoid receptor antagonists, angiotensin-converting enzyme (ACE) inhibitors, and angiotensin receptor blockers (ARBs) target the sympathetic nervous system (SNS) and the RAAS overactivity. However, these medications are imperfect and have several limitations. For example, beta-blockers, mineralocorticoid

receptor antagonists, and ARBs are receptor antagonists designed to compete with endogenous ligands without altering or decreasing the levels of endogenous ligands, which potentially limits their vascular protective effects (13,14). Distinct from receptor antagonists, ACE inhibitors reduce the endogenous production of angiotensin II (Ang II), but its use is contraindicated in patients with certain cardiorenal syndromes due to unwanted side effects (15). Current treatments inadequately manage the diverse population of HF patients in a manner that effectively extends longevity and quality of life. Therefore, examination of novel pharmacotherapies that have the ability to protect multiple organ systems in heart failure is warranted.

SEE PAGE 810

H₂S plays a key role in cardiovascular homeostasis and cardioprotection (16-20). H₂S wields a wide range of biological functions in the cardiovascular system, including mitochondria biogenesis, metabolic modulation, angiogenesis, vasodilation and blood pressure regulation, inhibition of inflammation, antioxidant, and antiapoptosis (21). Furthermore, H₂S reportedly exerts protective effects in multiple organ systems including vasculature and the kidney (22-24), both of which are involved in the pathogenesis and progression of HF. Taken together, H₂S represents a promising therapeutic target for the treatment of HF and its comorbidities.

JK-1 is a novel, phosphorothioate-based synthetic H₂S donor that releases H₂S upon hydrolysis. Recently we reported that administration of JK-1 at the time of



reperfusion significantly reduced infarct size and circulating troponin-I levels in a murine model of myocardial ischemia-reperfusion injury (20). In the present study, using a well-established pressure overload HF model, the efficacy of delayed therapy with JK-1 on cardiac, renal, and vascular injury and dysfunction in the setting of HF was investigated.

METHODS

EXPERIMENTAL COMPOUNDS. JK-1 and its control compound were designed and synthesized by

Dr. Ming Xian et al. (20). The optimal dose of JK-1 was selected based on the results of a previous study investigating the infarct-sparing effects of JK-1 (20). The structures of JK-1 and control are described in **Figure 1A**.

EXPERIMENTAL ANIMALS. Male C57BL/6J mice (Jackson Laboratory, Bar Harbor, Maine) were 9 weeks of age at the time of study initiation. All experimental protocols were approved by the Institute for Animal Care and Use Committee at LSU Health Sciences Center and conformed to the Guide for the Care and Use of Laboratory Animals.

All experimental animals received humane care in accordance with National Society of Medical Research and National Institutes of Health tenets.

TRANSVERSE AORTIC CONSTRICTION PROTOCOL. Cardiac hypertrophy and heart failure were induced through transverse aortic constriction (TAC) as previously described (19,25,26).

EXPERIMENTAL DESIGN AND TREATMENT REGIMEN. Sham or TAC procedures were performed 1 week after baseline echocardiography. All mice subjected to the TAC procedure were randomly assigned to 1 of 3 study groups. One group received control treatment starting at 3 weeks post TAC and continued until 18 weeks post TAC. The second group was treated with JK-1 starting at 3 weeks post TAC. The third group started with control at 3 weeks post TAC and were then switched to JK-1 at 10 weeks post TAC. All treatments were administered twice a day at a dosage of 100 µg/kg (200 µg/kg/day) through i.p. injection. A detailed experimental protocol is shown in [Figure 1B](#).

ECHOCARDIOGRAPHY. Echocardiography was performed at baseline and once every 3 weeks (model Vevo-2100 ultrasonography system; Visual Sonic, Baltimore, Maryland) as previously described (25). M-mode long-axis views were used to measure the thickness of interventricular septal walls at systole and diastole and left ventricle (LV) posterior wall at systole and diastole. LV chamber diameters at end systole and end diastole and LV ejection fraction (LVEF) were also measured using M-mode long-axis images.

LV HEMODYNAMICS ASSESSMENT. Invasive hemodynamic measurement was performed at the 18-week endpoint using a 1.2-F transonic pressure catheter. The pressure catheter was inserted into the right common carotid and advanced into the LV. Systolic and diastolic arterial blood pressure, LV end diastolic pressure, and relaxation constant Tau values were recorded. Equally elevated systolic blood pressure across groups indicates a successful TAC procedure ([Supplemental Figure S3A](#)).

HISTOLOGICAL ASSESSMENT. Cardiac and renal fibrosis were assessed at the 18-week endpoint as previously described (26). To quantify global fibrosis from the Picrosirius red-stained slides, fluorescent images were obtained (original magnification ×20) using an Olympus (model BX53-DP80, Tokyo, Japan) system and analyzed using ImageJ software (U.S. National Institutes of Health, Bethesda, Maryland).

QUANTITATIVE PCR. RNA was isolated from the heart and kidney tissues of control and JK-1-treated mice at 18 weeks post TAC. One milligram of RNA was

transcribed using I-script cDNA synthesis kit (Bio-Rad, Portland, Maine). TaqMan primers for collagen 1a1 (Col1a1), collagen 3a1 (Col3a1), interleukin (IL)-6, connective tissue growth factor (CTGF), and fibronectin (FN1) were used to amplify by quantitative polymerase chain reaction (qPCR). For real-time qPCR experiments, 18S ribosomal RNA was used as a housekeeping gene, and $2^{-\Delta\Delta\text{CT}}$ values corrected to 18S rRNA were used for data analysis.

CIRCULATING B-TYPE NATRIURETIC PEPTIDE QUANTIFICATION. Circulating B-type natriuretic peptide (BNP) levels were measured at 18 weeks post TAC using an enzyme immunoassay (catalog number EK-011-23; Phoenix Pharmaceuticals, Burlingame, California).

PLASMA CREATININE QUANTIFICATION. Creatinine levels were measured at 18 weeks post TAC using a commercially available assay kit (catalog number ab65340, Abcam, Cambridge, Massachusetts).

AORTIC VASCULAR REACTIVITY ASSESSMENT. Vascular function was measured in isolated segments (3 mm in length) of thoracic aorta as previously described (27) at 18 weeks following TAC in all study groups. Vascular rings were equilibrated for 60 min and then pre-contracted with phenylephrine (1 µM). The vascular rings were then challenged with increasing concentrations of acetylcholine (10^{-9} to 10^{-5}) and sodium nitroprusside (SNP) at concentrations from $10^{-10.5}$ to $10^{-6.5}$. Data are reported as percent of relaxation from maximum contraction to phenylephrine.

EXERCISE CAPACITY ASSESSMENT. Treadmill exercise capacity was measured using an rodent treadmill (IITC, Woodland Hills, California) at 18 weeks post TAC as previously described with modifications (28). Briefly, immediately before the test, mice were put on the treadmill to acclimate for 15 min. Then, the mice are subjected to a slow walk/run period to be familiarized with treadmill running. In this period, the treadmill is set to start at 0 m/min speed and increased by 1 m/min every min for 10 min. Finally, in the test phase, the treadmill was set to a 30° incline, and the starting speed was set to 10 m/min and programmed to increase by 2.67 m/min every min for 3 min, at which time the treadmill reached and maintained its top speed at 18 m/s. All mice were run at this setting until they reach a state of exhaustion. Duration and distance of the running and shocks per minute were recorded. Shocks per minute is a measurement of endurance. As mice felt tired and fell behind the speed of the treadmill, they received an electrical shock by the shocking grid. Higher shocks per min indicated lower endurance. A mouse was deemed to be exhausted when it spent more than 5 consecutive s at the back of

the treadmill despite receiving electrical shocks repeatedly.

CIRCULATING 8-ISOPROSTANE MEASUREMENT. Plasma, myocardial, and renal 8-isoprostane levels were measured at 18 weeks post TAC by using an enzyme-linked immunosorbent assay kit (catalog number 516351, Cayman Chemical, Ann Arbor, Michigan). Tissue levels of 8-isoprostane were adjusted based on protein concentration (catalog number 23225; Thermo Fisher Scientific, Halethorpe, Maryland).

SNS AND RAAS OVERACTIVATION ASSESSMENT. Renal norepinephrine levels were measured at 18 weeks post TAC, using an enzyme-linked immunosorbent assay kit (catalog number KA3836, Abnova, Taipei, Taiwan). Renin (catalog number EMREN1, Thermo Fisher Scientific), Ang II (catalog number ADI-900-204, Enzo Clinical Labs, Farmingdale, New York), and aldosterone (catalog number ADI-900-173, Enzo) levels were measured at 18 weeks post TAC using commercially available enzyme immunoassays. Renal levels of norepinephrine and aldosterone were adjusted based on protein concentration.

H₂S AND NO MEASUREMENTS. Plasma, myocardial, and renal H₂S and nitric oxide (NO) levels were measured at 18 weeks post TAC as previously described (18,29,30). Tissue levels of H₂S and NO were adjusted based on protein concentration.

cGMP MEASUREMENT. Cardiac and renal cyclic guanosine monophosphate (cGMP) levels were measured at 18 weeks post TAC as previously described (31).

STATISTICAL ANALYSIS. All data in this study are mean ± SEM. Differences in data among the groups were compared using 1-way ANOVA analysis or 2-way ANOVA analysis, using Prism 6 software (GraphPad, San Diego, California) followed by a Bonferroni multiple comparison test. A p value of <0.05 was considered statistically significant. Presented data may have different numbers of animals per endpoint due to procedural complications, limited sample collection (i.e., plasma volume), or lack of participation in involuntary treadmill running. Prior to conducting statistical analysis, an outlier test was performed using “ROUT” method developed by Prism 6 to identify and remove any outlier in the data set.


RESULTS

DELAYED TREATMENT WITH JK-1 ATTENUATES CARDIAC DILATION, PRESERVES LV EJECTION FRACTION, IMPROVES LV HEMODYNAMICS, AND REDUCES CARDIAC FIBROSIS AFTER TAC. The effects of delayed therapy with JK-1 on the failing heart

were evaluated using a TAC model of HF (Figure 1). Administration of JK-1 was initiated at 3 or 10 weeks post TAC and continued until 18 weeks post TAC. Mice treated with JK-1 initiated at 3 weeks or 10 weeks post TAC displayed significant reductions in LV chamber dimensions at end-diastole and end-systole compared to control mice (Figures 2A and 2B). In contrast, only the mice from 3-week post-TAC-treated group had preserved LVEF (Figure 2C), and there was no effect on LVEF in those that received JK-1 starting at 10 weeks following TAC. Invasive hemodynamic measurements revealed that delayed JK-1 treatment significantly decreased LV end-diastolic pressure and shortened relaxation constant Tau (Figures 2D and 2E). In addition, circulating BNP levels were significantly reduced in mice that received delayed JK-1 treatment (Figure 2F). Furthermore, the TAC-induced global LV hypertrophic response and subsequent wall thinning were attenuated in both of the JK-1 treatment groups (Supplemental Figures S1A to S1D). We also demonstrated that, at the 18-week endpoint, the whole heart weight and left atrial weight-to-tibia length ratio were significantly lower in JK-1-treated mice than in control mice, indicating that JK-1 reduced pathological remodeling in pressure overload HF (Supplemental Figures S2A to S2C).

Next, we evaluated the effects of delayed JK-1 treatment on cardiac fibrosis with PicroSirius red staining at 18 weeks post TAC. Representative photomicrographs of heart sections are shown in Figure 3A. Treatment with JK-1 initiated at 3 weeks post TAC significantly reduced cardiac fibrosis whereas there was no significant change when JK-1 treatment was initiated at 10 weeks post TAC (Figure 3B). Furthermore, 3-week-delayed JK-1 treatment mitigated the transcription of pro-fibrotic genes such as collagen 1a1, collagen 3a1, interleukin-6, connective tissue growth factor, and fibronectin in the heart (Figure 3C).

DELAYED TREATMENT WITH JK-1 AMELIORATES RENAL DYSFUNCTION AND REDUCES RENAL FIBROSIS IN HF. The effect of JK-1 on renal function in HF was assessed at 18 weeks post TAC by plasma creatinine measurements. Mice treated with Control exhibited impaired renal function as shown by elevated plasma creatinine levels that were compared to Sham mice. Mice treated with JK-1 initiated at 3 weeks post TAC and 10 weeks post TAC displayed significant reductions in plasma creatinine levels, indicating preserved renal function (Figure 4D). In addition, histological assessment of PicroSirius red-stained kidney sections revealed increased renal




fibrosis in mice subjected to the TAC procedure and those that received Control treatment. Contrary to our observation in cardiac fibrosis, we saw significantly lessened amounts of renal fibrosis in mice that received 3-week-delayed or 10-week-delayed JK-1 treatment (Figure 4A and 4B). Furthermore, both 3-week- and 10-week-delayed JK-1 treatment significantly mitigated the transcription of interleukin-6 and connective tissue growth factor in the kidney, whereas only 3-week-delayed treatment significantly reduced the transcription of collagen 1a1 and fibronectin (Figure 4C).

DELAYED TREATMENT WITH JK-1 ATTENUATES ENDOTHELIAL DYSFUNCTION IN HF. Vascular reactivity of isolated thoracic aortas to acetylcholine

and SNP is shown in Figure 5. Aortas from mice that received the Control compound displayed significantly worsened responses to acetylcholine, as shown in concentration-response curve, maximal relaxation reached, and Half maximal effective concentration (EC₅₀) values. Interestingly, delayed treatment with JK-1 (3 and 10 weeks delayed) resulted in better relaxation responses to acetylcholine (Figures 5A to 5C).

We also evaluated vascular relaxation responses of isolated thoracic aorta to SNP. Aortas from mice that received the Control compound displayed worsened vasodilation only at concentrations of 1 and 10 nM SNP, and no significant differences were observed in maximal relaxation and EC₅₀ (Figures 5D to 5F). Taken together, these data suggest that TAC-induced HF

FIGURE 3 Cardiac Fibrosis at 18 Weeks Post TAC


(A) Representative images (original magnification $\times 20$) of Picrosirius red-stained cardiac tissue sections from HF + Control and HF + JK-1 (3 weeks and 10 weeks, respectively) delayed treatment. (B) Quantified cardiac fibrosis. (C) Relative fold changes in myocardial mRNA levels of a panel of fibrosis-related genes in mice from HF + Control, HF + JK-1 3 weeks post TAC, and HF + JK-1 10 weeks post TAC groups at 18 weeks post TAC. All data in Figure 3 were analyzed with 1-way ANOVA. Circles inside bars denote sample size. Abbreviations as in Figure 1.

resulted in severe endothelial dysfunction, and delayed treatment with JK-1 partially corrected this dysfunction.

DELAYED TREATMENT WITH JK-1 ENHANCED TREADMILL EXERCISE PERFORMANCE. Reduced exercise capacity is one of the hallmark symptoms of HF in patients. We examined the effects of delayed therapy with JK-1 on exercise capacity using a rodent treadmill at 18 weeks post TAC (Figure 6). Mice from the HF + Control group displayed greatly reduced time to exhaustion and distance of running and experienced higher numbers of shocks per min than Sham mice, suggesting that TAC-induced HF resulted in reduced exercise capacity and lower endurance. Mice that received JK-1 treatment initiated at 3 or 10 weeks post TAC showed significantly improved exercise capacity as they had better treadmill running duration and distance and received fewer shocks per min.

DELAYED TREATMENT WITH JK-1 AUGMENTS H₂S AND NO BIOAVAILABILITY, AND REDUCED SYSTEMIC AND TISSUE LEVELS OF OXIDATIVE STRESS. Next, we examined the effects of delayed JK-1 treatment on the bioavailability of H₂S, NO, and levels of oxidative stress at 18 weeks post TAC (Figure 7). Mice treated with JK-1 starting at 3 or 10 weeks post TAC had significantly elevated levels of circulating H₂S (Figure 7A), myocardial tissue H₂S (Figure 7B), and renal tissue H₂S (Figure 7C), further verifying the ability of JK-1 to donate pharmacologically relevant concentrations of H₂S.

Multiple studies have previously reported that H₂S augments the bioavailability and signaling of NO (18,31-33). The current study found that JK-1 treatment initiated at 3 or 10 weeks post TAC resulted in significantly elevated levels of nitrite in circulation, heart, and kidney (Figures 7D to 7F). In addition, significantly elevated levels of cGMP were also observed in cardiac and renal tissue at 18 weeks post



TAC in mice receiving JK-1 treatment (Supplemental Figures S4A and S4B).


In addition, it has been reported that both H₂S and NO are powerful antioxidants (31,34). Therefore, we examined the levels of oxidative stress by using 8-isoprostane as a biomarker (Figures 7G to 7I). TAC-induced HF resulted in elevated oxidative stress as the mice receiving the Control compound exhibited significantly higher levels of 8-isoprostane in circulation, heart, and kidney. Such elevation in oxidative stress was ameliorated by both the 3- and 10-week-delayed JK-1 treatment.

DELAYED TREATMENT WITH JK-1 REDUCED RENAL NOREPINEPHRINE CONTENTS, ATTENUATED RAAS SIGNALING IN HF. Neurohumoral dysregulation is a significant contributor to the pathogenesis of HF and its co-morbidities (14,35). Specifically, overactivation of the SNS and the RAAS significantly exacerbates the severity of HF (4). In the current study, we evaluated the extent of SNS and RAAS overactivation by measuring renal or circulating levels of norepinephrine, renin, Ang II, and aldosterone (Figure 8). We observed that TAC-induced HF significantly activated

the SNS as mice receiving the Control compound displayed nearly 20-fold increase in renal norepinephrine content. JK-1 treatment initiated at 3 or 10 weeks post TAC significantly attenuated renal norepinephrine content compared to mice receiving Control (Figure 8A). Furthermore, delayed JK-1 treatment ameliorated RAAS activation as mice that received 3-week- and 10-week-delayed JK-1 treatment had significantly reduced levels of circulating renin (Figure 8B), circulating Ang II (Figure 8C), and circulating aldosterone (Figure 8D) compared to those that received Control. Similarly, renal aldosterone content was significantly reduced in mice that received either 3-week- or 10-week-delayed JK-1 therapy (Figure 8E).

DISCUSSION


It was previously shown that pharmacological H₂S therapy ameliorates TAC-induced HF severity (19,25,36,37). However, one major limitation of those studies is that the experimental H₂S therapies were administered prior to or right after the induction of HF

FIGURE 5 Aortic Vasorelaxation at 18 Weeks Post TAC

Freshly isolated thoracic aortic rings were allowed to equilibrate, and pre-contracted with 1 μ M phenylephrine. **(A)** Concentration-response curve (% relaxation from 10^{-9} to 10^{-5}). **(B)** EC₅₀ (nM). **(C)** Maximal relaxation (%) achieved for acetylcholine (ACh)-mediated vasorelaxation in mice from Sham, HF + Control, HF + JK-1 3 weeks post TAC, and HF + JK-1 10 weeks post TAC groups at 18 weeks post TAC. **(D)** Concentration-response curve (% relaxation from $10^{-10.5}$ to $10^{-6.5}$) **(E)** EC₅₀ (nM), and **(F)** maximal relaxation (%) achieved for SNP mediated vasorelaxation in mice from Sham, HF + Control, HF + JK-1 3 weeks post TAC, and HF + JK-1 10 weeks post TAC groups at 18 weeks post TAC. * $p < 0.05$ between Sham vs. HF + Control; **** $p < 0.0001$ between Sham vs. HF + Control; # $p < 0.05$ between HF + JK-1 3 weeks post TAC vs. HF + Control; ## $p < 0.01$ between HF + JK-1 3 weeks post TAC vs. HF + Control; ### $p < 0.0001$ between HF + JK-1 3 weeks post TAC vs. HF + Control; \$\$\$ $p < 0.001$ between HF + JK-1 10 weeks post TAC; \$\$\$\$ $p < 0.0001$ between HF + JK-1 10 weeks post TAC. Concentration-response curve were generated using a model of nonlinear regression with variable slope followed by 2-way-ANOVA analysis. EC₅₀ and maximal relaxation data were analyzed with 1-way ANOVA. **Circles inside bars** denote the sample size. EC₅₀ = half maximal effective concentration; SNP = sodium nitroprusside; other abbreviations as in [Figure 1](#).

by TAC. Such treatment regimens do not reflect clinical scenarios where medications are given to patients after the establishment of cardiac dysfunction or failure (8,38). The current study examined the efficacy of a novel H₂S donor, JK-1, in TAC-induced HF with a delayed treatment approach. To better simulate the clinical situation, we initiated the H₂S therapy with JK-1, either at 3 or 10 weeks post TAC. At 3 weeks post TAC, echocardiographic assessment revealed that all mice subjected to TAC started to develop cardiac hypertrophy. At 10 weeks following TAC, all mice that were subjected to TAC and received the Control compound had LVEFs below 45%.

We observed significant preservation of LVEF in mice treated with JK-1 starting at 3 weeks post TAC but not in those started at 10 weeks post TAC. This preservation in LVEF was accompanied by reduction in cardiac fibrosis, which was also seen only in mice starting to receive H₂S therapy at 3 weeks post TAC. These data suggest that 10-week-delayed H₂S therapy fails to reverse established cardiac fibrosis and pathological remodeling in the failing heart. Interestingly, despite its neutral effects in LVEF and cardiac fibrosis, 10-week-delayed JK-1 therapy ameliorated other aspects of cardiac dysfunction. We observed significantly attenuated LV chamber dilatation, lower




LVEDP, and shorter relaxation constant Tau with both the 3-week- and the 10-week-delayed treatment compared to Control mice. These improvements in LV chamber size translated into significant reductions in circulating BNP levels for both the 3-week- and the 10-week-delayed JK-1 treatments. Taken together, these data demonstrate that delayed H₂S therapy is efficacious in TAC-induced HF with greater improvements in LV function when treatment is initiated early.

HF is a systemic condition that deteriorates the function of multiple organ systems including the kidney and the vasculature (3-6,15,39,40). As chronic HF progresses, cardiac pathologies such as overactive SNS and RAAS, inflammation, and excessive oxidative stress induces renal injury and impairs normal renal function. Such phenomenon is defined as type II cardiorenal syndrome and often results in poorer prognosis and survival (5,35). To date, no pharmacological therapeutic has been developed to simultaneously target both the failing heart and the injured kidney. Previous studies have reported that H₂S therapies effectively ameliorated renal dysfunction in several animal models of kidney diseases through differential mechanisms, suggesting that H₂S-based therapies may be efficacious in treating both the heart and the kidney in HF (14,38,41-47). In consistent with previous literature, we observed that the mice received Control developed type-II cardiorenal syndrome as evidenced by elevated levels of plasma creatinine and renal fibrosis at 18 weeks post TAC (39). Delayed treatment with JK-1 started at 3 or 10 weeks post TAC significantly lowered plasma creatinine levels, renal fibrosis, as well as the transcriptional levels of pro-fibrotic genes such as collagen 1a1, collagen

3a1, interleukin 6, connective tissue growth factor, and fibronectin. Taken together, our data provides the first evidence that H₂S therapy is efficacious for the treatment of cardiorenal syndrome induced by TAC.

In addition to cardiorenal syndrome, endothelial injury and dysfunction is another commonly seen comorbidities in HF (3,4). Endothelium is a monolayer of cells covering the inner surface of blood vessels that mediates the vascular hemostasis through providing NO, antiproliferative and anti-inflammatory actions. Endothelial dysfunction results in the inactivation of endothelial NOS (eNOS) and attenuated NO bioavailability, over-produced inflammatory cytokines, and increased oxidative stress, which further exacerbate the progression of HF (6,48,49). Multiple studies have reported that H₂S augments endothelial function via reducing oxidative stress, promoting eNOS activity, and other mechanisms (16,33,50,51); However, there is a lack of information regarding the endothelium-protective actions of H₂S in the setting of HF. In the current study, we observed that the aortic rings of isolated from HF + Control mice exhibited diminished response to acetylcholine, verifying that TAC-induced HF indeed caused severe endothelium dysfunction. More importantly, delayed H₂S treatment with JK-1 attenuated endothelium dysfunction as shown by improved concentration-response curve, achieving greater maximal relaxation, and resulting in lower EC₅₀. This improvement in the vasodilatory response to acetylcholine was also reflected in enhanced NO bioavailability and signaling following JK-1 treatment.


Severe exercise intolerance is a primary chronic symptom associated with decreased quality of life in

FIGURE 7 H₂S and NO Bioavailability and 8-Isoprostane Levels at 18 Weeks Post TAC

H₂S levels (μM) in plasma (A), myocardial tissue (B), and renal tissue (C). Nitrite levels (μM) in plasma (D), myocardial tissue (E), and renal tissue (F). 8-Isoprostane (8-Iso) levels (pg/ml) in circulation (G), myocardial tissue (H), and renal tissue (I) in mice from Sham, HF + Control, HF + JK-1 3 weeks post TAC, and HF + JK-1 10 weeks post TAC groups at 18 weeks post TAC. All data were analyzed with 1-way ANOVA. Circles inside bars denote the sample size. Abbreviations as in Figure 1.

patients with HF, even when stable and well compensated (52). More recently, the endothelium has emerged as a therapeutic target for exercise intolerance in HF (6,53). Given the prominent endothelium-protective effects of JK-1 in HF, we measured the exercise capacity and we observed that TAC-induced HF significantly impaired the exercise capacity of mice treated with control, whereas JK-1 treatment initiated at 3 and 10 weeks post TAC significantly increased the duration and distance of treadmill running. Interestingly, despite the neutral effect on LVEF, the mice received 10-week-delayed JK-1 treatment displayed 3-fold increase in duration and distance of running. This suggests that vascular endothelial function plays an independent role from cardiac function in H₂S-mediated improvement in exercise capacity in TAC-induced HF.

Additionally, we examined the effects of delayed JK-1 treatment on the levels of SNS and RAAS activation. In the pathogenesis of HF, maladaptive SNS signaling increases in response to decreased cardiac output, which in turn, activates RAAS (8). SNS and RAAS overactivation are major maladaptive neurohumoral alterations in HF that accelerate the exacerbation of HF and its comorbidities (8,9). Previously, it was reported that NO exerts sympathoinhibitory effects via attenuating the release of norepinephrine from neural terminals (54). More recently, it was discovered that elevation in oxidative stress in central nervous system resulted in systemic sympathoexcitation (40). Given the powerful antioxidant and NO-promoting effects of H₂S, we propose that delayed therapy with JK-1 down regulates SNS and RAAS activation. We observed that renal



norepinephrine content was significantly reduced in kidneys from mice received 3-week- and 10-week-delayed JK-1 treatment when compared to HF + Control. Reduction in renal norepinephrine content was accompanied by reduced levels of renin, Ang II, and aldosterone in the circulation.

Although we observed similar degree of reductions in plasma creatinine levels, renal fibrosis, renal norepinephrine content, renin, and aldosterone levels for both 3-week- and 10-week-delayed JK-1-treated groups, the 3-week-delayed treatment offered greater reductions in circulating Ang II, and greater increases in NO bioavailability and signaling. This resulted in greater improvements in cardiac and vascular function as well as exercise capacity. This observation suggests that although H₂S therapy is efficacious in both early and late stages of the diseases, earlier intervention may offer better outcomes.

STUDY LIMITATIONS. We demonstrated the differential effects of 3-week-delayed versus 10-week-delayed JK-1

treatment in the context of pressure overload-induced heart failure; however, in order to verify the efficacy of JK-1 and fully understand its mechanism of action, further investigations using other heart failure models are warranted. In addition, although we showed that JK-1 treatment significantly reduced the plasma creatinine levels, a more clinically relevant index of kidney function would be glomerular filtration rate or urine creatinine-to-albumin ratio. Furthermore, the pharmacokinetics and effects of JK-1 treatment on naïve mice were not investigated in the current study. Such information would provide strong basis for the translation of JK-1 as a therapeutic agent.

CONCLUSIONS

We have shown that delayed treatment with a novel and potent H₂S donor, JK-1, protects multiple organs including the heart, peripheral arteries, and the kidneys against injury and dysfunctions in the setting of TAC- induced heart failure. Such protection in

multiple organs translated into improved exercise capacity. Furthermore, our results demonstrate that delayed treatment with JK-1 reduced the overactivation of the SNS and RAAS, possibly through enhancing NO bioavailability and signaling, and reducing oxidative stress. We conclude that delayed H₂S therapy is efficacious in HF, and is worthy of further development as a promising therapeutic for the treatment of HF and its comorbidities. Future studies will investigate the cellular and molecular mechanism by which H₂S mitigates the overactivation of SNS and RAAS in HF.

ACKNOWLEDGEMENTS The authors thank Ms. Jean Carnal for technical support and Dr. Luis Marrero and the Morphology and Imaging Core at LSUHSC for performing PicroSirius red staining of cardiac and renal tissues.

ADDRESS FOR CORRESPONDENCE: Dr. David J. Lefer, Cardiovascular Center of Excellence, Louisiana State University Health Sciences Center, 533 Bolivar Street, Room 408, New Orleans, Louisiana 70112. E-mail: dlefe1@lsuhsc.edu.

PERSPECTIVES

COMPETENCY IN MEDICAL KNOWLEDGE: In the setting of heart failure, the lack of proper perfusion and the activation of maladaptive signaling pathways such as RAAS lead to renal and endothelial dysfunction. Similar to NO, H₂S is an endogenous gaseous molecule that exerts protective effects on cardiovascular system. Here we used JK-1, a novel synthetic H₂S donor, to simultaneously protect the heart, the kidneys, the vessels, and to improved exercise performance in mice subjected to heart failure.

TRANSLATIONAL OUTLOOK: H₂S-based therapeutics have been tested in heart failure patients and proven to be safe and bio-active. Our study identified that JK-1, a novel H₂S donor, protects the heart against pressure overload-induced heart failure and reduces the collateral damages to the kidney and the endothelium. These findings support further clinical investigation of the efficacy of H₂S-based therapies for treating heart failure and its comorbidities.

REFERENCES

- Heineke J, Molkentin JD. Regulation of cardiac hypertrophy by intracellular signalling pathways. *Nat Rev Mol Cell Biol* 2006;7:589-600.
- Voigt J, Sasha John M, Taylor A, Krucoff M, Reynolds MR, Michael Gibson C. A reevaluation of the costs of heart failure and its implications for allocation of health resources in the United States. *Clin Cardiol* 2014;37:312-21.
- van Deursen VM, Urso R, Laroche C, et al. Comorbidities in patients with heart failure: an analysis of the European heart failure pilot survey. *Eur J Heart Fail* 2014;16:103-11.
- van Deursen VM, Damman K, van der Meer P, et al. Co-morbidities in heart failure. *Heart Fail Rev* 2014;19:163-72.
- Ronco C, Di Lullo L. Cardiorenal syndrome. *Heart Fail Clin* 2014;10:251-80.
- Van Craenenbroeck EM, Conraads VM. Mending injured endothelium in chronic heart failure: a new target for exercise training. *Int J Cardiol* 2013;166:310-4.
- Pagani M, Lucini D. Autonomic dysregulation in essential hypertension: insight from heart rate and arterial pressure variability. *Auton Neurosci* 2001;90:76-82.
- Muesan ML, Paini A, Agabiti Rosei C, Bertacchini F, Stassaldi D, Salvetti M. Current pharmacological therapies in heart failure patients. *High Blood Press Cardiovasc Prev* 2017;24:107-14.
- Grippo AJ, Johnson AK. Stress, depression and cardiovascular dysregulation: a review of neurobiological mechanisms and the integration of research from preclinical disease models. *Stress* 2009;12:1-21.
- Grippo AJ. Mechanisms underlying altered mood and cardiovascular dysfunction: the value of neurobiological and behavioral research with animal models. *Neurosci Biobehav Rev* 2009;33:171-80.
- Sabbah HN. Biologic rationale for the use of beta-blockers in the treatment of heart failure. *Heart Fail Rev* 2004;9:91-7.
- Grassi G, Seravalle G, Dell'Oro R, Mancia G. Sympathetic mechanisms, organ damage, and antihypertensive treatment. *Curr Hypertens Rep* 2011;13:303-8.
- Levy BI. How to explain the differences between renin angiotensin system modulators. *Am J Hypertens* 2005;18:1345-415.
- Polhemus DJ, Trivedi RK, Gao J, et al. Renal sympathetic denervation protects the failing heart via inhibition of neprilysin activity in the kidney. *J Am Coll Cardiol* 2017;70:2139-53.
- Koniaris K, Nikolaou M, Paraskevidis I, Parisis J. Therapeutic options for the management of the cardiorenal syndrome. *Int J Nephrol* 2010;2011:194910.
- Polhemus DJ, Lefer DJ. Emergence of hydrogen sulfide as an endogenous gaseous signaling molecule in cardiovascular disease. *Circ Res* 2014;114:730-7.
- Polhemus DJ, Calvert JW, Butler J, Lefer DJ. The cardioprotective actions of hydrogen sulfide in acute myocardial infarction and heart failure. *Scientifica* 2014;2014:768607.
- Predmore BL, Kondo K, Bhushan S, et al. The polysulfide diallyl trisulfide protects the ischemic myocardium by preservation of endogenous hydrogen sulfide and increasing nitric oxide bioavailability. *Am J Physiol Heart Circ Physiol* 2012;302:H2410-8.
- Polhemus D, Kondo K, Bhushan S, et al. Hydrogen sulfide attenuates cardiac dysfunction after heart failure via induction of angiogenesis. *Circ Heart Fail* 2013;6:1077-86.
- Kang J, Li Z, Organ CL, et al. pH-Controlled hydrogen sulfide release for myocardial ischemia-reperfusion injury. *J Am Chem Soc* 2016;138:6336-9.
- Predmore BL, Lefer DJ. Development of hydrogen sulfide-based therapeutics for cardiovascular disease. *J Cardiovasc Transl Res* 2010;3:487-98.
- Zhou Z, Martin E, Sharina I, et al. Regulation of soluble guanylyl cyclase redox state by hydrogen sulfide. *Pharmacol Res* 2016;111:556-62.
- Andreadou I, Iliodromitis EK, Rassaf T, Schulz R, Papapetropoulos A, Ferdinandy P. The role of gasotransmitters NO, H₂S and CO in myocardial ischaemia/reperfusion injury and cardioprotection by preconditioning, postconditioning and remote conditioning. *Br J Pharmacol* 2015;172:1587-606.
- Cao X, Bian JS. The role of hydrogen sulfide in renal system. *Front Pharmacol* 2016;7:385.

25. Kondo K, Bhushan S, King AL, et al. H₂S protects against pressure overload-induced heart failure via upregulation of endothelial nitric oxide synthase. *Circulation* 2013;127:1116-27.
26. Bradley JM, Li Z, Organ CL, et al. A novel mtDNA repair fusion protein attenuates maladaptive remodeling and preserves cardiac function in heart failure. *Am J Physiol Heart Circ Physiol* 2018; 314:H311-21.
27. Trivedi RK, Polhemus DJ, Li Z, et al. Combined angiotensin receptor-neprilysin inhibitors improve cardiac and vascular function via increased no bioavailability in heart failure. *J Am Heart Assoc* 2018 Mar 3;7(5). pii: e008268, <https://doi.org/10.1161/JAHA.117.008268>.
28. Sung MM, Byrne NJ, Robertson IM, et al. Resveratrol improves exercise performance and skeletal muscle oxidative capacity in heart failure. *Am J Physiol Heart Circ Physiol* 2017;312:H842-53.
29. Polhemus DJ, Li Z, Pattillo CB, et al. A novel hydrogen sulfide prodrug, SG1002, promotes hydrogen sulfide and nitric oxide bioavailability in heart failure patients. *Cardiovasc Ther* 2015;33: 216-26.
30. Shimizu Y, Nicholson CK, Lambert JP, et al. Sodium sulfide attenuates ischemic-induced heart failure by enhancing proteasomal function in an Nrf2-dependent manner. *Circ Heart Fail* 2016;9: e002368.
31. King AL, Polhemus DJ, Bhushan S, et al. Hydrogen sulfide cytoprotective signaling is endothelial nitric oxide synthase-nitric oxide dependent. *Proc Natl Acad Sci U S A* 2014;111: 3182-7.
32. Katsouda A, Bibli SI, Pyriochou A, Szabo C, Papapetropoulos A. Regulation and role of endogenously produced hydrogen sulfide in angiogenesis. *Pharmacol Res* 2016;113:175-85.
33. Bibli SI, Szabo C, Chatzianastasiou A, et al. Hydrogen sulfide preserves endothelial nitric oxide synthase function by inhibiting proline-rich kinase 2: implications for cardiomyocyte survival and cardioprotection. *Mol Pharmacol* 2017;92: 718-30.
34. Islam KN, Polhemus DJ, Donnarumma E, Brewster LP, Lefer DJ. Hydrogen sulfide levels and nuclear factor-erythroid 2-related factor 2 (NRF2) activity are attenuated in the setting of critical limb ischemia (CLI). *J Am Heart Assoc* 2015 May 14;4(5). pii: e001986.
35. Smilde TD, Hillege HL, Navis G, Boomsma F, de Zeeuw D, van Veldhuisen DJ. Impaired renal function in patients with ischemic and nonischemic chronic heart failure: association with neurohormonal activation and survival. *Am Heart J* 2004; 148:165-72.
36. Calvert JW, Jha S, Gundewar S, et al. Hydrogen sulfide mediates cardioprotection through Nrf2 signaling. *Circ Res* 2009;105: 365-74.
37. Meng G, Liu J, Liu S, et al. Hydrogen sulfide pretreatment improves mitochondrial function in myocardial hypertrophy via a SIRT3-dependent manner. *Br J Pharmacol* 2018;175:1126-45.
38. Li Z, Polhemus DJ, Lefer DJ. Evolution of hydrogen sulfide therapeutics to treat cardiovascular disease. *Circ Res* 2018;123.
39. Kamal FA, Travers JG, Schafer AE, Ma Q, Devarajan P, Blaxall BC. G protein-coupled receptor-G-protein betagamma-subunit signaling mediates renal dysfunction and fibrosis in heart failure. *J Am Soc Nephrol* 2017;28:197-208.
40. Gao L, Zimmerman MC, Biswal S, Zucker IH. Selective Nrf2 gene deletion in the rostral ventrolateral medulla evokes hypertension and sympathoexcitation in mice. *Hypertension* 2017; 69:1198-206.
41. Xue R, Hao DD, Sun JP, et al. Hydrogen sulfide treatment promotes glucose uptake by increasing insulin receptor sensitivity and ameliorates kidney lesions in type 2 diabetes. *Antioxid Redox Signal* 2013;19:5-23.
42. Wu D, Gao B, Li M, et al. Hydrogen sulfide mitigates kidney injury in high fat diet-induced obese mice. *Oxid Med Cell Longev* 2016;2016: 2715718.
43. Weber GJ, Pushpakumar SB, Sen U. Hydrogen sulfide alleviates hypertensive kidney dysfunction through an epigenetic mechanism. *Am J Physiol Heart Circ Physiol* 2017;312:H874-85.
44. John A, Kundu S, Pushpakumar S, et al. GYY4137, a hydrogen sulfide donor modulates miR194-dependent collagen realignment in diabetic kidney. *Sci Rep* 2017;7:10924.
45. Jiang D, Zhang Y, Yang M, Wang S, Jiang Z, Li Z. Exogenous hydrogen sulfide prevents kidney damage following unilateral ureteral obstruction. *NeuroUrol Urodyn* 2014;33:538-43.
46. Hsu CN, Tain YL. Hydrogen Sulfide in hypertension and kidney disease of developmental origins. *Int J Mol Sci* 2018 May 11;19(5). pii: E1438.
47. Hou CL, Wang MJ, Sun C, et al. Protective effects of hydrogen sulfide in the ageing kidney. *Oxid Med Cell Longev* 2016;2016:7570489.
48. Piepoli MF, Guazzi M, Boriani G, et al. Exercise intolerance in chronic heart failure: mechanisms and therapies. Part I. *Eur J Cardiovasc Prev Rehabil* 2010;17:637-42.
49. Piepoli MF, Guazzi M, Boriani G, et al. Exercise intolerance in chronic heart failure: mechanisms and therapies. Part II. *Eur J Cardiovasc Prev Rehabil* 2010;17:643-8.
50. Cheng Z, Garikipati VN, Nickoloff E, et al. Restoration of hydrogen sulfide production in diabetic mice improves reparative function of bone marrow cells. *Circulation* 2016;134: 1467-83.
51. Kanagy NL, Szabo C, Papapetropoulos A. Vascular biology of hydrogen sulfide. *Am J Physiol Cell Physiol* 2017;312:C537-49.
52. Esposito F, Mathieu-Costello O, Shabetai R, Wagner PD, Richardson RS. Limited maximal exercise capacity in patients with chronic heart failure: partitioning the contributors. *J Am Coll Cardiol* 2010;55:1945-54.
53. Marti CN, Gheorghide M, Kalogeropoulos AP, Georgiopoulou VV, Quyyumi AA, Butler J. Endothelial dysfunction, arterial stiffness, and heart failure. *J Am Coll Cardiol* 2012;60:1455-69.
54. Seilicovich A, Lasaga M, Befumo M, et al. Nitric oxide inhibits the release of norepinephrine and dopamine from the medial basal hypothalamus of the rat. *Proc Natl Acad Sci U S A* 1995;92: 11299-302.

KEY WORDS cardiorenal syndrome, exercise tolerance, heart failure, hydrogen sulfide, RAAS

APPENDIX For supplemental figures, please see the online version of this paper.