

Mutually Quadratically Invariant Information Structures in Two-Team Stochastic Dynamic Games

Marcello Colombino[†], Roy S. Smith[†], and Tyler H. Summers[‡]

Abstract— We formulate a two-team linear quadratic stochastic dynamic game featuring two opposing teams each with decentralized information structures. We introduce the concept of mutual quadratic invariance (MQI), which, analogously to quadratic invariance in (single team) decentralized control, defines a class of interacting information structures for the two teams under which optimal linear feedback control strategies are easy to compute. We show that, for zero-sum two-team dynamic games, structured state feedback Nash (saddle-point) equilibrium strategies can be computed from equivalent structured disturbance feedforward saddle point equilibrium strategies. However, for nonzero-sum games we show via a counterexample that a similar equivalence fails to hold. The results are illustrated with a simple yet rich numerical example that illustrates the importance of the information structure for dynamic games.

I. INTRODUCTION

Future cyber-physical systems (CPS) will feature cooperative networks of autonomous decision making agents equipped with embedded sensing, computation, communication, and actuation capabilities. These capabilities promise to significantly enhance performance, but also render the network vulnerable by increasing the number of access and influence points available to attackers. “Red team-blue team” scenarios, in which a defending team seeks to operate the network efficiently and securely while the attacking team seeks to disrupt network operation, have been used to qualitatively assess and improve security in military and intelligence organizations, but have not received formal mathematical analysis in a CPS context. Here we will study some fundamental properties in two-team stochastic dynamic games in cyber-physical networks, with a focus on interactions of the information structures of each team.

Dynamic game theory [1] offers a general framework for the study of optimal decision making in stochastic and non-cooperative environments. The theory can be viewed as a marriage of game theory [2], with a focus on interactions of multiple decision making agents, and optimal control theory [3], [4], with a focus on dynamics and feedback. The main elements of dynamic game theory are (1) a dynamical system along with a set of agents whose actions influence the state evolution of the system, (2) an objective function to be

optimized associated with each agent, and (3) an information structure that specifies information sets for each agent, i.e., who knows what and when.

Several special classes of dynamic games have been extensively studied. In team decision theory all agents cooperate to optimize the same objective function. Static team theory traces back to the seminal work of Marschak and Radner [5]–[7]. Decentralized control theory has developed from team decision theory and control theory and introduces dynamics. The presence of dynamics makes available information depend on the actions of agents and significantly complicates the problem. Dynamic aspects were studied in important early work by Witsenhausen [8]–[10] and Ho [11], [12]. Witsenhausen’s famous counterexample [8] vividly demonstrated the computational difficulties associated with team decision making in dynamic and stochastic environments. This still-unsolved counterexample described a simple team decision problem in which a nonlinear strategy strictly outperforms the optimal linear strategy and established deep connections between control, communication, and information theory. Research on decentralized and distributed control theory has continued, and there has been a recent resurgence of interest driven by the advent of large-scale cyber-physical networks. Recent work has elaborated on connections with communication and information theory [13], [14] and focused on computational and structural issues [15]–[20].

These important structural information aspects arising from cooperating agents have received much less attention in the dynamic game literature, which tends to focus only on non-cooperative and adversarial behavior. The most well studied case is the two-player problem, which features two opposing agents who have centralized information structures and has connections to robust control [21]. What is currently under explored is a comprehensive study of information structure aspects when there are *both* non-cooperative elements, as in general dynamic game theory, and cooperative elements, as in decentralized control theory. These aspects can be captured by a two-team stochastic dynamic game framework.

Two-team stochastic dynamic games feature two opposing teams with decentralized information structures for both the attacking and defending teams: each agent must act based on partial information measured or received locally in a way that coordinates its actions with team members and counters against the opposing team. This framework mathematically formalizes “red team-blue team” scenarios that qualitatively assess network security and resilience. In comparison to decentralized control theory, a team adversarial element is

[†]M. Colombino and R. Smith are with the Automatic Control Laboratory, ETH Zurich, Switzerland. [‡]T. Summers is with the Department of Mechanical Engineering, University of Texas at Dallas, USA. E-mail addresses: {mcolombi, rsmith}@control.ee.ethz.ch, tyler.summers@utdallas.edu. This research is supported by the National Science Foundation under grant CNS-1566127 and partially supported by the Swiss National Science Foundation grant 2-773337-12.

added. In comparison to general dynamic game theory, a sharp contrast between cooperation with teammates and conflict against adversaries is preserved. Further, the stochastic element (modeled by a “chance” or “Nature” player in the game) allows the inclusion of random component failures and disturbance signals.

There is currently a lack of deep theoretical and computational understanding in this class of dynamic games. A static version of the problem was studied in [22]. Many fundamental questions that been answered in static or single team decentralized control settings do not have counterparts in the two-team setting.

The main contributions of the present paper are as follows. We formulate a two-team stochastic dynamic game problem and introduce a concept of mutual quadratic invariance, which defines a class of interacting information structures for the two teams under which optimal linear feedback control strategies are easy to compute. This is analogous to the concept of quadratic invariance in (single team) decentralized control [15]. We show that for zero-sum two-team dynamic games, structured state feedback saddle point equilibrium strategies can be computed from equivalent structured disturbance feedforward saddle point equilibrium strategies. However, for nonzero-sum games we show via a counterexample that a similar equivalence fails to hold for structured Nash equilibrium strategies. Finally, we present a numerical example, which illustrates the importance of the information structure on the value of the game.

The rest of the paper is structured as follows. Section II provides preliminaries on static team games. Section III formulates a two-team stochastic dynamic game. Sections IV and V develop results on disturbance feedforward and state feedback strategies and introduce the concept of mutual quadratic invariance. Section VI presents illustrative numerical experiments, and Section VII gives some concluding remarks and future research directions.

II. TWO-TEAM STOCHASTIC STATIC GAMES

In this section we review basic results for a two-team stochastic static game. In this setting, two teams who both know the distribution parameters of a Gaussian random vector w need to decide strategies to compute vectors $u \in \mathbb{R}^m$ and $v \in \mathbb{R}^q$ as a function of the realization $w \in \mathbb{R}^n$ in order to minimize the expectation of *different* quadratic forms in w, u, v . Each team is composed of multiple agents, each of which observes a different linear function of w and decides a portion of the vectors u or v .

More formally, given a vector $w \sim \mathcal{N}(m_w, \Sigma_w)$, where $\Sigma_w \succ 0$, consider the following game

$$T_1 : \begin{cases} \min_{\kappa_i(\cdot)} \mathbb{E}_w (J_1(w, u, v)) \\ \text{s. t. } u_i = \kappa_i(C_i w) \\ \forall i \in \mathbb{Z}_{[1, N]} \end{cases}, \quad T_2 : \begin{cases} \min_{\lambda_j(\cdot)} \mathbb{E}_w (J_2(w, u, v)) \\ \text{s. t. } v_j = \lambda_j(\Gamma_j w) \\ \forall j \in \mathbb{Z}_{[1, M]} \end{cases} \quad (1)$$

where $J_i(w, u, v) :=$

$$\begin{bmatrix} w \\ u \\ v \end{bmatrix}^\top \begin{bmatrix} \mathcal{H}_{i \ w} & \mathcal{H}_{i \ wu} & \mathcal{H}_{i \ wv} \\ \mathcal{H}_{i \ wu}^\top & \mathcal{H}_{i \ uu} & \mathcal{H}_{i \ uv} \\ \mathcal{H}_{i \ wv}^\top & \mathcal{H}_{i \ uv}^\top & \mathcal{H}_{i \ vv} \end{bmatrix} \begin{bmatrix} w \\ u \\ v \end{bmatrix}, \quad i \in \{1, 2\}$$

are the objective functions of each team, and $\kappa_i(\cdot), i \in \mathbb{Z}_{[1, N]}$ and $\lambda_j(\cdot), j \in \mathbb{Z}_{[1, M]}$ are Borel measurable functions corresponding to the decision strategies of agents on team 1 and 2, respectively.

Assumption 1: We assume

$$\begin{bmatrix} \mathcal{H}_{1 \ uu} & \mathcal{H}_{1 \ uv} \\ \mathcal{H}_{1 \ uv}^\top & \mathcal{H}_{1 \ vv} \end{bmatrix} \succ 0, \quad \begin{bmatrix} \mathcal{H}_{1 \ uu} & \mathcal{H}_{2 \ uv} \\ \mathcal{H}_{2 \ uv}^\top & \mathcal{H}_{2 \ vv} \end{bmatrix} \succ 0.$$

Note that in the zero-sum case ($J_1 = -J_2$), Assumption 1 is standard to guarantee the existence of a saddle point equilibrium to the game without decentralized information structure [23, condition 6.3.9]. If $J_1 = J_2$, Assumption 1 reduces to the standard positive definite assumption of team theory [6].

We now define the set of Nash optimal strategies for the game in (1).

Definition 1: A pair of strategies $(\kappa^*(\cdot), \lambda^*(\cdot))$ of the form $[\kappa_1^{*\top}(C_1 \cdot), \dots, \kappa_N^{*\top}(C_n \cdot)]^\top$ and $[\lambda_1^{*\top}(\Gamma_1 \cdot), \dots, \lambda_M^{*\top}(\Gamma_n \cdot)]^\top$ is Nash optimal for the game in (1) if

$$\begin{cases} \kappa^*(\cdot) \in \arg \min_{\kappa(\cdot)} \mathbb{E}_w J_1(w, \lambda^*(w), \kappa(w)) \\ \lambda^*(\cdot) \in \arg \min_{\lambda(\cdot)} \mathbb{E}_w J_2(w, \lambda(w), \kappa^*(w)). \end{cases} \quad (2)$$

Under Assumption 1, the game in (1) admits a unique set of linear Nash optimal strategies, which can be computed by solving a set of linear equations derived from stationarity conditions [22]. This turns out to be a special case of a general multi-player, multi-objective linear quadratic static game considered in [24].

III. TWO-TEAM STOCHASTIC DYNAMIC GAMES

Problem (1) is a static game. There is no concept of time and causality of the information pattern. In this section we formulate a dynamic game, where two teams can influence the state evolution of a dynamical system. The agents on each team decide a portion of an input signal based on different observations of the system state over time. Decision must be causal: each player is only allowed to use past or, at most, present information.

Our focus will be on the role of information structures for both teams in determining equilibrium strategies. Dynamic games offer a rich variety of information structures. Specific instances have been considered in the literature, with much work on various types of centralized structures [1] and some work on structures defined by spatiotemporal decentralization patterns. For example, a one-step-delay observation sharing pattern was shown in [24] to admit unique linear optimal strategies. There has been recent progress in (single team) decentralized control on information structure issues, including a characterization of information structures called quadratically invariant that yield convex control design

problems [15]. Here we seek an analogous result in a two-team game setting.

Consider the system

$$x(t+1) = Ax(t) + B_1u(t) + B_2v(t) + w(t), \quad (3)$$

where $x(t) \in \mathbb{R}^n$ is the system state at time t with $x(0) \sim \mathcal{N}(0, \Sigma_0)$, $u(t) \in \mathbb{R}^{m_1}$ is the input for team 1 at time t , $v(t) \in \mathbb{R}^{m_2}$ is the input for team 2 at time t , and $w(t) \sim \mathcal{N}(0, \Sigma_t)$ is a random disturbance. The cost functions for each team are given by

$$J_i := \mathbb{E} \left(\sum_{t=0}^{N-1} x(t)^\top M_i(t)x(t) + u(t)^\top R_i(t)u(t) + v(t)^\top V_i(t)v(t) \right) + x(N)^\top M_i(N)x(N), \quad i \in \{1, 2\}, \quad (4)$$

where $M_i(0) = \mathbf{0}_{n \times n}$, $M_i(t) = M_i(t)^\top \in \mathbb{R}^{n \times n}$, $R_i(t) = R_i(t)^\top \in \mathbb{R}^{m_1 \times m_1}$ and $V_i(t) = V_i(t)^\top \in \mathbb{R}^{m_2 \times m_2}$. By defining the matrices $\mathcal{A} = \text{blockdiag}(A, \dots, A) \in \mathbb{R}^{n(N+1) \times n(N+1)}$,

$$B_i = \begin{bmatrix} B_i & 0 & 0 \\ 0 & \ddots & 0 \\ \vdots & & B_i \\ 0 & \dots & 0 \end{bmatrix} \in \mathbb{R}^{n(N+1) \times m_j N}, \quad i \in \{1, 2\},$$

$\mathcal{M}_i = \text{blockdiag}(0, M_i(1), \dots, M_i(N)) \in \mathbb{R}^{n(N+1) \times n(N+1)}$, $\mathcal{R}_i = \text{blockdiag}(R_i(0), \dots, R_i(N-1)) \in \mathbb{R}^{m_1 N \times m_1 N}$ and $\mathcal{V}_i = \text{blockdiag}(V_i(0), \dots, V_i(N-1)) \in \mathbb{R}^{m_2 N \times m_2 N}$ for $i \in \{1, 2\}$, the vectors $\mathbf{x} = (x(0), \dots, x(N)) \in \mathbb{R}^{n(N+1)}$, $\mathbf{u} = (u(0), \dots, u(N-1)) \in \mathbb{R}^{m_1 N}$, $\mathbf{v} = (v(0), \dots, v(N-1)) \in \mathbb{R}^{m_2 N}$ and $\mathbf{w} = (x(0), w(0), \dots, w(N-1)) \in \mathbb{R}^{n(N+1)}$, and the shift matrix

$$\mathcal{Z} := \begin{bmatrix} 0 & & & & \\ I & \ddots & & & \\ & \ddots & \ddots & & \\ & & \ddots & \ddots & \\ & & & I & 0 \end{bmatrix} \in \mathbb{R}^{n(N+1) \times n(N+1)},$$

we can write system (3) as

$$\mathbf{x} = \mathcal{Z}\mathcal{A}\mathbf{x} + \mathcal{Z}\mathcal{B}_1\mathbf{u} + \mathcal{Z}\mathcal{B}_2\mathbf{v} + \mathbf{w}. \quad (5)$$

The system in (5) can be rewritten compactly as

$$\mathbf{x} = \begin{bmatrix} \mathcal{P}_{11} & \mathcal{P}_{12} & \mathcal{P}_{13} \end{bmatrix} \begin{bmatrix} \mathbf{w} \\ \mathbf{u} \\ \mathbf{v} \end{bmatrix}, \quad (6)$$

where $\mathcal{P}_{11} = (I - \mathcal{Z}\mathcal{A})^{-1}$, $\mathcal{P}_{12} = (I - \mathcal{Z}\mathcal{A})^{-1}\mathcal{Z}\mathcal{B}_1$ and $\mathcal{P}_{13} = (I - \mathcal{Z}\mathcal{A})^{-1}\mathcal{Z}\mathcal{B}_2$. The cost functions in (4) can be written in function of the vectorized inputs as

$$\mathbf{J}_i(\mathbf{u}, \mathbf{v}) = \mathbb{E}_{\mathbf{w}} \left(\begin{bmatrix} \mathbf{w} \\ \mathbf{u} \\ \mathbf{v} \end{bmatrix}^\top \mathcal{H}_i \begin{bmatrix} \mathbf{w} \\ \mathbf{u} \\ \mathbf{v} \end{bmatrix} \right), \quad i \in \{1, 2\},$$

where

$$\mathcal{H}_i = \begin{bmatrix} \mathcal{P}_{11}^\top \mathcal{M}_i \mathcal{P}_{11} & \mathcal{P}_{11}^\top \mathcal{M}_i \mathcal{P}_{12} & \mathcal{P}_{11}^\top \mathcal{M}_i \mathcal{P}_{13} \\ \mathcal{P}_{12}^\top \mathcal{M}_i \mathcal{P}_{11} & \mathcal{P}_{12}^\top \mathcal{M}_i \mathcal{P}_{12} + \mathcal{R}_i & \mathcal{P}_{12}^\top \mathcal{M}_i \mathcal{P}_{13} \\ \mathcal{P}_{13}^\top \mathcal{M}_i \mathcal{P}_{11} & \mathcal{P}_{13}^\top \mathcal{M}_i \mathcal{P}_{12} & \mathcal{P}_{13}^\top \mathcal{M}_i \mathcal{P}_{13} + \mathcal{V}_i \end{bmatrix},$$

for $i \in \{1, 2\}$.

We are interested in the finite horizon, two-team stochastic dynamic game where:

- Team 1 minimizes $\mathbf{J}_1(\mathbf{u}, \mathbf{v})$
- Team 2 minimizes $\mathbf{J}_2(\mathbf{u}, \mathbf{v})$
- Each team choses a structured causal state feedback strategy of the form

$$\mathbf{u} = \mathcal{K}_1(\mathbf{x}), \quad \mathbf{v} = \mathcal{K}_2(\mathbf{x}). \quad \mathcal{K}_1 \in \mathcal{S}_1, \quad \mathcal{K}_2 \in \mathcal{S}_2,$$

where $\mathcal{K}_i : \mathbb{R}^{n(N+1)} \rightarrow \mathbb{R}^{m_i N}$, for $i \in \{1, 2\}$ are measurable functions and \mathcal{S}_1 and \mathcal{S}_2 define an information structure for each team.

We define an information structure $\mathcal{S}_i \in \{0, 1\}^{n(N+1) \times m_i N}$ as a binary matrix. $\mathcal{K}_i \in \mathcal{S}_i$ indicates that, if $[\mathcal{S}_i]_{jk} = 0$, then the j^{th} element of \mathcal{K}_i is not a function of \mathbf{x}_k . By choosing the information structures one can enforce causality and a prescribed spatiotemporal structure on the controller strategies.

IV. MUTUAL QUADRATIC INVARIANCE

In decentralized control with quadratically invariant information structures, the controller structure can be enforced on an affine parameter that defines the achievable set of closed-loop systems and recover a structured feedback controller. We now follow a similar approach in the two-team setting.

A. Disturbance feedforward strategies

By searching for measurable disturbance feedforward strategies of the type $\mathbf{u} = \mathcal{Q}_1(\mathcal{P}_{11}\mathbf{w})$ and $\mathbf{v} = \mathcal{Q}_2(\mathcal{P}_{11}\mathbf{w})$, where $\mathcal{Q}_1 \in \mathcal{S}_1$ and $\mathcal{Q}_2 \in \mathcal{S}_2$, we recover the formulation of (1). Provided that Assumption 1 is satisfied, there exists a unique Nash equilibrium of the form $\mathbf{u} = \bar{\mathcal{Q}}_1\mathcal{P}_{11}\mathbf{w}$, $\mathbf{v} = \bar{\mathcal{Q}}_2\mathcal{P}_{11}\mathbf{w}$ in the space of linear strategies [22]. The matrices $\bar{\mathcal{Q}}_1, \bar{\mathcal{Q}}_2$ can be easily computed by solving a linear system of equations or a sequence of semidefinite programs [22], [24]. Assumption 1 for the dynamic game problem becomes

$$\begin{bmatrix} \mathcal{P}_{12}^\top \mathcal{M}_1 \mathcal{P}_{12} + \mathcal{R}_1 & \mathcal{P}_{12}^\top \mathcal{M}_1 \mathcal{P}_{13} \\ \mathcal{P}_{13}^\top \mathcal{M}_1 \mathcal{P}_{12} & \mathcal{P}_{13}^\top \mathcal{M}_1 \mathcal{P}_{13} + \mathcal{S}_2 \end{bmatrix} \succ 0, \\ \begin{bmatrix} \mathcal{P}_{12}^\top \mathcal{M}_2 \mathcal{P}_{12} + \mathcal{R}_2 & \mathcal{P}_{12}^\top \mathcal{M}_2 \mathcal{P}_{13} \\ \mathcal{P}_{13}^\top \mathcal{M}_2 \mathcal{P}_{12} & \mathcal{P}_{13}^\top \mathcal{M}_2 \mathcal{P}_{13} + \mathcal{S}_1 \end{bmatrix} \succ 0.$$

We can define new cost functions that depend on the matrices \mathcal{Q}_1 and \mathcal{Q}_2 describing linear disturbance feedforward strategies as

$$\mathcal{J}_i \left(\begin{bmatrix} \mathcal{Q}_1 \\ \mathcal{Q}_2 \end{bmatrix} \right) = \mathbf{J}_i(\mathbf{u}, \mathbf{v}) \Big|_{\mathbf{u}=\mathcal{Q}_1\mathcal{P}_{11}\mathbf{w}, \mathbf{v}=\mathcal{Q}_2\mathcal{P}_{11}\mathbf{w}}. \quad (7)$$

In particular,

$$\begin{aligned} \mathcal{J}_i \left(\begin{bmatrix} \mathcal{Q}_1 \\ \mathcal{Q}_2 \end{bmatrix} \right) &= \|\mathcal{M}_i^{\frac{1}{2}} (I + \mathcal{P}_{12}\mathcal{Q}_1 + \mathcal{P}_{13}\mathcal{Q}_2) \mathcal{P}_{11}\Sigma_{\mathbf{w}}^{\frac{1}{2}}\|_F^2 \\ &\quad + \|\mathcal{R}_i^{\frac{1}{2}}\mathcal{Q}_1\Sigma_{\mathbf{w}}^{\frac{1}{2}}\|_F^2 + \|\mathcal{V}_i^{\frac{1}{2}}\mathcal{Q}_2\Sigma_{\mathbf{w}}^{\frac{1}{2}}\|_F^2, \end{aligned}$$

where $\Sigma_{\mathbf{w}}^{\frac{1}{2}}$ is the covariance of \mathbf{w} .

B. Equivalent state feedback strategies

It is easy to show that there exists a bijective relationship between a pair of linear disturbance feedforward strategies $(\mathcal{Q}_1, \mathcal{Q}_2)$ and an equivalent pair of linear state feedback strategies described by the matrices $(\mathcal{K}_1, \mathcal{K}_2)$. More precisely, using (6) we obtain

$$\begin{aligned} \begin{bmatrix} \mathbf{u} \\ \mathbf{v} \end{bmatrix} &= \begin{bmatrix} \mathcal{Q}_1 \\ \mathcal{Q}_2 \end{bmatrix} \mathcal{P}_{11}\mathbf{w} \\ &= \begin{bmatrix} \mathcal{Q}_1 \\ \mathcal{Q}_2 \end{bmatrix} \mathbf{x} - \begin{bmatrix} \mathcal{Q}_1 \\ \mathcal{Q}_2 \end{bmatrix} \begin{bmatrix} \mathcal{P}_{12} & \mathcal{P}_{13} \end{bmatrix} \begin{bmatrix} \mathbf{u} \\ \mathbf{v} \end{bmatrix}. \end{aligned} \quad (8)$$

We can define the function g such that

$$\begin{pmatrix} \mathcal{K}_1 \\ \mathcal{K}_2 \end{pmatrix} = g \left(\begin{bmatrix} \mathcal{Q}_1 \\ \mathcal{Q}_2 \end{bmatrix} \right),$$

where

$$g \left(\begin{bmatrix} \mathcal{Q}_1 \\ \mathcal{Q}_2 \end{bmatrix} \right) = \left(I + \begin{bmatrix} \mathcal{Q}_1 \\ \mathcal{Q}_2 \end{bmatrix} \begin{bmatrix} \mathcal{P}_{12} & \mathcal{P}_{13} \end{bmatrix} \right)^{-1} \begin{bmatrix} \mathcal{Q}_1 \\ \mathcal{Q}_2 \end{bmatrix}.$$

Using a similar approach to (8), one can construct the inverse mapping that, given a pair of feedback strategies, recovers the equivalent feedforward strategies.

$$\begin{bmatrix} \mathcal{Q}_1 \\ \mathcal{Q}_2 \end{bmatrix} = g^{-1} \left(\begin{bmatrix} \mathcal{K}_1 \\ \mathcal{K}_2 \end{bmatrix} \right),$$

where the map g^{-1} takes the form

$$\begin{aligned} g^{-1} \left(\begin{bmatrix} \mathcal{K}_1 \\ \mathcal{K}_2 \end{bmatrix} \right) &= \\ &= \begin{bmatrix} \mathcal{K}_1 \\ \mathcal{K}_2 \end{bmatrix} \left(I - \begin{bmatrix} \mathcal{P}_{12} & \mathcal{P}_{13} \end{bmatrix} \begin{bmatrix} \mathcal{K}_1 \\ \mathcal{K}_2 \end{bmatrix} \right)^{-1}. \end{aligned}$$

Given a pair of linear feedback strategies \mathcal{K}_1 and \mathcal{K}_2 , the cost for player i can be evaluated by considering the equivalent feedforward strategies as

$$\mathbf{J}_i(\mathbf{u}, \mathbf{v}) \Big|_{\mathbf{u}=\mathcal{K}_1\mathbf{x}, \mathbf{v}=\mathcal{K}_2\mathbf{x}} = \mathcal{J}_i \left(g^{-1} \left(\begin{bmatrix} \mathcal{K}_1 \\ \mathcal{K}_2 \end{bmatrix} \right) \right),$$

where \mathcal{J}_i is defined in (7).

Now that we have a way to construct feedback strategies which are equivalent to any set of linear feedforward strategies, we need to establish a condition that guarantees that such equivalent feedback strategies will preserve the desired structure.

C. Mutual Quadratic Invariance

We know from the quadratic invariance literature [15], [25] that $\mathcal{Q}_1 \in \mathcal{S}_1$ and $\mathcal{Q}_2 \in \mathcal{S}_2 \iff \mathcal{K}_1 \in \mathcal{S}_1$ and $\mathcal{K}_2 \in \mathcal{S}_2$ if and only if for all $(\mathcal{K}_1, \mathcal{K}_2) \in \mathcal{S}_1 \times \mathcal{S}_2$ it holds that

$$\begin{bmatrix} \mathcal{K}_1 \\ \mathcal{K}_2 \end{bmatrix} \begin{bmatrix} \mathcal{P}_{12} & \mathcal{P}_{13} \end{bmatrix} \begin{bmatrix} \mathcal{K}_1 \\ \mathcal{K}_2 \end{bmatrix} \in \mathcal{S}_1 \times \mathcal{S}_2, \quad (9)$$

in other words $\mathcal{S}_1 \times \mathcal{S}_2$ is quadratically invariant under $\begin{bmatrix} \mathcal{P}_{12} & \mathcal{P}_{13} \end{bmatrix}$. We define this property as **mutual quadratic invariance**. We can expand (9) as

$$\begin{bmatrix} \mathcal{K}_1 \\ \mathcal{K}_2 \end{bmatrix} \in \mathcal{S}_1 \times \mathcal{S}_2 \implies \begin{cases} \mathcal{K}_1\mathcal{P}_{12}\mathcal{K}_1, & \mathcal{K}_1\mathcal{P}_{13}\mathcal{K}_2 \in \mathcal{S}_1 \\ \mathcal{K}_2\mathcal{P}_{12}\mathcal{K}_1, & \mathcal{K}_2\mathcal{P}_{13}\mathcal{K}_2 \in \mathcal{S}_2. \end{cases} \quad (10)$$

By observing (9) we note that MQI is equivalent to QI for a control problem where both decisions \mathbf{u} and \mathbf{v} are taken by a single decision maker. MQI information structures will allow us to compute equilibrium strategies in two-team games.

V. COMPUTING EQUILIBRIUM STRATEGIES

We have seen in Section IV that given structures \mathcal{S}_1 and \mathcal{S}_2 which are mutually quadratically invariant under \mathcal{P}_{12} and \mathcal{P}_{13} , one can easily obtain a Nash equilibrium in the disturbance feedforward strategies. Furthermore once we recover the equivalent state feedback strategies the structure is preserved. There is still a nontrivial question that needs to be answered.

Problem 1: Given a Nash equilibrium in the feedforward strategies $(\bar{\mathcal{Q}}_1, \bar{\mathcal{Q}}_2)$, are the equivalent feedback strategies

$$\begin{pmatrix} \bar{\mathcal{K}}_1 \\ \bar{\mathcal{K}}_2 \end{pmatrix} = g \left(\begin{bmatrix} \bar{\mathcal{Q}}_1 \\ \bar{\mathcal{Q}}_2 \end{bmatrix} \right), \quad (11)$$

a Nash equilibrium in the feedback strategies?

In order to understand why the answer to Problem 1 is nontrivial, we first present a counterexample for a nonzero sum game.

A. Nonzero sum game

Consider the following problem instance with

$$\begin{aligned} N &= 2, & A &= 2, & B_1 &= 0.4, & B_2 &= 0.1, \\ \Sigma_0 &= 1, & \Sigma_t &= 1 \quad \forall t, \\ M_1(t) &= R_1(t) = S_1(t) = 1 \quad \forall t, \\ M_2(t) &= 70 \quad \forall t, R_2(t) = S_2(t) = 1 \quad \forall t. \end{aligned} \quad (12)$$

This is a single state, two-stage, two-player problem, and we will consider centralized, causal strategies, which are readily verified to be mutually quadratically invariant. Using disturbance feedforward strategies, the problem can be reduced to a static game whose unique Nash Equilibrium strategies are readily computed using methods from [22], [24]. This yields the Nash pair $(\bar{\mathcal{Q}}_1, \bar{\mathcal{Q}}_2)$, where

$$\begin{aligned} \bar{\mathcal{Q}}_1 &= \begin{bmatrix} -0.6795 & 0 & 0 \\ 0.6283 & -0.4301 & 0 \end{bmatrix}, \\ \bar{\mathcal{Q}}_2 &= \begin{bmatrix} -11.890 & 0 & 0 \\ 10.996 & -7.5269 & 0 \end{bmatrix}. \end{aligned} \quad (13)$$

The corresponding equilibrium value for player 1 is $\mathcal{J}_1^* \left(\begin{bmatrix} \bar{Q}_1 \\ \bar{Q}_2 \end{bmatrix} \right) = 220$. The corresponding state feedback strategies are

$$\begin{aligned} \bar{\mathcal{K}}_1 &= \begin{bmatrix} -0.6795 & 0 & 0 \\ 0 & -0.4301 & 0 \end{bmatrix}, \\ \bar{\mathcal{K}}_2 &= \begin{bmatrix} -11.890 & 0 & 0 \\ 0 & -7.5269 & 0 \end{bmatrix}, \end{aligned} \quad (14)$$

However, $(\bar{\mathcal{K}}_1, \bar{\mathcal{K}}_2)$ is not a Nash Equilibrium in state feedback strategies since it is readily verified that $\mathcal{J}_1 \left(g^{-1} \left(\begin{bmatrix} \hat{\mathcal{K}}_1 \\ \hat{\mathcal{K}}_2 \end{bmatrix} \right) \right) = 206.1$, where

$$\hat{\mathcal{K}}_1 = \begin{bmatrix} -1.853 & 0 & 0 \\ 0 & -0.4301 & 0 \end{bmatrix}. \quad (15)$$

Although the sparsity structure is preserved between the disturbance feedforward and state feedback strategies since the information structure is mutually quadratically invariant, the Nash Equilibrium property is not preserved. Thus, for this particular non-zero sum dynamic game, the answer to the question posed in Problem 1 is negative. We show next that this situation does not occur in zero-sum games.

B. Zero sum game

Let us now consider the zero sum game case, where the objective function of one team is precisely the negative of that of the other team. Such problems can be re-written as a min-max problem of the form.

$$\min_{\mathbf{u}} \max_{\mathbf{v}} \mathbf{J}(\mathbf{u}, \mathbf{v}) := \mathbb{E}_{\mathbf{w}} \left(\begin{bmatrix} \mathbf{w} \\ \mathbf{u} \\ \mathbf{v} \end{bmatrix}^\top \mathcal{H} \begin{bmatrix} \mathbf{w} \\ \mathbf{u} \\ \mathbf{v} \end{bmatrix} \right) \quad (16)$$

As before we are interested in strategies of the form

$$\mathbf{u} = \mathcal{K}_1(\mathbf{x}), \quad \mathbf{v} = \mathcal{K}_2(\mathbf{x}). \quad \mathcal{K}_1 \in \mathcal{S}_1, \mathcal{K}_2 \in \mathcal{S}_2,$$

where \mathcal{S}_1 and \mathcal{S}_2 are prescribed sets of structured causal controllers. As zero-sum games are a special case of nonzero sum games, provided Assumption 1 holds, we can find structured Nash equilibria (rather saddle point equilibrium in the zero sum context) if we consider linear strategies of the form $\mathbf{u} = Q_1 \mathcal{P}_{11} \mathbf{w}$ and $\mathbf{v} = Q_2 \mathcal{P}_{11} \mathbf{w}$. For the zero sum game of the form (16), Assumption 1 reads

$$\begin{bmatrix} \mathcal{P}_{12}^\top \mathcal{M} \mathcal{P}_{12} + \mathcal{R} & \mathcal{P}_{12}^\top \mathcal{M} \mathcal{P}_{13} \\ \mathcal{P}_{13}^\top \mathcal{M} \mathcal{P}_{12} & -(\mathcal{P}_{13}^\top \mathcal{M} \mathcal{P}_{13} + \mathcal{S}) \end{bmatrix} \succ 0.$$

As before, given the saddle point equilibrium in the feedforward strategies, when \mathcal{S}_1 and \mathcal{S}_2 are mutually quadratically invariant, we can compute equivalent linear feedback strategies that preserve the structure. In the zero-sum case, however, we can relate the equilibrium property of the feedforward strategies to that of the state feedback strategies.

We start with a result that shows that the maps g and g^{-1} preserve stationary points. We begin by noting that for a zero-sum game $\mathcal{J}_1 = -\mathcal{J}_2$.

Lemma 1: Given

$$\begin{bmatrix} \bar{Q}_1 \\ \bar{Q}_2 \end{bmatrix} = g^{-1} \left(\begin{bmatrix} \bar{\mathcal{K}}_1 \\ \bar{\mathcal{K}}_2 \end{bmatrix} \right)$$

then

$$\begin{aligned} \frac{\partial \mathcal{J}_1 \left(\begin{bmatrix} Q_1 \\ Q_2 \end{bmatrix} \right)}{\partial Q_1} \Big|_{Q_1=\bar{Q}_1, Q_2=\bar{Q}_2} &\in \mathcal{S}_1^\perp, \\ \frac{\partial \mathcal{J}_2 \left(\begin{bmatrix} Q_1 \\ Q_2 \end{bmatrix} \right)}{\partial Q_2} \Big|_{Q_1=\bar{Q}_1, Q_2=\bar{Q}_2} &\in \mathcal{S}_2^\perp, \end{aligned} \quad (17)$$

if and only if

$$\begin{aligned} \frac{\partial \mathcal{J}_1 \left(g^{-1} \left(\begin{bmatrix} \mathcal{K}_1 \\ \mathcal{K}_2 \end{bmatrix} \right) \right)}{\partial \mathcal{K}_1} \Big|_{\mathcal{K}_1=\bar{\mathcal{K}}_1, \mathcal{K}_2=\bar{\mathcal{K}}_2} &\in \mathcal{S}_1^\perp, \\ \frac{\partial \mathcal{J}_2 \left(g^{-1} \left(\begin{bmatrix} \mathcal{K}_1 \\ \mathcal{K}_2 \end{bmatrix} \right) \right)}{\partial \mathcal{K}_2} \Big|_{\mathcal{K}_1=\bar{\mathcal{K}}_1, \mathcal{K}_2=\bar{\mathcal{K}}_2} &\in \mathcal{S}_2^\perp, \end{aligned} \quad (18)$$

Proof: Let us simplify the notation and define

$$Q := \begin{bmatrix} Q_1 \\ Q_2 \end{bmatrix}, \quad \mathcal{K} := \begin{bmatrix} \mathcal{K}_1 \\ \mathcal{K}_2 \end{bmatrix}, \quad \mathcal{P} := \begin{bmatrix} \mathcal{P}_{12} & \mathcal{P}_{13} \end{bmatrix}.$$

We start by proving the *only if* part. Note that since $\mathcal{J}_1 = -\mathcal{J}_2$,

$$\frac{\partial \mathcal{J}_2 \left(\begin{bmatrix} Q_1 \\ Q_2 \end{bmatrix} \right)}{\partial Q_2} \Big|_{Q_1=\bar{Q}_1, Q_2=\bar{Q}_2} \in \mathcal{S}_2^\perp$$

if and only if

$$\frac{\partial \mathcal{J}_1 \left(\begin{bmatrix} Q_1 \\ Q_2 \end{bmatrix} \right)}{\partial Q_2} \Big|_{Q_1=\bar{Q}_1, Q_2=\bar{Q}_2} \in \mathcal{S}_2^\perp.$$

Assume (18) holds, and suppose (17) does not hold. Then there exist $\tilde{Q} \in \mathcal{S}_1 \times \mathcal{S}_2$, with $\tilde{Q} \neq 0$ such that

$$\lim_{\varepsilon \rightarrow 0} \frac{\mathcal{J}_1(\tilde{Q} + \varepsilon \tilde{Q}) - \mathcal{J}(\tilde{Q})}{\varepsilon} = \kappa \neq 0,$$

or equivalently

$$\lim_{\varepsilon \rightarrow 0} \frac{\mathcal{J}_1 \left(g^{-1} \left(g \left(\tilde{Q} + \varepsilon \tilde{Q} \right) \right) \right) - \mathcal{J}(\tilde{Q})}{\varepsilon} = \kappa \neq 0. \quad (19)$$

We know that

$$\begin{aligned} g \left(\tilde{Q} + \varepsilon \tilde{Q} \right) &= \left(I + \left(\tilde{Q} + \varepsilon \tilde{Q} \right) \mathcal{P} \right)^{-1} \left(\tilde{Q} + \varepsilon \tilde{Q} \right) \\ &= \left(I + \tilde{Q} \mathcal{P} + \varepsilon \tilde{Q} \mathcal{P} \right)^{-1} \left(\tilde{Q} + \varepsilon \tilde{Q} \right) \\ &= \left[\left(I + \tilde{Q} \mathcal{P} \right)^{-1} + \varepsilon \left(I + \tilde{Q} \mathcal{P} \right)^{-1} \tilde{Q} \mathcal{P} \left(I + \tilde{Q} \mathcal{P} \right)^{-1} \right. \\ &\quad \left. + \mathcal{O}(\varepsilon^2) \right] \left(\tilde{Q} + \varepsilon \tilde{Q} \right) \\ &= \bar{\mathcal{K}} + \varepsilon \tilde{\mathcal{K}} + \mathcal{O}(\varepsilon^2), \end{aligned} \quad (20)$$

where

$$\tilde{\mathcal{K}} = \left(I + \tilde{Q} \mathcal{P} \right)^{-1} \tilde{Q} + \left(I + \tilde{Q} \mathcal{P} \right)^{-1} \tilde{Q} \mathcal{P} \left(I + \tilde{Q} \mathcal{P} \right)^{-1} \tilde{Q}.$$

- [10] H. Witsenhausen, "On information structures, feedback and causality," *SIAM Journal on Control*, vol. 9, no. 2, pp. 149–160, 1971.
- [11] Y. C. Ho and K.-C. Chu, "Team decision theory and information structures in optimal control problems—part i," *IEEE Transactions on Automatic Control*, vol. 17, no. 1, pp. 15–22, 1972.
- [12] Y.-C. Ho, "Team decision theory and information structures," *Proceedings of the IEEE*, vol. 68, no. 6, pp. 644–654, 1980.
- [13] P. Grover, S. Y. Park, and A. Sahai, "Approximately optimal solutions to the finite-dimensional Witsenhausen counterexample," *IEEE Transactions on Automatic Control*, vol. 58, no. 9, pp. 2189–2204, 2013.
- [14] P. Grover, A. B. Wagner, and A. Sahai, "Information embedding and the triple role of control," *IEEE Transactions on Information Theory*, vol. 61, no. 4, pp. 1539–1549, 2015.
- [15] M. Rotkowitz and S. Lall, "A characterization of convex problems in decentralized control," *Automatic Control, IEEE Transactions on*, vol. 51, no. 2, pp. 274–286, 2006.
- [16] A. Nayyar, A. Mahajan, and D. Teneketzis, "Decentralized stochastic control with partial history sharing: A common information approach," *IEEE Transactions on Automatic Control*, vol. 58, no. 7, pp. 1644–1658, 2013.
- [17] S. Yüksel and T. Başar, "Stochastic networked control systems," *AMC*, vol. 10, p. 12, 2013.
- [18] J. Swigart and S. Lall, "Optimal controller synthesis for decentralized systems over graphs via spectral factorization," *IEEE Transactions on Automatic Control*, vol. 59, no. 9, pp. 2311–2323, 2014.
- [19] L. Lessard and S. Lall, "Optimal control of two-player systems with output feedback. to appear," *IEEE Transactions on Automatic Control*, vol. 60, no. 8, pp. 2129 – 2144, 2015.
- [20] A. Gattami, B. M. Bernhardsson, and A. Rantzer, "Robust team decision theory," *Automatic Control, IEEE Transactions on*, vol. 57, no. 3, pp. 794–798, 2012.
- [21] T. Başar and P. Bernhard, *H-infinity optimal control and related minimax design problems: a dynamic game approach*. Springer Science & Business Media, 1990.
- [22] M. Colombino, T. Summers, and R. Smith, "Quadratic two-team games," in *IEEE Conference on Decision and Control, Osaka, Japan*, pp. 3784–3789, 2015.
- [23] B. Hassibi, A. H. Sayed, and T. Kailath, "Indefinite-quadratic estimation and control," *Studies in Applied and Numerical Mathematics*, 1999.
- [24] T. Basar, "Decentralized multicriteria optimization of linear stochastic systems," *Automatic Control, IEEE Transactions on*, vol. 23, no. 2, pp. 233–243, 1978.
- [25] J. Swigart and S. Lall, "An explicit state-space solution for a decentralized two-player optimal linear-quadratic regulator," in *Proceedings of the 2010 American Control Conference*, pp. 6385–6390, IEEE, 2010.