Computer-aided whole-cell design: taking a holistic approach by integrating synthetic with systems biology

Lucia Marucci^{1,2,3,+}, Matteo Barberis^{4,5,6,*,+}, Jonathan Karr^{7,*,+}, Oliver Ray^{8,*,+}, Paul R. Race^{9,3,*,+}, Miguel de Souza Andrade^{10,11}, Claire Grierson^{12,3,+}, Stefan Andreas Hoffmann¹³, Sophie Landon^{1,3}, Elibio Rech^{10,+}, Joshua Rees-Garbutt^{12,3}, Richard Seabrook^{14,+}, William Shaw¹⁵, Christopher Woods^{16,3,+}.

¹Department of Engineering Mathematics, University of Bristol, BS81UB, Bristol, UK

²School of Cellular and Molecular Medicine, University of Bristol, BS81TD, Bristol, UK

³Bristol Centre for Synthetic Biology (BrisSynBio), University of Bristol, BS81TQ, Bristol, UK

⁴Systems Biology, School of Biosciences and Medicine, Faculty of Health and Medical Sciences, University of Surrey, GU2 7XH Guildford, Surrey, UK

⁵Centre for Mathematical and Computational Biology, CMCB, University of Surrey, GU2 7XH Guildford, Surrey, UK

⁶Synthetic Systems Biology and Nuclear Organization, Swammerdam Institute for Life Sciences, University of Amsterdam, 1098 XH Amsterdam, The Netherlands

⁷ Icahn Institute for Data Science and Genomic Technology, Department of Genetics and Genomic Sciences,

Icahn School of Medicine at Mount Sinai, 1255 5th Avenue, Suite C2, New York 10029, US

⁸Department of Computer Science, University of Bristol, BS81UB, Bristol, UK

⁹School of Biochemistry, University of Bristol, BS8 1TD, Bristol, UK

¹⁰Brazilian Agriculture Research Corporation/National Institute of Science and Technology - Synthetic Biology, Brasília, DF, 70770-917, Brazil

¹¹Laboratório de Biologia Teórica e Computacional (LBTC), Universidade de Brasília DF, Brazil

¹²School of Biological Sciences, University of Bristol, BS8 1TQ, Bristol, UK

¹³Manchester Institute of Biotechnology, University of Manchester, 131 Princess Street, M1 7DN, Manchester, UK

¹⁴Elizabeth Blackwell Institute for Health Research (EBI), University of Bristol, BS8 1UH, Bristol, UK

¹⁵Department of Bioengineering, Imperial College London, London SW7 2AZ, UK

¹⁶School of Chemistry, University of Bristol, BS8 1TS, Bristol, UK

* These authors contributed equally to this work

- ⁺ Correspondence should be addressed to L.M. (lucia.marucci@bristol.ac.uk), C.G.
- (claire.grierson@bristol.ac.uk), minimal genome design; M.B. (m.barberis@surrey.ac.uk),

network design and multiscale model derivation; O.R. (csxor@bristol.ac.uk), symbolic modelling; J.K.

(karr@mssm.edu), whole-cell model development; P.R.R., E.R., R.S. (Paul.Race@bristol.ac.uk,

elibio.rech@embrapa.br, richard.seabrook@bristol.ac.uk), bioindustrial applications;

C.W.(Christopher.Woods@bristol.ac.uk), software engineering.

Keywords: whole-cell models; synthetic biology; systems biology; bioengineering; biodesign.

Abstract

Computer-aided design for synthetic biology promises to accelerate the rational and robust engineering of biological systems; it requires both detailed and quantitative mathematical and experimental models of the processes to (re)design, and software and tools for genetic engineering and DNA assembly. Ultimately, the increased precision in the design phase will have a dramatic impact on the production of designer cells and organisms with bespoke functions and increased modularity. Computer-aided design strategies require quantitative representations of cells, able to capture multiscale processes and link genotypes to phenotypes. Here, we present a perspective on how whole-cell, multiscale models could transform design-build-test-learn cycles in synthetic biology. We show how these models could significantly aid in the design and learn phases while reducing experimental testing by presenting case studies spanning from genome minimization to cell-free systems, and we discuss several challenges for the realization of our vision. The possibility to describe and build *in silico* whole-cells offers an opportunity to develop increasingly automatized, precise and accessible computer-aided design tools and strategies throughout novel interdisciplinary collaborations.

1. Introduction

Whole-cell models (WCMs) are state-of-the-art Systems Biology formalisms: they aim at representing and integrating all cellular functions within a unique computational framework, ultimately enabling a holistic and quantitative understanding of cell biology (1, 2). Quantitative and high-throughput *in silico* experiments generated from WCMs promise to significantly shorten the distance between hypothesis/design formulation and testing (3).

While simplified models for specific cellular functions were first developed over 30 years ago (e.g. formalisms for gene expression regulation (4), signaling (5) and metabolic (6) pathways, cell growth (7) and the cell cycle (8-10)), the first WCM, namely the E-Cell model, was only derived in the 1990s for *Mycoplasma genitalium*, the smallest living organisms with a 580 kb genome, following publication of the organism genome sequence (11). The so-called virtual self-surviving cell (SSC) partially stochastic model includes only a subset of protein-coding genes (105) and enables dynamic simulations which encompass various subcellular processes, including enzymatic reactions, complex formation and substance translocation. In parallel, the first genome-scale metabolic models (GSMMs) were developed by Palsson's group (12) using flux balance analysis (FBA) in the 1990s.

More recently, hundreds of GSMMs have been reconstructed for different organisms, with an increasing number of represented genes (13-15). GSMMs have been complemented with a mathematical description of other processes, such as transcription, translation and signaling (16, 17). Less than a decade ago the first hybrid dynamic model, representing all genes and molecular functions known for an organism, was proposed by Covert's group (18). In this pioneering work, Karr and colleagues integrated 28 sub-models within a unique MATLAB WCM to represent one cell cycle of *M. genitalium*; each sub-model is represented with a distinct formalism, including ordinary differential equations (ODEs), flux balance analysis (FBA), stochastic simulations and Boolean rules.

Substantial research and effort are still needed to improve WCMs' descriptive power and to increase the complexity of organisms they can represent. A complete WCM should ideally integrate multiscale interactions at the cellular level (18, 19) while accounting for the overall cellular structure (20), the dynamic structure of molecular interactions (21-23) and the spatial compartment of the subcellular components (24-26). Ensuring an accurate representation of all the mentioned processes is highly challenging, especially when aiming to model increasingly complex organisms (27-29). We refer the reader to recent efforts which provide an overview of the state-of-the-art in the development of WCMs (30, 31).

Here, we focus on the enormous potential we believe WCMs have for design-build-test cycles integrating synthetic with systems biology (Figure 1). While the discussed applications are diverse, they share a high degree of complexity which would imply extensive trial and error experimental cycles in the absence of robust computational design algorithms based on predictive models. We conclude by considering relevant challenges that interdisciplinary communities should address to fully realize our vision, discussing future directions for the integration of WCM development through synthetic and systems biology applications.

2. Whole-cell design strategies in synthetic biology

2.1 Model granularity of gene network (re)design

Mathematical models can be instrumental to (re)design network circuits that recapitulate definite biological functions. Knowledge of regulatory mechanisms in biological pathways has been gained by considering living systems as a composition of functional modules, which are investigated through minimal computer models. Examples include controllable oscillators (32-35), circadian clocks (36, 37), signaling networks (38), the metabolism (39, 40) and transcriptional regulation (41). The granularity of biochemical details uncovered about a definite regulatory network may differ between minimal and detailed computer models.

However, one may expect that if the core design of a minimal and a detailed model is similar, their general properties will match.

Understanding of a living organism at a system's level may be reached through decomposing a system into functional modules, or modular circuits that perform a definite task (42-44). The capability to sustain viability through autonomously generated offspring is essential for organismal functions and, therefore, is a feature WCMs shall account through modeling cell division, which is intimately integrated with various layers of cellular regulation (metabolism, signaling, gene regulation, transcription, etc.). A number of minimal models have been developed for the eukaryotic cell cycle by Barberis's, Tyson's and Novák's groups (45-50).

Currently, the majority of multiscale models (not WCMs) lack components able to bridge cellular networks/function (cell cycle, metabolism, signaling, gene regulation, etc.). Identification of hubs, i.e. elements with high connectivity in the cellular environment that integrate cellular networks, is a critical feature of WCMs. Transcription factors have recently been identified as hubs that integrate multiscale networks, potentially connecting the cell cycle to metabolism (51), and can be among the parts of a system that influence its state as a whole. Multiscale frameworks coupling a variable granularity of these networks are being developed, by identifying the relevant regulations occurring among common network nodes and through the use of different mathematical formalisms (52). These and other strategies are developed to progressively integrate networks of other cellular functional modules (53). Together with the identification of network designs underlying cell's autonomous oscillations, these strategies can rationalize the proper timing of offspring generation accounted by WCMs.

Designing synthetic gene networks by integrating them within WCM formalisms (as in (33)) could be critical to investigate how gene expression correlates with codon usage, explore possible cell burden effects (54), and predict modularity of synthetic gene networks and tools to modulate gene expression across different chassis (55-57).

2.2 Design and engineering of reduced genomes

Minimal genomes can be defined as reduced genomes containing only the genetic material which is essential for a cell to reproduce (58). Studying and engineering minimal genomes can be instrumental both to understand the most essential tasks a cell must perform to sustain life, and to obtain optimal chassis for synthetic biology applications, with reduced cell burden and superior robustness (59-63). Genome-scale computational models of cells can be instrumental to fully understand the dynamic and context-dependent nature of gene essentiality (64), and to rationally design minimized genomes *in silico*. Computer-aided minimal genome engineering could significantly reduce the time and cost of approaches used so far to produce reduced genomes, which can otherwise be generated by extensive experimental iterations (60, 65-70).

To the best of our knowledge, two top-down genome reduction approaches have been proposed so far based on computational genome-scale models. The MinGenome algorithm applies a mixed-integer linear programming (MILP) algorithm to a GSMM of *E. coli*, using information pertaining to essential genes and synthetic lethal pairs within the optimization (71). In contrast, Minesweeper and GAMA are top-down genome minimization algorithms based on the *M. genitalium* WCM. They exploit a divide-and-conquer approach and a genetic algorithm, respectively, to iteratively simulate reduced genomes (72); their *in silico* predictions have not been tested in the laboratory yet.

Applying GSMM-based genome reduction algorithms such as MinGenome or analogous metaheuristic equivalents which could be easily adapted for this aim (e.g. (73-75) is, at the current stage, more broadly applicable across organisms given the large availability of these formalisms. Still, as more WCMs become available, we expect WCM-based genome reduction algorithms to provide superior predictions of cellular processes and genetic interactions, thanks to their richness of multiscale cellular process representation.

2.3 Design and prototyping of cell-free systems

Cell-free transcription/translation systems, based on crude cellular extracts, are a valuable platform to address fundamental biological questions in a controllable and reproducible way. In recent years, the decrease of costs associated with this technology and significant improvements in synthesis yield capabilities (76) have made cell-free systems increasingly popular in synthetic biology for the prototyping and testing of engineered biological parts (13-15, 70) and networks (77-79). As the possible applications of cell-free systems grow (see (80) for a recent review where a spectrum of cell-free probing of cellular functions and of applications are discussed in detail), mathematical models are developed to quantitatively formalize how biological processes perform within cell-free platforms (81).

So far, deterministic models (ordinary differential equations-, or constraint-based) have been proposed to describe specific processes within cell-free platforms such as transcription and translation (78, 82, 83), resource competition (84-87) and metabolism (88). The integration of mathematical formalisms across scales for cell-free platforms, building towards WCMs, could be highly beneficial to both facilitate *de novo* design of circuits, and to quantitatively compare *in vitro* cell-free products with their *in vivo* counterparts.

2.4 Whole-cell biosensor design and testing

Biosensors are analytical devices which can convert a biochemical reaction into a measurable signal. The recognition unit in a biosensor can be composed of whole cells, nucleic acids, enzymes, proteins, antibodies or combinations thereof. Synthetic biology has significantly accelerated biosensor development; new generation whole-cell biosensors (i.e. sensors implemented throughout living cells) have been engineered, for example, to detect arsenic (89) and other analytes (including pollutants and antibiotics (90)), for microbial detection in industrial settings (91) and for *in vivo* diagnostic applications (e.g. detection of environmental signals in the gut (92) and diagnosis of liver metastases (93); see (94) for an overview of translation of promising technologies into diagnostic tools).

The application of WCMs to the design, prototyping and testing of whole-cell biosensors could suggest rational approaches to tune their sensitivity, stability and dynamic range while facilitating the choice of the ideal chassis and, if needed, guide its re-engineering to optimize biosensor performance (95). If WCMs become available for different chassis and entire organisms, they could also support the design of optimized targeted delivery of genetically encoded biosensors.

2.5 Industrial implications of whole-cell models

Although the intellectual merit of pursuing a computer-aided whole-cell design approach is unquestioned, it is clear that the success of this endeavor will ultimately be judged by its impact on industry. The increasing drive towards 'green' chemistry approaches, allied to increases in gene synthesis speed and capability and associated cost reductions, are making biosynthesis an increasingly appealing route for the manufacture of high-value chemicals (96). This includes a plethora of opportunities across the pharmaceutical, agrochemical, commodity chemical, and materials sectors, amongst others.

A major challenge, however, remains the development of robust, scalable microbial chassis, whose metabolic processes can be predictably tuned for a desired outcome (97). Currently, chassis choice is largely restricted to a subset of genetically tractable microorganisms, whose physiology and performance during fermentation are well understood, and for whom effective molecular genetic tools required for their manipulation exist. Chassis

optimization to date has relied exclusively on incremental, stepwise improvements in desired host strain characteristics, including growth rate, feedstock utilization and product yield (98). For these reasons, the process of chassis optimization remains prohibitively slow and expensive, accounting in part for the paucity of high-value small molecules that are currently manufactured using synthetic biology processes. Targeted manipulations often lead to unanticipated off-target effects, linked to the co-dependency of metabolic processes, which generally function in concert within interdependent cellular networks (99): perturbations may compromise rather than enhance desirable characteristics, leading to undesired outcomes. Clearly, robust, predictable WCMs represent an attractive solution to the problem of chassis optimization, affording a catch-all tool that can be used to unpick dependencies and ensure that performance criteria can be met. Besides, the complexities associated with population heterogeneity during chassis fermentation must be resolved (100). For fermentation-based industrial processes to be tractable, product yields must be sufficiently high to make biosynthesis financially viable. The emergence of 'cheaters' or slow-growers within microbial populations must be addressed. This will undoubtably be best achieved through tunable regulatory processes that operate throughout populations. The introduction of such characteristics is a major challenge to conventional chassis design approaches. However, this could be much more easily implemented and tested by employing WCM driven approaches.

Critical to the success of a computer-aided whole-cell design approach is the quality of the employed model (101). Microbial systems with small genomes represent a compelling entry point for study, with model development possibly being facilitated by ongoing studies focused on establishing the core constituents of a functional genome. These studies are being predominantly driven by genome minimization experiments, which in turn can be used to further refine model performance. Importantly, fundamental gaps remain in our understanding of microbial metabolic processes, and this will unquestionably hinder progress (102). However, the capacity of WCMs to predict previously unidentified metabolic dependencies should be viewed as an acid test of model validity. Indeed, GSMMs often fail due to their inability to account for interdependencies, a feature which has led to skepticism within industrial circles, questioning the value of such models. Whole-cell approaches offer a mechanism to circumvent this issue and could increase reliability and promote confidence in the adoption of WCMs. This is of particular significance when developing chassis for 'non-natural' products whose chemistries sit outside those of metabolites found in nature (98). Expanding the metabolic capacity of chassis organisms to deliver such novel products risks introducing additional complexities, including excessive depletion of core metabolite pools or the generation of toxic products or intermediates. Design approaches driven by WCMs are uniquely placed to identify such issues and provide a route to their circumvention.

The capacity to design-in explicit control over cellular behavior is also critical for industrial adoption of model-derived chassis. It can be argued that the ability to regulate cellular processes is as important as defining the processes themselves. Tunable regulatory systems must afford a degree of both intrinsic and extrinsic control. Synthetic biology-based approaches for the construction of genetic circuitry are now placing us on a path to broad-reaching cellular regulation, though issues still exist. These systems are often insufficiently orthogonal, with bespoke designs required for different chassis due to variations in core metabolic process (39). Again, whole-cell design approaches offer a solution to this issue, as such systems can be predefined and tested for functionality *in silico* prior to undertaking costly lab experimentation.

3. What's next? Going beyond the prototype

In recent years, advances in genomic measurement technologies for data generation, the establishment of data repositories, and the development of WCM simulation platforms have significantly facilitated the derivation of WCMs (see (30) for a review). Nevertheless, the

implementation of WCM-based design-build-test cycles for genome-scale engineering requires further challenges to be addressed (103).

If a model has to be used for the design and prototyping of an engineered living system, the model needs to be reliable. Even for a simple organism, the number of kinetic parameters surges as the complexity and the level of detail of a mathematical model increase; constraining parameters thus becomes harder and requires extensive experimental data. To set the 1,462 quantitative parameters of the *M. genitalium* WCM, values from other species were incorporated due to a lack of organism-specific data (104); a combination of parameter values reported from previous experiments and numerical optimization on a reduced model was performed. While, ideally, we would like to measure all kinetic parameters directly from experiments, we still lack techniques to measure each state in individual cells over time, and across all possible environmental conditions. Mathematical models can be used to produce predictions of missing data, however they often abstract physical processes using simplifying assumptions which might hold in specific conditions only (105). A combination of direct experimental estimation and parameter inference might be needed for genome-scale formalisms and WCMs, for example using scalable inference techniques to parameterize submodels.

Sensitivity analysis, usually performed by perturbing parameters to understand how uncertainties affect the model outputs (106), can become extremely computationally expensive when applied to genome-scale models. Alternatively, statistical approaches such as those based on Bayesian methods (107) or on the Fisher information matrix (108) could be carefully carried out at least at the sub-model level, and possibly scaled up to WCMs. The Reverse Engineering Assessments and Methods (DREAM8) parameter estimation challenge (109) was organized to develop new parameter estimation techniques specific for WCMs. It suggested possible interesting avenues for WCM parameterization (i.e. model reduction and a combination of differential evolution and random forests), and highlighted that the availability of comprehensive data-set is critical to ensure the model is practically identifiable (110).

It is also important to consider the structural uncertainties in the model, which depend on model assumptions. While, for certain sets of models (e.g. small ordinary differential equation systems for signaling pathways), likelihood- and Bayesian-based approaches have been proposed for model selection (111) (112) and Semidefinite Programming for model invalidation (113), no suitable techniques for WCM selection have been proposed to date.

We foresee that automation will play a fundamental role in the derivation of WCMs for eukaryotic organisms and in their application to design complex processes. Ideally, we would like to introduce automation at different stages, such as data extraction from the literature, model derivation, and model/data integration both within the model fitting and validation steps, and when comparing *in vitro* design prediction with *in vivo* tests (103). This, in turn, will require the adoption of standards for both data and model repositories.

Researchers have started to collect data needed for WCM development into public repositories (e.g. (114-118)); still, data needed to derive and fit WCMs are dispersed across many repositories and publications and often not annotated or normalized, ultimately requiring a massive manual effort. Federated archives of repositories, such as the PDB-Dev system to deposit Integrative/Hybrid models and corresponding data (119), might be well placed to archive and disseminate both data and models, while enabling different researchers to attempt alternative modelling/parameterization approaches. Covert's group developed the WholeCellKB database (120) to organize the quantitative measurements (over 1,400) from which the *M. genitalium* WCM was derived; it would be ideal to enable automatic access and querying in such databases.

To enhance WCM reproducibility, new standards and simulations software are also needed (121). Researchers should invest efforts to use standard formats such as the Systems

Biology Markup Language (SBML) (122) and the Systems Biology Graphical Notation (SBGN) (123). At present, various aspects of the *M. genitalium* WCM cannot be represented by SBML, mainly due to the multi-algorithmic nature of the model itself (124). Further development of standard modelling formats is needed to enable reproducible WCM simulations, e.g. by including in the SMBL Hierarchical Model Composition package ontologies which could represent the algorithm needed for specific sub-models (125). In the context of synthetic biology applications, we believe it would be appropriate and beneficial to report and deposit data related to various iterations of WCM-generated *in silico* predictions, *in vivo* testing and possible model/design refinement; this would establish the predictive power of WCMs and illuminate steps to make design-build-test-learn cycles more effective.

To facilitate the adoption of WCMs for synthetic biology applications, highperformance parallelized computer clusters are required to coordinate and run the models with lengthy runtimes and large corresponding databases across computer clusters, to parameterize and validate the models, and then to integrate them in design cycles in combination with optimization algorithms (104) (126).

The implementation of standardized tools to share data and simulate WCMs would, in turn, facilitate model validation. This should involve the definition of proper metrics and formal model verification techniques such as those developed for SBML-encoded models (127).

4. (Re)thinking system approaches: a collaborative effort

In addressing the aforementioned challenges, we believe there is a tremendous opportunity to rethink approaches used so far to generate genome-scale models, including WCMs, and to integrate with broader communities including software engineers, computer scientists, structural biologists, bioinformaticians, and systems and synthetic biologists.

We do anticipate that, as diverse communities synergize on WCM-related research, different kinds of formalisms might be integrated within genome-scale models. Symbolic reasoning (e.g. satisfiability solving, model checking, theorem proving, formal methods, logic programming and Boolean networks) provide a range of expressive and intuitive logical frameworks that could potentially complement and help glue together sub-models at different scales. Such methods are routinely applied to complex systems in the electronics and software industries, and have been tentatively applied to biological systems for nearly a decade (128). Recent work showed the feasibility of applying logic programming methods to signaling pathways (129), metabolic networks (130) and automating a mechanistic philosophy of scientific discovery in simulated organisms (131); it should be feasible to integrate such sub-models within a WCM framework.

We believe there is scope to further increase the descriptive and predictive ability of WCMs across spatial and temporal scales by integrating the structural biology and the molecular modelling communities to carefully consider not only the biochemical, but also the physical, molecular and structural components of cells. The development of the so-called "physical" WCMs (see (31) and (132) for comprehensive reviews) is an emerging field, with the first models describing minimal cellular environments in full atomistic detail (21, 133). With the final aim to integrate cellular and physical WCMs within a multiscale framework (134), we need approaches which can cope with the limitations of atomistic models of biomolecules (mainly in terms of computational resources), possibly exploiting coarse-grained (135, 136) or continuum (137) approaches.

By collaborating with software engineers, we need to develop tools which can enable, and possibly automate, the integration of different data types across scales, model derivation, fitting and validation, and visualization and interpretation of results (27). Rule-based models might become the new standard to represent each molecular species with the required level of granularity and multi-algorithmic sub-models (e.g. flux balance analysis and stochastic

dynamical models). Frameworks where intuitive logic is coupled to rule-based models have started to be developed recently (52).

As we produce ever-increasing amounts of experimental data and increasingly sophisticated computational tools to realize detailed and complex representations of actual cells, approaches instead focusing on deliberately abstract and parsimonious simulations of artificial cellular systems provide a valuable change of perspective. Such "toy models" might be a valuable tool to test different algorithms for model derivation and fitting, while offering an opportunity to engage with broader research communities and with the public (138).

Finally, we believe there is tremendous potential for applying machine learning techniques to both WCM derivation and their applications in synthetic biology. Two recent works (139, 140) showed that deep neural networks are well placed to reconstruct the architecture of living systems (namely, the hierarchical organization of nuclear transcriptional factors in the nucleus (139) and of a basic eukaryotic cell (140)) and predict cell states and phenotypes. In both cases, the configuration of network layers and thus the biological structure were formulated using extensive prior knowledge, ultimately enabling fully "visible" systems, where all the internal biological states can be interrogated mechanistically (141). Machine learning could be beneficial to systematically process large *in vivo* and *in silico* whole-cell data-sets, for example by applying Bayesian inference, to integrate data from diverse sources and supplement sparse data (142), and to help to automatically classify WCM simulations and link phenotypes to genotypes (143). Optimal experimental design techniques might also offer a valuable methodology to select the best experimental datasets for both model identification and validation (144).

5. Discussion

We have shown that WCMs are likely to be instrumental to inform design-build-test cycles across synthetic biology applications. WCMs can accelerate the realization of "designer" cells and organisms tailored to specific functions, reducing experimental iterations and increasing the predictive power of computational formalisms used so far.

In the (re)design of cellular network functionalities, it is therefore important to quantitatively analyze and predict, through dedicated modelling strategies, the dynamics of interactions between various layers of cellular regulation. Thus, WCMs shall take into account how different cellular layers are integrated, and how regulatory feedback among these layers occurs in time. These challenges are tackled through integrative computational and experimental collaborative efforts aimed, respectively, towards: (i) engineering *in vivo* network designs which, through predictive systems biology, may be able to autonomously oscillate, sustaining generation of offspring, and (ii) extraction, visualization and functional exploration of regulatory interactions among cellular layers through novel multiscale modeling frameworks.

As synthetic biology moves toward the (re)engineering of entire genomes and multicellular systems, interdisciplinary communities need to collaborate for the development of tools that are required to improve the predictive power of WCMs. Although challenges remain, it is clear that the adoption of model-based methods has the potential to transform both basic research and the current bioproduction development process, leading to marked improvements in host performance and product yield on an industrial scale.

Ultimately, as the development of human genome-scale kinetic models becomes more feasible (27, 145), whole-cell formalisms might become an indispensable tool to study human variation, and design bespoke treatments and synthetic cellular screening systems.




Figure 1 Integrated design-build-test-learn cycles in synthetic biology encompassing whole-cell model-guided approaches, and relative applications.

Author contributions

L.M., M.B., J.K., O.R. and P.R.R. wrote the manuscript; M.d.S.A. prepared the figure; all other authors participated to discussion within the workshop, helped with editing and/or provided feedback.

Acknowledgements

This work captures discussions between participants at the "Computer-Aided Whole-Cell Design and Engineering" Workshop held on the 02-03 July 2019 at the University of Bristol, UK, and funded by the Engineering and Physical Sciences Research Council (EPSRC) within the remits of the Big Ideas initiative. We sincerely thank Dr Kathleen Sedgley for her support with the workshop organization, and Dr Thomas Gorochowski for participating in discussions.

Funding

L.M. was funded by the Engineering and Physical Sciences Research Council (EPSRC, grants EP/R041695/1 and EP/S01876X/1) and Horizon 2020 (CosyBio, grant agreement 766840); M.B. was funded by the Systems Biology Grant of the University of Surrey; J.K. was funded by the National Institutes of Health (award R35GM119771); P.R.R. was funded by EPSRC (EP/R020957/1) and the Biotechnology and Biological Sciences Research Council (BBSRC, BB/T001968/1); C.G., L.M. and P.R.R. were funded by BrisSynBio, a BBSRC/EPSRC Synthetic Biology Research Centre (BB/L01386X/); S.L. and J.R. were funded by EPSRC Future Opportunity PhD scholarships; E.R. was funded by INCT BioSyn (National Institute of Science and Technology in Synthetic Biology), CNPq (National Council for Scientific and Technological Development), CAPES (Coordination for the Improvement of Higher Education Personnel), Brazilian Ministry of Health, and FAPDF (Research Support Foundation of the Federal District), Brazil.

Conflicts of interest

The authors declare no competing financial interests.

References

1. Karr JR, Takahashi K, Funahashi A. The principles of whole-cell modeling. Curr Opin Microbiol. 2015;27:18-24.

2. Tomita M. Whole-cell simulation: a grand challenge of the 21st century. Trends Biotechnol. 2001;19(6):205-10.

3. Carrera J, Covert MW. Why Build Whole-Cell Models? Trends Cell Biol. 2015;25(12):719-22.

4. McAdams HH, Arkin A. Stochastic mechanisms in gene expression. Proc Natl Acad Sci U S A. 1997;94(3):814-9.

5. Morton-Firth CJ, Bray D. Predicting temporal fluctuations in an intracellular signalling pathway. J Theor Biol. 1998;192(1):117-28.

6. Cornish-Bowden A, Hofmeyr JH. MetaModel: a program for modelling and control analysis of metabolic pathways on the IBM PC and compatibles. Comput Appl Biosci. 1991;7(1):89-93.

7. Shu J, Shuler ML. A mathematical model for the growth of a single cell of E. coli on a glucose/glutamine/ammonium medium. Biotechnol Bioeng. 1989;33(9):1117-26.

8. Novak B, Tyson JJ. Numerical analysis of a comprehensive model of M-phase control in Xenopus oocyte extracts and intact embryos. J Cell Sci. 1993;106 (Pt 4):1153-68.

9. Goldbeter A. A minimal cascade model for the mitotic oscillator involving cyclin and cdc2 kinase. Proc Natl Acad Sci U S A. 1991;88(20):9107-11.

10. Tyson JJ. Modeling the cell division cycle: cdc2 and cyclin interactions. Proc Natl Acad Sci U S A. 1991;88(16):7328-32.

11. Tomita M, Hashimoto K, Takahashi K, Shimizu TS, Matsuzaki Y, Miyoshi F, et al. E-CELL: software environment for whole-cell simulation. Bioinformatics. 1999;15(1):72-84.

12. Varma A, Palsson BO. Stoichiometric flux balance models quantitatively predict growth and metabolic by-product secretion in wild-type Escherichia coli W3110. Appl Environ Microbiol. 1994;60(10):3724-31.

13. McCloskey D, Palsson BO, Feist AM. Basic and applied uses of genome-scale metabolic network reconstructions of Escherichia coli. Mol Syst Biol. 2013;9:661.

14. Mendoza SN, Olivier BG, Molenaar D, Teusink B. A systematic assessment of current genome-scale metabolic reconstruction tools. Genome Biol. 2019;20(1):158.

15. Yilmaz LS, Walhout AJ. Metabolic network modeling with model organisms. Curr Opin Chem Biol. 2017;36:32-9.

16. Lee JM, Gianchandani EP, Eddy JA, Papin JA. Dynamic analysis of integrated signaling, metabolic, and regulatory networks. PLoS Comput Biol. 2008;4(5):e1000086.

17. Thiele I, Jamshidi N, Fleming RM, Palsson BO. Genome-scale reconstruction of Escherichia coli's transcriptional and translational machinery: a knowledge base, its mathematical formulation, and its functional characterization. PLoS Comput Biol. 2009;5(3):e1000312.

18. Karr JR, Sanghvi JC, Macklin DN, Gutschow MV, Jacobs JM, Bolival B, Jr., et al. A whole-cell computational model predicts phenotype from genotype. Cell. 2012;150(2):389-401.

19. King ZA, Lu J, Drager A, Miller P, Federowicz S, Lerman JA, et al. BiGG Models: A platform for integrating, standardizing and sharing genome-scale models. Nucleic Acids Res. 2016;44(D1):D515-22.

20. Betts MJ, Russell RB. The hard cell: from proteomics to a whole cell model. FEBS Lett. 2007;581(15):2870-6.

21. Yu I, Mori T, Ando T, Harada R, Jung J, Sugita Y, et al. Biomolecular interactions modulate macromolecular structure and dynamics in atomistic model of a bacterial cytoplasm. Elife. 2016;5.

22. McGuffee SR, Elcock AH. Diffusion, crowding & protein stability in a dynamic molecular model of the bacterial cytoplasm. PLoS Comput Biol. 2010;6(3):e1000694.

23. Noske AB, Costin AJ, Morgan GP, Marsh BJ. Expedited approaches to whole cell electron tomography and organelle mark-up in situ in high-pressure frozen pancreatic islets. J Struct Biol. 2008;161(3):298-313.

24. Ander M, Beltrao P, Di Ventura B, Ferkinghoff-Borg J, Foglierini M, Kaplan A, et al. SmartCell, a framework to simulate cellular processes that combines stochastic approximation with diffusion and localisation: analysis of simple networks. Syst Biol (Stevenage). 2004;1(1):129-38.

25. Takahashi K, Arjunan SN, Tomita M. Space in systems biology of signaling pathways-towards intracellular molecular crowding in silico. FEBS Lett. 2005;579(8):1783-8.

26. Thul PJ, Akesson L, Wiking M, Mahdessian D, Geladaki A, Ait Blal H, et al. A subcellular map of the human proteome. Science. 2017;356(6340).

27. Szigeti B, Roth YD, Sekar JAP, Goldberg AP, Pochiraju SC, Karr JR. A blueprint for human whole-cell modeling. Curr Opin Syst Biol. 2018;7:8-15.

28. Bouhaddou M, Barrette AM, Stern AD, Koch RJ, DiStefano MS, Riesel EA, et al. A mechanistic pan-cancer pathway model informed by multi-omics data interprets stochastic cell fate responses to drugs and mitogens. PLoS Comput Biol. 2018;14(3):e1005985.

29. Singla J, McClary KM, White KL, Alber F, Sali A, Stevens RC. Opportunities and Challenges in Building a Spatiotemporal Multi-scale Model of the Human Pancreatic beta Cell. Cell. 2018;173(1):11-9.

30. Goldberg AP, Szigeti B, Chew YH, Sekar JA, Roth YD, Karr JR. Emerging whole-cell modeling principles and methods. Curr Opin Biotechnol. 2018;51:97-102.

31. Feig M, Sugita Y. Whole-Cell Models and Simulations in Molecular Detail. Annu Rev Cell Dev Biol. 2019;35:191-211.

32. Tomazou M, Barahona M, Polizzi KM, Stan GB. Computational Re-design of Synthetic Genetic Oscillators for Independent Amplitude and Frequency Modulation. Cell Syst. 2018;6(4):508-20 e5.

33. Purcell O, Jain B, Karr JR, Covert MW, Lu TK. Towards a whole-cell modeling approach for synthetic biology. Chaos. 2013;23(2):025112.

34. Purcell O, Savery NJ, Grierson CS, di Bernardo M. A comparative analysis of synthetic genetic oscillators. J R Soc Interface. 2010;7(52):1503-24.

35. Marucci L, Barton DA, Cantone I, Ricci MA, Cosma MP, Santini S, et al. How to turn a genetic circuit into a synthetic tunable oscillator, or a bistable switch. PLoS One. 2009;4(12):e8083.

36. Ananthasubramaniam B, Schmal C, Herzel H. Amplitude Effects Allow Short Jet Lags and Large Seasonal Phase Shifts in Minimal Clock Models. J Mol Biol. 2020.

37. Gerard C, Gonze D, Goldbeter A. Dependence of the period on the rate of protein degradation in minimal models for circadian oscillations. Philos Trans A Math Phys Eng Sci. 2009;367(1908):4665-83.

38. Prescott AM, Abel SM. Combining in silico evolution and nonlinear dimensionality reduction to redesign responses of signaling networks. Phys Biol. 2017;13(6):066015.

39. Pandit AV, Srinivasan S, Mahadevan R. Redesigning metabolism based on orthogonality principles. Nat Commun. 2017;8:15188.

40. Castellanos M, Wilson DB, Shuler ML. A modular minimal cell model: purine and pyrimidine transport and metabolism. Proc Natl Acad Sci U S A. 2004;101(17):6681-6.

41. Carrera J, Rodrigo G, Jaramillo A. Model-based redesign of global transcription regulation. Nucleic Acids Res. 2009;37(5):e38.

42. Hartwell LH, Hopfield JJ, Leibler S, Murray AW. From molecular to modular cell biology. Nature. 1999;402(6761 Suppl):C47-52.

43. Ravasz E, Somera AL, Mongru DA, Oltvai ZN, Barabasi AL. Hierarchical organization of modularity in metabolic networks. Science. 2002;297(5586):1551-5.

44. Kitano H. Computational systems biology. Nature. 2002;420(6912):206-10.

45. Mondeel T, Ivanov O, Westerhoff HV, Liebermeister W, Barberis M. Clb3-centered regulations are recurrent across distinct parameter regions in minimal autonomous cell cycle oscillator designs. NPJ Syst Biol Appl. 2020;6(1):8.

46. Linke C, Chasapi A, Gonzalez-Novo A, Al Sawad I, Tognetti S, Klipp E, et al. A Clb/Cdk1-mediated regulation of Fkh2 synchronizes CLB expression in the budding yeast cell cycle. NPJ Syst Biol Appl. 2017;3:7.

47. Barberis M, Linke C, Adrover MA, Gonzalez-Novo A, Lehrach H, Krobitsch S, et al. Sic1 plays a role in timing and oscillatory behaviour of B-type cyclins. Biotechnol Adv. 2012;30(1):108-30.

48. Battogtokh D, Tyson JJ. Bifurcation analysis of a model of the budding yeast cell cycle. Chaos. 2004;14(3):653-61.

49. Gerard C, Tyson JJ, Novak B. Minimal models for cell-cycle control based on competitive inhibition and multisite phosphorylations of Cdk substrates. Biophys J. 2013;104(6):1367-79.

50. Gerard C, Tyson JJ, Coudreuse D, Novak B. Cell cycle control by a minimal Cdk network. PLoS Comput Biol. 2015;11(2):e1004056.

51. Mondeel T, Holland P, Nielsen J, Barberis M. ChIP-exo analysis highlights Fkh1 and Fkh2 transcription factors as hubs that integrate multi-scale networks in budding yeast. Nucleic Acids Res. 2019;47(15):7825-41.

52. van der Zee L, Barberis M. Advanced Modeling of Cellular Proliferation: Toward a Multi-scale Framework Coupling Cell Cycle to Metabolism by Integrating Logical and Constraint-Based Models. Methods Mol Biol. 2019;2049:365-85.

53. Prescott TP, Lang M, Papachristodoulou A. Quantification of Interactions between Dynamic Cellular Network Functionalities by Cascaded Layering. PLoS Comput Biol. 2015;11(5):e1004235.

54. Borkowski O, Ceroni F, Stan GB, Ellis T. Overloaded and stressed: whole-cell considerations for bacterial synthetic biology. Curr Opin Microbiol. 2016;33:123-30.

55. Pedone E, Postiglione L, Aulicino F, Rocca DL, Montes-Olivas S, Khazim M, et al. A tunable dual-input system for on-demand dynamic gene expression regulation. Nat Commun. 2019;10(1):4481.

56. Gomide MS, Sales TT, Barros LRC, Limia CG, de Oliveira MA, Florentino LH, et al. Genetic switches designed for eukaryotic cells and controlled by serine integrases. Commun Biol. 2020;3(1):255.

57. Way JC, Collins JJ, Keasling JD, Silver PA. Integrating biological redesign: where synthetic biology came from and where it needs to go. Cell. 2014;157(1):151-61.

58. Glass JI, Merryman C, Wise KS, Hutchison CA, 3rd, Smith HO. Minimal Cells-Real and Imagined. Cold Spring Harb Perspect Biol. 2017;9(12).

59. Ceroni F, Ellis T. The challenges facing synthetic biology in eukaryotes. Nat Rev Mol Cell Biol. 2018;19(8):481-2.

60. Hutchison CA, 3rd, Chuang RY, Noskov VN, Assad-Garcia N, Deerinck TJ, Ellisman MH, et al. Design and synthesis of a minimal bacterial genome. Science. 2016;351(6280):aad6253.

61. Landon S, Rees-Garbutt J, Marucci L, Grierson C. Genome-driven cell engineering review: in vivo and in silico metabolic and genome engineering. Essays Biochem. 2019;63(2):267-84.

62. Mol M, Kabra R, Singh S. Genome modularity and synthetic biology: Engineering systems. Prog Biophys Mol Biol. 2018;132:43-51.

63. Moya A, Gil R, Latorre A, Pereto J, Pilar Garcillan-Barcia M, de la Cruz F. Toward minimal bacterial cells: evolution vs. design. FEMS Microbiol Rev. 2009;33(1):225-35.

64. Rancati G, Moffat J, Typas A, Pavelka N. Emerging and evolving concepts in gene essentiality. Nat Rev Genet. 2018;19(1):34-49.

65. Breuer M, Earnest TM, Merryman C, Wise KS, Sun L, Lynott MR, et al. Essential metabolism for a minimal cell. Elife. 2019;8.

66. Hirokawa Y, Kawano H, Tanaka-Masuda K, Nakamura N, Nakagawa A, Ito M, et al. Genetic manipulations restored the growth fitness of reduced-genome Escherichia coli. J Biosci Bioeng. 2013;116(1):52-8.

67. Iwadate Y, Honda H, Sato H, Hashimoto M, Kato J. Oxidative stress sensitivity of engineered Escherichia coli cells with a reduced genome. FEMS Microbiol Lett. 2011;322(1):25-33.

68. Posfai G, Plunkett G, 3rd, Feher T, Frisch D, Keil GM, Umenhoffer K, et al. Emergent properties of reduced-genome Escherichia coli. Science. 2006;312(5776):1044-6.

69. Zhou J, Wu R, Xue X, Qin Z. CasHRA (Cas9-facilitated Homologous Recombination Assembly) method of constructing megabase-sized DNA. Nucleic Acids Res. 2016;44(14):e124.

70. Reuss DR, Altenbuchner J, Mader U, Rath H, Ischebeck T, Sappa PK, et al. Large-scale reduction of the Bacillus subtilis genome: consequences for the transcriptional network, resource allocation, and metabolism. Genome Res. 2017;27(2):289-99.

71. Wang L, Maranas CD. MinGenome: An In Silico Top-Down Approach for the Synthesis of Minimized Genomes. ACS Synth Biol. 2018;7(2):462-73.

72. Rees-Garbutt J, Chalkley O, Landon S, Purcell O, Marucci L, Grierson C. Designing minimal genomes using whole-cell models. Nat Commun. 2020;11(1):836.

73. Mutturi S. FOCuS: a metaheuristic algorithm for computing knockouts from genome-scale models for strain optimization. Mol Biosyst. 2017;13(7):1355-63.

74. Tang PW, Chua PS, Chong SK, Mohamad MS, Choon YW, Deris S, et al. A Review of Gene Knockout Strategies for Microbial Cells. Recent Pat Biotechnol. 2015;9(3):176-97.

75. Burgard AP, Pharkya P, Maranas CD. Optknock: a bilevel programming framework for identifying gene knockout strategies for microbial strain optimization. Biotechnol Bioeng. 2003;84(6):647-57.

76. Calhoun KA, Swartz JR. Energizing cell-free protein synthesis with glucose metabolism. Biotechnol Bioeng. 2005;90(5):606-13.

77. Noireaux V, Bar-Ziv R, Libchaber A. Principles of cell-free genetic circuit assembly. Proc Natl Acad Sci U S A. 2003;100(22):12672-7.

78. Siegal-Gaskins D, Tuza ZA, Kim J, Noireaux V, Murray RM. Gene circuit performance characterization and resource usage in a cell-free "breadboard". ACS Synth Biol. 2014;3(6):416-25.

79. Takahashi MK, Chappell J, Hayes CA, Sun ZZ, Kim J, Singhal V, et al. Rapidly characterizing the fast dynamics of RNA genetic circuitry with cell-free transcription-translation (TX-TL) systems. ACS Synth Biol. 2015;4(5):503-15.

80. Silverman AD, Karim AS, Jewett MC. Cell-free gene expression: an expanded repertoire of applications. Nat Rev Genet. 2020;21(3):151-70.

81. Koch M, Faulon J-L, Borkowski O. Models for Cell-Free Synthetic Biology: Make Prototyping Easier, Better, and Faster. Frontiers in Bioengineering and Biotechnology. 2018;6(182).

82. Karzbrun E, Shin J, Bar-Ziv RH, Noireaux V. Coarse-grained dynamics of protein synthesis in a cell-free system. Phys Rev Lett. 2011;106(4):048104.

83. Stogbauer T, Windhager L, Zimmer R, Radler JO. Experiment and mathematical modeling of gene expression dynamics in a cell-free system. Integr Biol (Camb). 2012;4(5):494-501.

84. Borkowski O, Bricio C, Murgiano M, Rothschild-Mancinelli B, Stan GB, Ellis T. Cellfree prediction of protein expression costs for growing cells. Nat Commun. 2018;9(1):1457.

85. Underwood KA, Swartz JR, Puglisi JD. Quantitative polysome analysis identifies limitations in bacterial cell-free protein synthesis. Biotechnol Bioeng. 2005;91(4):425-35.

86. Moore SJ, MacDonald JT, Wienecke S, Ishwarbhai A, Tsipa A, Aw R, et al. Rapid acquisition and model-based analysis of cell-free transcription-translation reactions from nonmodel bacteria. Proc Natl Acad Sci U S A. 2018;115(19):E4340-E9.

87. Matsuura T, Hosoda K, Shimizu Y. Robustness of a Reconstituted Escherichia coli Protein Translation System Analyzed by Computational Modeling. ACS Synth Biol. 2018;7(8):1964-72.

88. Vilkhovoy M, Horvath N, Shih CH, Wayman JA, Calhoun K, Swartz J, et al. Sequence Specific Modeling of E. coli Cell-Free Protein Synthesis. ACS Synth Biol. 2018;7(8):1844-57.

89. Diesel E, Schreiber M, van der Meer JR. Development of bacteria-based bioassays for arsenic detection in natural waters. Anal Bioanal Chem. 2009;394(3):687-93.

90. van der Meer JR, Belkin S. Where microbiology meets microengineering: design and applications of reporter bacteria. Nat Rev Microbiol. 2010;8(7):511-22.

91. Lu TK, Bowers J, Koeris MS. Advancing bacteriophage-based microbial diagnostics with synthetic biology. Trends Biotechnol. 2013;31(6):325-7.

92. Kotula JW, Kerns SJ, Shaket LA, Siraj L, Collins JJ, Way JC, et al. Programmable bacteria detect and record an environmental signal in the mammalian gut. Proc Natl Acad Sci U S A. 2014;111(13):4838-43.

93. Danino T, Prindle A, Kwong GA, Skalak M, Li H, Allen K, et al. Programmable probiotics for detection of cancer in urine. Sci Transl Med. 2015;7(289):289ra84.

94. Slomovic S, Pardee K, Collins JJ. Synthetic biology devices for in vitro and in vivo diagnostics. Proc Natl Acad Sci U S A. 2015;112(47):14429-35.

95. Hicks M, Bachmann TT, Wang B. Synthetic Biology Enables Programmable Cell-Based Biosensors. Chemphyschem. 2020;21(2):131.

96. El Karoui M, Hoyos-Flight M, Fletcher L. Future Trends in Synthetic Biology-A Report. Front Bioeng Biotechnol. 2019;7:175.

97. Xu X, Liu Y, Du G, Ledesma-Amaro R, Liu L. Microbial Chassis Development for Natural Product Biosynthesis. Trends Biotechnol. 2020.

98. Calero P, Nikel PI. Chasing bacterial chassis for metabolic engineering: a perspective review from classical to non-traditional microorganisms. Microb Biotechnol. 2019;12(1):98-124.

99. Woolston BM, Edgar S, Stephanopoulos G. Metabolic engineering: past and future. Annu Rev Chem Biomol Eng. 2013;4:259-88.

100. Danchin A. Scaling up synthetic biology: Do not forget the chassis. FEBS Lett. 2012;586(15):2129-37.

101. Fernandez-Castane A, Feher T, Carbonell P, Pauthenier C, Faulon JL. Computer-aided design for metabolic engineering. J Biotechnol. 2014;192 Pt B:302-13.

102. Price MN, Wetmore KM, Waters RJ, Callaghan M, Ray J, Liu H, et al. Mutant phenotypes for thousands of bacterial genes of unknown function. Nature. 2018;557(7706):503-9.

103. Bartley BA, Beal J, Karr JR, Strychalski EA. Organizing genome engineering for the gigabase scale. Nat Commun. 2020;11(1):689.

104. Macklin DN, Ruggero NA, Covert MW. The future of whole-cell modeling. Curr Opin Biotechnol. 2014;28:111-5.

105. Babtie AC, Stumpf MPH. How to deal with parameters for whole-cell modelling. J R Soc Interface. 2017;14(133).

106. Erguler K, Stumpf MP. Practical limits for reverse engineering of dynamical systems: a statistical analysis of sensitivity and parameter inferability in systems biology models. Mol Biosyst. 2011;7(5):1593-602.

107. Vernon I, Liu J, Goldstein M, Rowe J, Topping J, Lindsey K. Bayesian uncertainty analysis for complex systems biology models: emulation, global parameter searches and evaluation of gene functions. BMC Syst Biol. 2018;12(1):1.

108. Rand DA. Mapping global sensitivity of cellular network dynamics: sensitivity heat maps and a global summation law. J R Soc Interface. 2008;5 Suppl 1:S59-69.

109. Karr JR, Williams AH, Zucker JD, Raue A, Steiert B, Timmer J, et al. Summary of the DREAM8 Parameter Estimation Challenge: Toward Parameter Identification for Whole-Cell Models. PLoS Comput Biol. 2015;11(5):e1004096.

110. Ashyraliyev M, Fomekong-Nanfack Y, Kaandorp JA, Blom JG. Systems biology: parameter estimation for biochemical models. FEBS J. 2009;276(4):886-902.

111. Kirk P, Thorne T, Stumpf MP. Model selection in systems and synthetic biology. Curr Opin Biotechnol. 2013;24(4):767-74.

112. Wilkinson DJ. Bayesian methods in bioinformatics and computational systems biology. Brief Bioinform. 2007;8(2):109-16.

113. Anderson J, Papachristodoulou A. On validation and invalidation of biological models. BMC Bioinformatics. 2009;10:132.

114. UniProt Consortium T. UniProt: the universal protein knowledgebase. Nucleic Acids Res. 2018;46(5):2699.

115. Caspi R, Billington R, Keseler IM, Kothari A, Krummenacker M, Midford PE, et al. The MetaCyc database of metabolic pathways and enzymes - a 2019 update. Nucleic Acids Res. 2020;48(D1):D445-D53.

116. Sajed T, Marcu A, Ramirez M, Pon A, Guo AC, Knox C, et al. ECMDB 2.0: A richer resource for understanding the biochemistry of E. coli. Nucleic Acids Res. 2016;44(D1):D495-501.

117. Wittig U, Kania R, Golebiewski M, Rey M, Shi L, Jong L, et al. SABIO-RK--database for biochemical reaction kinetics. Nucleic Acids Res. 2012;40(Database issue):D790-6.

118. Kolesnikov N, Hastings E, Keays M, Melnichuk O, Tang YA, Williams E, et al. ArrayExpress update--simplifying data submissions. Nucleic Acids Res. 2015;43(Database issue):D1113-6.

119. Burley SK, Kurisu G, Markley JL, Nakamura H, Velankar S, Berman HM, et al. PDB-Dev: a Prototype System for Depositing Integrative/Hybrid Structural Models. Structure. 2017;25(9):1317-8.

120. Karr JR, Sanghvi JC, Macklin DN, Arora A, Covert MW. WholeCellKB: model organism databases for comprehensive whole-cell models. Nucleic Acids Res. 2013;41(Database issue):D787-92.

121. Medley JK, Goldberg AP, Karr JR. Guidelines for Reproducibly Building and Simulating Systems Biology Models. IEEE Trans Biomed Eng. 2016;63(10):2015-20.

122. Hucka M, Finney A, Sauro HM, Bolouri H, Doyle JC, Kitano H, et al. The systems biology markup language (SBML): a medium for representation and exchange of biochemical network models. Bioinformatics. 2003;19(4):524-31.

123. Le Novere N, Hucka M, Mi H, Moodie S, Schreiber F, Sorokin A, et al. The Systems Biology Graphical Notation. Nat Biotechnol. 2009;27(8):735-41.

124. Waltemath D, Karr JR, Bergmann FT, Chelliah V, Hucka M, Krantz M, et al. Toward Community Standards and Software for Whole-Cell Modeling. IEEE Trans Biomed Eng. 2016;63(10):2007-14.

125. Courtot M, Juty N, Knupfer C, Waltemath D, Zhukova A, Drager A, et al. Controlled vocabularies and semantics in systems biology. Mol Syst Biol. 2011;7:543.

126. Chalkley O, Purcell O, Grierson C, Marucci L. The genome design suite: enabling massive in-silico experiments to design genomes. bioRxiv. 2019:681270.

127. Kwiatkowska M, Norman G, Parker D, editors. PRISM 4.0: Verification of Probabilistic Real-Time Systems2011; Berlin, Heidelberg: Springer Berlin Heidelberg.

128. Iyengar S. Symbolic Systems Biology: Theory and Methods: Jones and Bartlett Learning; 2011 2011.

129. Ray O, Soh T, Inoue K. Analysing Pathways using ASP-Based approaches. In: Springer, editor. Proc 2010 Conf on Algebraic and Numeric Biology (ANB)2011. p. 167-83.

130. Bragagli S, Ray O, editors. Nonmonotonic Learning in Large Biological Networks. Proc 24th Int Conf on Inductive Logic Programming (ILP); 2015: Springer.

131. Rozanski R, Ray O, King R, Bragaglia S. Automating development of metabolic network models. Proc 13th Int Conf on Computational Methods in Systems Biology (CMSB): Springer; 2015.

132. Feig M, Sugita Y. Reaching new levels of realism in modeling biological macromolecules in cellular environments. J Mol Graph Model. 2013;45:144-56.

133. Feig M, Harada R, Mori T, Yu I, Takahashi K, Sugita Y. Complete atomistic model of a bacterial cytoplasm for integrating physics, biochemistry, and systems biology. J Mol Graph Model. 2015;58:1-9.

134. Sali A, Berman HM, Schwede T, Trewhella J, Kleywegt G, Burley SK, et al. Outcome of the First wwPDB Hybrid/Integrative Methods Task Force Workshop. Structure. 2015;23(7):1156-67.

135. Hyeon C, Thirumalai D. Capturing the essence of folding and functions of biomolecules using coarse-grained models. Nat Commun. 2011;2:487.

136. Ando T, Skolnick J. Crowding and hydrodynamic interactions likely dominate in vivo macromolecular motion. Proc Natl Acad Sci U S A. 2010;107(43):18457-62.

137. Solernou A, Hanson BS, Richardson RA, Welch R, Read DJ, Harlen OG, et al. Fluctuating Finite Element Analysis (FFEA): A continuum mechanics software tool for mesoscale simulation of biomolecules. PLoS Comput Biol. 2018;14(3):e1005897.

138. Castiglione F, Pappalardo F, Bianca C, Russo G, Motta S. Modeling biology spanning different scales: an open challenge. Biomed Res Int. 2014;2014:902545.

139. Lin C, Jain S, Kim H, Bar-Joseph Z. Using neural networks for reducing the dimensions of single-cell RNA-Seq data. Nucleic Acids Res. 2017;45(17):e156.

140. Ma J, Yu MK, Fong S, Ono K, Sage E, Demchak B, et al. Using deep learning to model the hierarchical structure and function of a cell. Nat Methods. 2018;15(4):290-8.

141. Yu MK, Ma J, Fisher J, Kreisberg JF, Raphael BJ, Ideker T. Visible Machine Learning for Biomedicine. Cell. 2018;173(7):1562-5.

142. Perdikaris P, Karniadakis GE. Model inversion via multi-fidelity Bayesian optimization: a new paradigm for parameter estimation in haemodynamics, and beyond. J R Soc Interface. 2016;13(118).

143. Alber M, Buganza Tepole A, Cannon WR, De S, Dura-Bernal S, Garikipati K, et al. Integrating machine learning and multiscale modeling-perspectives, challenges, and opportunities in the biological, biomedical, and behavioral sciences. NPJ Digit Med. 2019;2:115.

144. Smucker B, Krzywinski M, Altman N. Optimal experimental design. Nat Methods. 2018;15(8):559-60.

145. Bordbar A, McCloskey D, Zielinski DC, Sonnenschein N, Jamshidi N, Palsson BO. Personalized Whole-Cell Kinetic Models of Metabolism for Discovery in Genomics and Pharmacodynamics. Cell Syst. 2015;1(4):283-92.