


2D transformations (a.k.a. warping)



Course announcements

- Homework 1 posted on course website.
 - Due on February 5th at 23:59.
 - This homework is in Matlab.
 - How many of you have looked at/started/finished homework 1?
- Homework 2 will be posted tonight on the course website.
 - It will be due on February 19th at 23:59 pm.
 - Start early because it is much larger and more difficult than homework 1.
- Second theory quiz on course website and due on February 10th, at 23:59.
- Starting with quiz 3, release/due dates will be on Sunday.
- Reminder: All office hours are at the Smith Hall 200 conference room (not my office!).
- How was the first quiz?

Overview of today's lecture

- Reminder: image transformations.
- 2D transformations.
- Projective geometry 101.
- Transformations in projective geometry.
- Classification of 2D transformations.
- Determining unknown 2D transformations.
- Determining unknown image warps.

Slide credits

Most of these slides were adapted from:

- Kris Kitani (16-385, Spring 2017).

Reminder: image transformations


What is an image?



grayscale image

What is the range of the image function f ?

$f(\mathbf{x})$



domain $\mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix}$

A (grayscale) image is a 2D function.

What types of image transformations can we do?




Filtering



changes pixel *values*



Warping



changes pixel *locations*

What types of image transformations can we do?

F



Filtering



$$G(\mathbf{x}) = h\{F(\mathbf{x})\}$$

G



changes *range* of image function

F




Warping



$$G(\mathbf{x}) = F(h\{\mathbf{x}\})$$

G



changes *domain* of image function


Warping example: feature matching



Warping example: feature matching



Warping example: feature matching



- object recognition
- 3D reconstruction
- augmented reality
- image stitching

How do you compute the transformation?

Warping example: feature matching

Given a set of matched feature points:

$$\{x_i, x'_i\}$$

point in one
image

point in the
other image

and a transformation:


$$x' = f(x; p)$$

transformation
function

parameters

find the best estimate of the parameters

p



What kind of transformation functions f are there?

2D transformations

2D transformations



translation



rotation



aspect



affine



perspective




cylindrical


2D planar transformations




2D planar transformations




2D planar transformations




2D planar transformations




2D planar transformations



2D planar transformations




2D planar transformations




How would you implement rotation?

2D planar transformations



2D planar transformations



Polar coordinates...

$$x = r \cos(\phi)$$

$$y = r \sin(\phi)$$

$$x' = r \cos(\phi + \theta)$$

$$y' = r \sin(\phi + \theta)$$

Trigonometric Identity...

$$x' = r \cos(\phi) \cos(\theta) - r \sin(\phi) \sin(\theta)$$


$$y' = r \sin(\phi) \cos(\theta) + r \cos(\phi) \sin(\theta)$$

Substitute...

$$x' = x \cos(\theta) - y \sin(\theta)$$

$$y' = x \sin(\theta) + y \cos(\theta)$$

2D planar transformations



2D planar and linear transformations

$$\mathbf{x}' = f(\mathbf{x}; p)$$



$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \mathbf{M} \begin{bmatrix} x \\ y \end{bmatrix}$$

parameters p

point \mathbf{x}

2D planar and linear transformations

Scale

$$\mathbf{M} = \begin{bmatrix} s_x & 0 \\ 0 & s_y \end{bmatrix}$$

Flip across y

$$\mathbf{M} = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$$

Rotate

$$\mathbf{M} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

Flip across origin

$$\mathbf{M} = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$$


Shear

$$\mathbf{M} = \begin{bmatrix} 1 & s_x \\ s_y & 1 \end{bmatrix}$$

Identity


$$\mathbf{M} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

2D translation




How would you implement translation?

2D translation



2D translation



Projective geometry 101

Homogeneous coordinates

heterogeneous
coordinates

homogeneous
coordinates

$$\begin{bmatrix} x \\ y \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

← add a 1 here

- Represent 2D point with a 3D vector

Homogeneous coordinates


heterogeneous
coordinates

homogeneous
coordinates

$$\begin{bmatrix} x \\ y \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \stackrel{\text{def}}{=} \begin{bmatrix} ax \\ ay \\ a \end{bmatrix}$$


- Represent 2D point with a 3D vector
- 3D vectors are only defined up to scale

2D translation



2D translation

y



$$x' = x + t_x$$

$$y' = y + t_y$$


What about matrix representation using heterogeneous coordinates?

$$\begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \mathbf{M} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix}$$

x

2D translation using homogeneous coordinates

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} x + t_x \\ y + t_y \\ 1 \end{bmatrix}$$



Homogeneous coordinates

Conversion:

- heterogeneous \rightarrow homogeneous

$$\begin{bmatrix} x \\ y \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

- homogeneous \rightarrow heterogeneous

$$\begin{bmatrix} x \\ y \\ w \end{bmatrix} \Rightarrow \begin{bmatrix} x/w \\ y/w \end{bmatrix}$$

- scale invariance

$$\begin{bmatrix} x & y & w \end{bmatrix}^\top = \lambda \begin{bmatrix} x & y & w \end{bmatrix}^\top$$

Special points:

- point at infinity

$$\begin{bmatrix} x & y & 0 \end{bmatrix}$$

- undefined

$$\begin{bmatrix} 0 & 0 & 0 \end{bmatrix}$$

Projective geometry

image point in
pixel coordinates

$$\mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix}$$





image point in
homogeneous
coordinates

$$\mathbf{X} = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$



What does scaling \mathbf{X} correspond to?

Transformations in projective geometry

2D transformations in heterogeneous coordinates

Re-write these transformations as 3x3 matrices:

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

translation

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} ? \\ ? \\ ? \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

scaling

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} ? \\ ? \\ ? \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

rotation

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} ? \\ ? \\ ? \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

shearing

2D transformations in heterogeneous coordinates

Re-write these transformations as 3x3 matrices:

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

translation

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

scaling

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} & & \\ & ? & \\ & & \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

rotation

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} & & \\ & ? & \\ & & \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

shearing

2D transformations in heterogeneous coordinates

Re-write these transformations as 3x3 matrices:

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

translation

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

scaling

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} & & \\ & ? & \\ & & \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

rotation

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & \beta_x & 0 \\ \beta_y & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

shearing

2D transformations in heterogeneous coordinates

Re-write these transformations as 3x3 matrices:

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

translation

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

scaling

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} \cos \Theta & -\sin \Theta & 0 \\ \sin \Theta & \cos \Theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

rotation

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & \beta_x & 0 \\ \beta_y & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

shearing

Matrix composition

Transformations can be combined by matrix multiplication:

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \left(\begin{bmatrix} 1 & 0 & tx \\ 0 & 1 & ty \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos \Theta & -\sin \Theta & 0 \\ \sin \Theta & \cos \Theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} sx & 0 & 0 \\ 0 & sy & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$

$\mathbf{p}' = \quad ? \quad ? \quad ? \quad \mathbf{p}$

Matrix composition

Transformations can be combined by matrix multiplication:

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \left(\begin{bmatrix} 1 & 0 & tx \\ 0 & 1 & ty \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos \Theta & -\sin \Theta & 0 \\ \sin \Theta & \cos \Theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} sx & 0 & 0 \\ 0 & sy & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$

\mathbf{p}' = translation(t_x, t_y)

rotation(θ)


scale(s, s)

\mathbf{p}

Does the multiplication order matter?

Classification of 2D transformations

Classification of 2D transformations



Classification of 2D transformations


Name	Matrix	# D.O.F.
translation	$\left[\begin{array}{c c} \mathbf{I} & \mathbf{t} \end{array} \right]$?
rigid (Euclidean)	$\left[\begin{array}{c c} \mathbf{R} & \mathbf{t} \end{array} \right]$?
similarity	$\left[\begin{array}{c c} s\mathbf{R} & \mathbf{t} \end{array} \right]$?
affine	$\left[\begin{array}{c} \mathbf{A} \end{array} \right]$?
projective	$\left[\begin{array}{c} \tilde{\mathbf{H}} \end{array} \right]$?

Classification of 2D transformations

Translation:

$$\begin{bmatrix} 1 & 0 & t_1 \\ 0 & 1 & t_2 \\ 0 & 0 & 1 \end{bmatrix}$$

How many degrees of freedom?




Classification of 2D transformations

Euclidean (rigid):
rotation + translation

$$\begin{bmatrix} r_1 & r_2 & r_3 \\ r_4 & r_5 & r_6 \\ 0 & 0 & 1 \end{bmatrix}$$

Are there any values that are related?




Classification of 2D transformations

Euclidean (rigid):
rotation + translation

$$\begin{bmatrix} \cos \theta & -\sin \theta & r_3 \\ \sin \theta & \cos \theta & r_6 \\ 0 & 0 & 1 \end{bmatrix}$$

How many degrees of freedom?




Classification of 2D transformations

which other matrix values will
change if this increases?

Euclidean (rigid):
rotation + translation

$$\begin{matrix} \downarrow \\ \begin{bmatrix} \cos \theta & -\sin \theta & r_3 \\ \sin \theta & \cos \theta & r_6 \\ 0 & 0 & 1 \end{bmatrix} \end{matrix}$$




Classification of 2D transformations

what will happen to the image if this increases?



Euclidean (rigid):
rotation + translation

$$\begin{bmatrix} \cos \theta & -\sin \theta & r_3 \\ \sin \theta & \cos \theta & r_6 \\ 0 & 0 & 1 \end{bmatrix}$$




Classification of 2D transformations

what will happen to the
image if this increases?

Euclidean (rigid):
rotation + translation

$$\begin{bmatrix} \cos \theta & -\sin \theta & r_3 \\ \sin \theta & \cos \theta & r_6 \\ 0 & 0 & 1 \end{bmatrix}$$




Classification of 2D transformations

Similarity:
uniform scaling + rotation
+ translation

$$\begin{bmatrix} r_1 & r_2 & r_3 \\ r_4 & r_5 & r_6 \\ 0 & 0 & 1 \end{bmatrix}$$

Are there any values that are related?



Classification of 2D transformations


multiply these four by scale s



Similarity:
uniform scaling + rotation
+ translation

$$\begin{bmatrix} \cos \theta & -\sin \theta & r_3 \\ \sin \theta & \cos \theta & r_6 \\ 0 & 0 & 1 \end{bmatrix}$$

How many degrees of freedom?




Classification of 2D transformations

what will happen to the
image if this increases?

Similarity:
uniform scaling + rotation
+ translation

$$\begin{matrix} \downarrow \\ \begin{bmatrix} r_1 & r_2 & r_3 \\ r_4 & r_5 & r_6 \\ 0 & 0 & 1 \end{bmatrix} \end{matrix}$$




Classification of 2D transformations

Affine transform:
uniform scaling + shearing
+ rotation + translation

$$\begin{bmatrix} a_1 & a_2 & a_3 \\ a_4 & a_5 & a_6 \\ 0 & 0 & 1 \end{bmatrix}$$

Are there any values that are related?




Classification of 2D transformations

Affine transform:
uniform scaling + shearing
+ rotation + translation

$$\begin{bmatrix} a_1 & a_2 & a_3 \\ a_4 & a_5 & a_6 \\ 0 & 0 & 1 \end{bmatrix}$$

Are there any values that are related?

$$\begin{matrix} \text{similarity} \\ \begin{bmatrix} sr_1 & sr_2 \\ sr_3 & sr_4 \end{bmatrix} \end{matrix} \begin{matrix} \text{shear} \\ \begin{bmatrix} 1 & h_1 \\ h_2 & 1 \end{bmatrix} \end{matrix} = \begin{bmatrix} sr_1 + h_2 sr_2 & sr_2 + h_1 sr_1 \\ sr_3 + h_2 sr_4 & sr_4 + h_1 sr_3 \end{bmatrix}$$




Classification of 2D transformations

Affine transform:
uniform scaling + shearing
+ rotation + translation

$$\begin{bmatrix} a_1 & a_2 & a_3 \\ a_4 & a_5 & a_6 \\ 0 & 0 & 1 \end{bmatrix}$$

How many degrees of freedom?

$$\begin{bmatrix} sr_1 & sr_2 \\ sr_3 & sr_4 \end{bmatrix} \begin{bmatrix} 1 & h_1 \\ h_2 & 1 \end{bmatrix} = \begin{bmatrix} sr_1 + h_2 sr_2 & sr_2 + h_1 sr_1 \\ sr_3 + h_2 sr_4 & sr_4 + h_1 sr_3 \end{bmatrix}$$



Affine transformations


Affine transformations are combinations of

- arbitrary (4-DOF) linear transformations; and
- translations

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$

Properties of affine transformations:

- origin does not necessarily map to origin
- lines map to lines
- parallel lines map to parallel lines
- ratios are preserved
- compositions of affine transforms are also affine transforms



Does the last coordinate w ever change?

Affine transformations


Affine transformations are combinations of

- arbitrary (4-DOF) linear transformations; and
- translations

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$

Properties of affine transformations:

- origin does not necessarily map to origin
- lines map to lines
- parallel lines map to parallel lines
- ratios are preserved
- compositions of affine transforms are also affine transforms



Nope! But what does that mean?

How to interpret affine transformations here?

image point in
pixel coordinates

$$\mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix}$$





image point in
heterogeneous
coordinates

$$\mathbf{X} = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$



\mathbf{X} is a projection of a point
 \mathbf{P} on the image plane

Projective transformations

Projective transformations are combinations of


- affine transformations; and
- projective wraps

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$

How many degrees of freedom?

Properties of projective transformations:

- origin does not necessarily map to origin
- lines map to lines
- parallel lines do not necessarily map to parallel lines
- ratios are not necessarily preserved
- compositions of projective transforms are also projective transforms



Projective transformations

Projective transformations are combinations of


- affine transformations; and
- projective wraps

$$\begin{bmatrix} x' \\ y' \\ w' \end{bmatrix} = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \begin{bmatrix} x \\ y \\ w \end{bmatrix}$$

Properties of projective transformations:

- origin does not necessarily map to origin
- lines map to lines
- parallel lines do not necessarily map to parallel lines
- ratios are not necessarily preserved
- compositions of projective transforms are also projective transforms

8 DOF: vectors (and therefore matrices) are defined up to scale)



How to interpret projective transformations here?

image point in
pixel coordinates

$$\mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix}$$





image point in
heterogeneous
coordinates

$$\mathbf{X} = \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$




X is a projection of a point
 P on the image plane

Determining unknown 2D transformations

Determining unknown transformations



Suppose we have two triangles: ABC and DEF.



Determining unknown transformations

Suppose we have two triangles: ABC and DEF.


- What type of transformation will map A to D, B to E, and C to F?



Determining unknown transformations

Suppose we have two triangles: ABC and DEF.

- What type of transformation will map A to D, B to E, and C to F?
- How do we determine the unknown parameters?



Affine transform:
uniform scaling + shearing
+ rotation + translation


$$\begin{bmatrix} a_1 & a_2 & a_3 \\ a_4 & a_5 & a_6 \\ 0 & 0 & 1 \end{bmatrix}$$

How many degrees of freedom do we have?

Determining unknown transformations

Suppose we have two triangles: ABC and DEF.

- What type of transformation will map A to D, B to E, and C to F?
- How do we determine the unknown parameters?



unknowns

$$\mathbf{x}' = \mathbf{M}\mathbf{x}$$


point correspondences

- One point correspondence gives how many equations?
- How many point correspondences do we need?

Determining unknown transformations

Suppose we have two triangles: ABC and DEF.

- What type of transformation will map A to D, B to E, and C to F?
- How do we determine the unknown parameters?




unknowns

$$\mathbf{x}' = \mathbf{M}\mathbf{x}$$


point correspondences

How do we solve this for \mathbf{M} ?



Least Squares Error

$$E_{\text{LS}} = \sum_i \|\mathbf{f}(\mathbf{x}_i; \mathbf{p}) - \mathbf{x}'_i\|^2$$




Least Squares Error

$$E_{\text{LS}} = \sum_i \left\| \mathbf{f}(x_i; \mathbf{p}) - x'_i \right\|^2$$

What is this?

What is this?

What is this?



$$\|x\| = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2}$$

Least Squares Error


$$E_{\text{LS}} = \sum_i \left\| \mathbf{f}(\mathbf{x}_i; \mathbf{p}) - \mathbf{x}'_i \right\|^2$$

Euclidean
(L2) norm

squared!


predicted
location

measured
location



Least Squares Error

$$E_{\text{LS}} = \sum_i \underbrace{\|f(x_i; p) - x'_i\|}_{\text{Residual (projection error)}}^2$$




Least Squares Error

$$E_{\text{LS}} = \sum_i \|\mathbf{f}(\mathbf{x}_i; \mathbf{p}) - \mathbf{x}'_i\|^2$$

What is the free variable?

What do we want to optimize?



Find parameters that minimize squared error

$$\hat{\mathbf{p}} = \arg \min_{\mathbf{p}} \sum_i \|\mathbf{f}(\mathbf{x}_i; \mathbf{p}) - \mathbf{x}'_i\|^2$$

General form of linear least squares

(**Warning:** change of notation. \mathbf{x} is a vector of parameters!)

$$\begin{aligned} E_{\text{LLS}} &= \sum_i |\mathbf{a}_i \mathbf{x} - \mathbf{b}_i|^2 \\ &= \|\mathbf{A} \mathbf{x} - \mathbf{b}\|^2 \quad (\text{matrix form}) \end{aligned}$$

Determining unknown transformations

Affine transformation:

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} p_1 & p_2 & p_3 \\ p_4 & p_5 & p_6 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

Why can we drop
the last line?

Vectorize transformation
parameters:

$$\begin{bmatrix} x' \\ y' \\ x' \\ y' \\ \vdots \\ x' \\ y' \end{bmatrix} = \begin{bmatrix} x & y & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & x & y & 1 \\ x & y & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & x & y & 1 \\ \vdots & & & \vdots & & \\ x & y & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & x & y & 1 \end{bmatrix} \begin{bmatrix} p_1 \\ p_2 \\ p_3 \\ p_4 \\ p_5 \\ p_6 \end{bmatrix}$$

Stack equations from point
correspondences:

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} x & y & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & x & y & 1 \end{bmatrix} \begin{bmatrix} p_1 \\ p_2 \\ p_3 \\ p_4 \\ p_5 \\ p_6 \end{bmatrix}$$

$$\underbrace{\begin{bmatrix} x' \\ y' \end{bmatrix}}_{\mathbf{b}} = \underbrace{\begin{bmatrix} x & y & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & x & y & 1 \end{bmatrix}}_{\mathbf{A}} \underbrace{\begin{bmatrix} p_1 \\ p_2 \\ p_3 \\ p_4 \\ p_5 \\ p_6 \end{bmatrix}}_{\mathbf{x}}$$

Notation in system form:

General form of linear least squares

(Warning: change of notation. \mathbf{x} is a vector of parameters!)

$$\begin{aligned} E_{\text{LLS}} &= \sum_i |\mathbf{a}_i \mathbf{x} - \mathbf{b}_i|^2 \\ &= \|\mathbf{A} \mathbf{x} - \mathbf{b}\|^2 \quad (\text{matrix form}) \end{aligned}$$

This function is quadratic.

How do you find the root of a quadratic?

Solving the linear system

Convert the system to a linear least-squares problem:

$$E_{LLS} = \|\mathbf{A}\mathbf{x} - \mathbf{b}\|^2$$

Expand the error:

$$E_{LLS} = \mathbf{x}^\top (\mathbf{A}^\top \mathbf{A}) \mathbf{x} - 2\mathbf{x}^\top (\mathbf{A}^\top \mathbf{b}) + \|\mathbf{b}\|^2$$

Minimize the error:

Set derivative to 0 $(\mathbf{A}^\top \mathbf{A})\mathbf{x} = \mathbf{A}^\top \mathbf{b}$

Solve for \mathbf{x} $\mathbf{x} = (\mathbf{A}^\top \mathbf{A})^{-1} \mathbf{A}^\top \mathbf{b}$ ←

In Matlab:

$$\mathbf{x} = \mathbf{A} \setminus \mathbf{b}$$

Note: You almost never want to compute the inverse of a matrix.

Linear least squares estimation only works when the transform function is ?

Linear least squares estimation only works when the transform function is **linear!** (duh)


Also doesn't deal well with outliers

Determining unknown image warps

Determining unknown image warps

Suppose we have two images.


- How do we compute the transform that takes one to the other?



Forward warping

Suppose we have two images.

- How do we compute the transform that takes one to the other?




1. Form enough pixel-to-pixel correspondences between two images ← later lecture
2. Solve for linear transform parameters as before
3. Send intensities $f(x, y)$ in first image to their corresponding location in the second image

Forward warping

Suppose we have two images.

- How do we compute the transform that takes one to the other?



what is the problem with this?


1. Form enough pixel-to-pixel correspondences between two images
2. Solve for linear transform parameters as before
3. Send intensities $f(x, y)$ in first image to their corresponding location in the second image



Forward warping

Pixels may end up between two points


- How do we determine the intensity of each point?



Forward warping

Pixels may end up between two points

- How do we determine the intensity of each point?
- ✓ We distribute color among neighboring pixels (x',y') (“splatting”)




- What if a pixel (x',y') receives intensity from more than one pixels (x,y) ?

Forward warping

Pixels may end up between two points

- How do we determine the intensity of each point?
- ✓ We distribute color among neighboring pixels (x',y') (“splatting”)




- What if a pixel (x',y') receives intensity from more than one pixels (x,y) ?
- ✓ We average their intensity contributions.

Inverse warping

Suppose we have two images.

- How do we compute the transform that takes one to the other?



what is the problem with this?


1. Form enough pixel-to-pixel correspondences between two images
2. Solve for linear transform parameters as before, then compute its inverse
3. Get intensities $g(x', y')$ in the second image from point $(x, y) = T^{-1}(x', y')$ in first image



Inverse warping

Pixel may come from between two points


- How do we determine its intensity?



Inverse warping


Pixel may come from between two points

- How do we determine its intensity?
- ✓ Use interpolation




Bilinear interpolation

Grayscale example



1. Interpolate to find R_2
2. Interpolate to find R_1
3. Interpolate to find P



In matrix form (with adjusted coordinates)

$$f(x, y) \approx \begin{bmatrix} 1-x & x \end{bmatrix} \begin{bmatrix} f(0,0) & f(0,1) \\ f(1,0) & f(1,1) \end{bmatrix} \begin{bmatrix} 1-y \\ y \end{bmatrix}.$$


In Matlab:

call `interp2`

Forward vs inverse warping

Suppose we have two images.

- How do we compute the transform that takes one to the other?




Pros and cons of each?

Forward vs inverse warping

Suppose we have two images.

- How do we compute the transform that takes one to the other?



- Inverse warping eliminates holes in target image
- Forward warping does not require existence of inverse transform

References

Basic reading:

- Szeliski textbook, Section 3.6.

Additional reading:

- Hartley and Zisserman, “Multiple View Geometry in Computer Vision,” Cambridge University Press 2004.
a comprehensive treatment of all aspects of projective geometry relating to computer vision, and also a very useful reference for the second part of the class.
- Richter-Gebert, “Perspectives on projective geometry,” Springer 2011.
a beautiful, thorough, and very accessible mathematics textbook on projective geometry (available online for free from CMU’s library).