

REVIEW Article

Porosome in Cystic Fibrosis

Bhanu P. Jena*

Wayne State University School of Medicine, Department of Physiology, Detroit, MI, USA

*Corresponding author: Bhanu P. Jena, Ph.D., D.Sc., (dr. h.c. mult.), George E. Palade University Professor, Distinguished Professor, Department of Physiology, Wayne State University School of Medicine, Director of the NanoBioScience Institute, 540 E. Canfield, 5245 Gordon Scott Hall, Detroit, MI 48201-1928, Phone: 313-577-1532, Fax: 313-993-4177, E-mail: bjena@med.wayne.edu

Submitted: June 09, 2014; Revised: Sept. 04, 2014; Accepted: Sept. 05, 2014; Published: Sept. 28, 2014;
Citation: Jena B. Porosome in Cystic Fibrosis. Discoveries 2014, Jul-Sep; 2(3): e24. DOI: 10.15190/d.2014.16

ABSTRACT

Macromolecular structures embedded in the cell plasma membrane called 'porosomes', are involved in the regulated fractional release of intravesicular contents from cells during secretion. Porosomes range in size from 15 nm in neurons and astrocytes to 100-180 nm in the exocrine pancreas and neuroendocrine cells. Porosomes have been isolated from a number of cells, and their morphology, composition, and functional reconstitution well documented. The 3D contour map of the assembly of proteins within the porosome complex, and its native X-ray solution structure at sub-nm resolution has also advanced. This understanding now provides a platform to address diseases that may result from secretory defects. Water and ion binding to mucin impart hydration, critical for regulating viscosity of the mucus in the airways epithelia. Appropriate viscosity is required for the movement of mucus by the underlying cilia. Hence secretion of more viscous mucus prevents its proper transport, resulting in chronic and fatal airways disease such as cystic fibrosis (CF). CF is caused by the malfunction of CF transmembrane conductance regulator (CFTR), a chloride channel transporter, resulting in viscous mucus in the airways. Studies in mice lacking functional CFTR secrete highly viscous mucous that adhered to the epithelium. Since CFTR is known to interact with the t-SNARE protein syntaxin-1A, and with the chloride channel CLC-3, which are also components of the porosome complex, the interactions between CFTR and the

porosome complex in the mucin-secreting human airway epithelial cell line Calu-3 was hypothesized and tested. Results from the study demonstrate the presence of approximately 100 nm in size porosome complex composed of 34 proteins at the cell plasma membrane in Calu-3 cells, and the association of CFTR with the complex. In comparison, the nuclear pore complex measures 120 nm and is comprised of over 500 protein molecules. The involvement of CFTR in porosome-mediated mucin secretion is hypothesized, and is currently being tested.

Keywords: Porosome Complex, CFTR, mucin secretion

SUMMARY

1. Introduction
2. Porosome in Calu-3 cell
3. Ongoing studies

1. Introduction

It is well established that cup-shaped macromolecular lipoprotein structures called *porosomes* are secretory portals embedded in the cell plasma membrane in cells, where membrane-bound secretory vesicles transiently dock and fuse to expel intravesicular contents during secretion¹⁻¹⁰. Porosomes have been isolated from a number of cells, including the exocrine pancreas^{5,6} (Figure 1), neurons³ (Figure 2), and in the mucin-secreting human airway epithelial cell line Calu-3 (Figure




Figure 1. Atomic force microscopy (AFM) micrographs demonstrate the presence of microvilli and interspersed mucin-secreting porosomes at plasma membrane in Calu-3 cells¹¹. Microvilli measuring on average 96 nm in thickness (mean ± SEM; 96 ± 3.3, n=50) are densely packed at the cell plasma membrane exposed to the medium, and is demonstrated both in low (A-C) and high (D-H) resolution AFM images. Interspersed among the microvilli are the 102 nm in diameter porosome openings (mean ± SEM; 102 ± 3, n=50) shown in figure G (red and green arrowheads). Similarly, the microvilli shown in figure H (red and green arrowheads) demonstrate some that appear coiled around each other, possibly as a consequence of secreted mucus¹¹. ©Bhanu Jena.

3,4)¹¹. The morphology, composition, and reconstitution of porosomes in the exocrine pancreas (Figure 5-7) and in neurons are well documented²⁻¹², and the 3D contour map of the assembly of proteins within the structure has also been determined in great detail¹⁰. This new understanding of the secretory machinery in cells now provides a platform to address diseases resulting from secretory defects. The structure, and composition of the porosome complex in Calu-3 cells expressing cystic fibrosis (CF) transmembrane conductance regulator (CFTR) has been determined for the first time¹¹, with promise to help better understand cystic fibrosis. CFTR is a plasma membrane chloride selective cyclic AMP-activated ion channel, localized at the apical membrane of secretory epithelial cells, including the conducting airways¹³. Besides mediating the secretion of Cl⁻, CFTR also regulates several other transport proteins, including K⁺ channels, aquaporin water channels, anion exchangers, the membrane fusion protein syntaxin-1A, and sodium bicarbonate transporters¹⁴⁻²⁶. Accordingly, studies show that CFTR and its associated proteins are present in large macromolecular signaling complexes via scaffolding proteins containing PDZ domains^{13,25,27}. The C-terminus of CFTR in humans contains the sequence Asp-Thr-Arg-Leu, that mediate binding to several PDZ domain proteins¹³. For example, ezrin

and moesin present in the Calu-3 porosome complex¹¹ are also known CFTR-PDZ binding protein¹³. In addition, CFTR has several other regions that mediate protein-protein interactions, such as a domain at its N-terminus that binds to syntaxin-1A and SNAP-23^{15,23}. CFTR also contains a protein phosphatase-2A (PP2A)-binding, and an AMP kinase (AMPK)-binding domain¹². Similarly, CFTR has a regulatory domain that is a substrate to both protein kinase A (PKA) and C (PKC)²⁸. These interactions facilitate CFTR to form large CFTR-associated macromolecular signaling complexes at the plasma membrane. CF as a disease was first identified as cysts observed in the pancreas and the highly viscous mucus found in the lung of patients. However, since discovery that these observed defects are a result of a dysfunction of the CFTR chloride channel^{29,30}, there has been little progress in our understanding of the link between CFTR dysfunction and the secretion of such highly viscous mucin in the lung of CF patients³¹. The surface of the airways is coated with a thin film of mucous composed of essentially mucin, salt, proteases, antioxidants, and antibodies^{31,32}. Mucin lubricates, trap foreign particles and pathogens, and assists in the clearance of foreign particles from the airways via ciliary transport^{31,32}. A key property of mucus is its appropriate viscosity that enables its movement by the underlying cilia. Secretion of




Figure 2. Representative electron micrographs of Calu-3 cells in culture demonstrating the presence of microvilli (MV) and porosomes (P) at the cell plasma membrane [11]. (A) Calu-3 cells demonstrate the presence of dense microvilli and scattered porosomes at the cell plasma membrane. (B-D) Note the flask-shaped porosomes measuring nearly 100 nm in diameter (E) and from 200-300 nm in depth, with openings to the cell surface (red arrowhead). Mucus (C), is found at the opening of the porosome to the cell exterior. Of the two porosomes shown in (D), the one to the center appears to be sectioned right through the center of the organelle, where as the porosome to the left, has been sectioned at its base. (E) Similar to the AFM images in Figure 1, the microvilli measure on average 92 nm in diameter [11]. ©Bhanu Jena.

more viscous mucus disallows its proper transport, resulting in chronic and fatal airways disease such as CF³². Similar to other secretory cells that undergo secretory vesicle volume increase during secretion³⁵⁻⁴⁵, goblet cells of the airways epithelia that store mucin in a dehydrated state within membrane-bound secretory granules are no exception. Since vesicle swelling is a requirement for cell secretion⁴¹, and both ion channels and water channels or aquaporins regulate this process^{42,44}, altered chloride transport would impair secretory vesicle hydration and optimal release. Furthermore, recent studies in mice lacking functional CFTR³¹ showed that these animals secrete highly viscous mucous that adhered to the epithelium. Since CFTR is known to interact with syntaxin-1A, chloride channel CLC-3, and aquaporins¹⁴⁻²⁶, which are components of the porosome complex^{1,2,5-7,46}, the possible interactions between CFTR and the porosome in goblet cells was hypothesized and tested in a recent study¹¹. Results from the study demonstrate the presence of approximately 100 nm in size porosomes and microvilli at the surface of the plasma membrane in Calu-3 cells (Figure 1,2)¹¹.

The t-SNARE SNAP-25 specific antibody conjugated to protein A-sepharose® has been utilized to isolate the porosome complex from Calu-3 cells¹¹. For each immunoisolation, 1 mg of Triton-Lubrol-solubilized Calu-3 cells was used. The Triton/Lubrol solubilization buffer contained 0.5% Lubrol, 0.5% Triton X-100, 1 mM benzamidine, 5 mM Mg-ATP, and 5 mM EDTA in PBS at pH 7.5, supplemented with protease inhibitor mix (Sigma, St. Louis, MO). Ten micrograms of SNAP-25 antibody conjugated to the protein A-sepharose® were incubated with the 1 mg of the solubilized cells for 1 h at room temperature followed by three washes of 10 volumes of wash buffer (500 mM NaCl, 10 mM Tris, 2 mM EDTA, pH 7.5). The immunoprecipitated porosome attached to the immunosepharose beads was eluted using low pH buffer (pH 3.5) to dissociate the complex from the antibody bound to the bead, and the eluted sample immediately returned to neutral pH and stored at -80 degrees¹¹. A combination of proteomics, Western blot analysis, and immunocytochemistry, were all used to determine the composition and distribution of the porosome complex in Calu-3




Figure 3. Photon correlation spectroscopy (PCS) demonstrate the immunisolated porosome complex from Calu-3 cells to measure on average 271 nm (trimmers), and both immunoprecipitation and immunoblot analysis demonstrate the interaction of CFTR with the porosome complex in the cell¹¹. (A) PCS on isolated porosomes from Calu-3 cells demonstrate an average size of 271.2 nm. (B) Immunoblot analysis using CFTR-specific antibody of CFTR-expressing HEK cell proteins resolved using SDS-PAGE, followed by electrotransfer to nitrocellulose membrane, demonstrates the presence of a 180 kD band representing CFTR (lane 1, positive control). SDS-PAGE resolved immunisolated porosome complexes also demonstrate immunopositive for CFTR (lane 2). (C) Immunoblot analysis of the total Calu-3 cell homogenate (CH) and isolated porosome complex (P), demonstrate the presence of porosome proteins actin, Gαi3, and vimentin. Note the enriched presence of the proteins in the porosome complex. (D) Similarly, immunisolated CFTR complex using the CFTR-specific antibody, results in the pull-down of porosome associated proteins such as Syntaxin-1A (present as 70 kDa t-/v-SNARE complex), SNAP-25 (present as 70 kDa t-/v-SNARE complex), SNAP-23, present as 68 kDa t-/v-SNARE complex, and actin¹¹. ©Bhanu Jena.

cells¹¹. Proteomic analysis of isolated Calu-3 porosomes using mass spectrometry demonstrate the presence of CFTR as well as several proteins found in the neuronal porosome complex, including Syntaxin-1A, actin, rabs, heterotrimeric G-protein, and the GTPase activating protein GAP (Table I)¹¹. Immunoblot analysis (Figure 3) of the isolated Calu-3 porosome complex, and immunocytochemistry (Figure 4) further confirms CFTR association with the porosome complex, reflecting important implication of CFTR in both normal mucus secretion in the airway epithelium in health, and in the impaired state in CF disease. In the past two decades, employing a combination of approaches including AFM, biochemistry, molecular biology, electrophysiology, EM, mass spectrometry, SAXS analysis, and database searches such as STRING 9.1 of known functional and predicted protein-protein interactions, further structural details of the porosome complex have been determined¹⁻¹². Although great progress have been made in our understanding of the porosome, of

Ca²⁺ and SNARE-mediated membrane fusion⁴⁷⁻⁷⁶, and on secretory vesicle volume regulation required for regulated fractional release of intravesicular contents³⁵⁻⁴⁵ during cell secretion, a molecular level understanding of porosome-mediated secretion in mucin-secreting cells remains to be determined. Therefore, a clear understanding of the porosome in mucin secreting in Calu-3 cells, and the role of CFTR in porosome-mediated mucin secretion is critical in revealing how mucin secretion is precisely regulated.

2. Porosome in Calu-3 cell

High resolution imaging using atomic force microscopy (AFM) (Figure 1) reveal in great detail the surface topology of Calu-3 cells, demonstrate the presence of approximately 102 nm in diameter porosome openings (mean ± SEM; 102.4 nm ± 3.0 nm), and 96 nm thick (mean ± SEM; 96 nm ± 3.3 nm) microvilli, at the cell plasma membrane. Nearly the entire cell surface is covered with the




Figure 4. Schematic drawing depicting the evidence view of predicted interactions between identified proteins within the mucous-secreting Calu-3 porosome proteome and other regulatory proteins. These interactions are generated from inputs of the identified proteins in the Calu-3 porosome, using STRING 9.1^{7,11}. STRING 9.1 is a database of known functional and predicted protein-protein interactions. The interactions include direct (physical) and indirect (functional) associations derived from genomic, high-throughput, conserved co-expression, and earlier knowledge. Note the two clusters of protein-protein interactions identified in the porosome complex. The one cluster to the top, and most likely present at the apical end of the porosome cup are cytoskeletal structure and signalling proteins. The bottom cluster represents proteins that are primarily involved in membrane fusion including SNARE proteins and CFTR, and therefore their location would be at the base of the porosome complex facing the cytosol. The confidence of the predicted functional interactions shown are >99%. ©Bhanu Jena.

microvilli, with interspersed porosome openings. In certain areas of the cell surface devoid of microvilli or porosome openings, cytoskeletal structures underlying the cell plasma membrane are observed (Figure 1B)¹¹. Transmission electron microscopy (TEM) performed on Calu-3 cells confirms the AFM results (Figure 2), demonstrating the presence of dense microvilli (Figure 2A), and cup-shaped porosomes (Figure 2B-D) at the cell plasma membrane. Immunisolated Calu-3 porosome complexes demonstrate a particle size of

approximately 300 nm (Figure 3A) using photon correlation spectroscopy (PCS), possibly a result of trimerization in their isolated state in suspension¹¹. Immunoblot analysis of isolated Calu-3 porosome complex (Figure 3B), demonstrate the co-association of CFTR with the porosome complex that contains among other proteins Gαi3, actin, and vimentin (Figure 3C). Similarly, when CFTR is immunisolated from solubilized Calu-3 cells, the porosome complex is co-immunisolated with CFTR, as demonstrated by the presence of




Figure 5. Small angle X-ray solution scattering structure of a native 35 nm synaptic vesicle (violet) docked with a 15 nm neuronal porosome complex (pink) at the presynaptic membrane. Note the prominent central plug of the porosome, which has been implicated in the rapid closing and opening of the complex⁸⁰. ©Bhanu Jena.

porosome-associated proteins Syntaxin-1A, SNAP-25, SNAP-23, actin, and vimentin (Figure 3D). Mass spectrometry on the immunisolated porosome complex from Calu-3 cells, demonstrates the presence of various porosome-associated proteins (found in the neuronal and pancreatic porosome complexes) as well as CFTR¹¹. Furthermore, mass spectrometry results confirm the interaction between CFTR and the porosome complex in Calu-3 cells, as determined using immunoprecipitation and immunoblot analysis. Using the STRING 9.1 database search⁷⁷ similar to the neuronal porosome complex¹², two clusters of protein-protein interactions within the mucin-secreting Calu-3 porosome have been identified (Figure 4). The cluster to the top-right in figure 4, represent primarily cytoskeletal and signaling proteins, whereas the bottom-left cluster represents proteins that are primarily involved in membrane fusion, including SNAREs, ion channels and CFTR. Therefore the bottom cluster is located at the porosome base facing the cytosol where mucin-containing vesicles dock and fuse, and the top cluster at the porosome opening to the outside.

Therefore in these studies, the porosome proteome in human airways epithelia has been determined. The interaction between CFTR and the porosome complex in the human airways epithelia is further demonstrated. The possible regulation by CFTR on the quality of mucus secretion via the porosome complex at the cell plasma membrane is hypothesized. These new findings will facilitate understanding of CFTR-porosome interactions influencing mucus secretion, and provide critical insights into the etiology of CF disease.

In view of this, an integrated approach is being used to characterize the molecular architecture of the mucin-secreting porosome complex of the human airways epithelia cell line Calu-3; determine the distribution of CFTR and its interaction with proteins and lipids within the mucin-secreting porosome complex; characterize the molecular architecture of the mucin-secreting porosome; and build and test a functional architectural model to determine how SNAREs, lipids, and calcium, establish continuity between the secretory vesicle membrane and the porosome. These studies will allow a molecular understanding of the porosome function in mucin secretion, and the role of CFTR in the process.

3. Ongoing studies

Among the 34 core porosome proteins in mucin-secreting Calu-3 cells, are included CFTR, actin, vimentin, annexin, filamin, G α 3, tubulin, syntaxin-1A, profiling, ezrin, spectrin, chloride channels CLC-1 and CLC-3, rab1A and rab3A, myosin, SNAP-25, and the ADP-ribosylation factor ARF3¹¹. It is anticipated that due to the nature of proteome studies and the dynamics of porosomes¹² that this initial analysis includes most of the core porosome proteins, but not all the peripheral proteins associated with the complex. The STRING 9.1⁷⁷ database search utilizing known physical and functional associations between proteins suggests additional candidates likely involved in protein-protein interactions within the Calu-3 porosome complex (Figure 4). Similarly, preliminary lipidomic studies using lipid overlay assays on isolated Calu-3 porosome complex, demonstrate the enriched presence of PA and PIP2. Furthermore, interactions of PA and other polyphosphoinositides with syntaxin-1A, and their involvement in cell secretion have previously been reported⁷⁸. These




Figure 6 Schematic drawing depicting the presence and increased association of dynamin with the porosome complex following stimulation of neurotransmitter release¹², which may be similar in the mucin-secreting Calu-3 cell. Following stimulation of secretion, synaptic vesicles would dock at the porosome base, develop intravesicular pressure via active transport of water through water channels or aquaporins (AQP) at the vesicle membrane, transiently fuse at the porosome base via SNAREs and calcium, and expel neurotransmitters. After secretion, NSF an ATPase, and dynamin a GTPase, would work synchronously to disassembly t-/v-SNARE complexes and fission the neck of fused vesicles at the porosome base respectively. By this mechanism, partially empty vesicles could go through multiple rounds of docking-fusion-expulsion-dissociation. Unlike protein and peptide containing vesicles, synaptic vesicles have neurotransmitter transporters at the vesicle membrane to rapidly refill vesicles¹². In case of the Calu-3 cell, once mucin containing vesicles empty, they may recycle via the endosome or lysosomal pathway. ©Bhanu Jena.

observations indicate the importance of lipid interactions in the structure and function of the porosome. Based on this information, and since mucin containing vesicles dock at the porosome base, and the t-SNARE syntaxin-1A and SNAP-25 associated with CFTR, rab3C, CLC3, and SNAP91 are among other proteins present at the porosome base (Figure 4), it could be surmised that PA and PIP2 together with these proteins are present at the porosome base. The protein cluster composed of cytoskeletal and signaling proteins on the other hand, is likely associated with the porosome opening to the outside of the cell, regulating dilation of the porosome opening during mucin secretion. Since in the presence of the actin depolymerizing agent cytochalasin, there is loss of

mucin secretion (unpublished observation) as in the case of the exocrine pancreas⁷ or growth hormone secreting cells¹, further supports the presence of signaling and motor proteins at the porosome opening, that regulate dilation of the porosome opening and mucin secretion.

Unlike individual proteins or lipids, determination of the atomic structure of such dynamic macromolecular lipoprotein complexes such as the porosome, poses a difficult challenge, requiring the use of several experimental and computational approaches to maximize resolution and accuracy. Although recently the isolated Calu-3 porosome has been functionally reconstituted as in case of the exocrine pancreas or neurons, further functional reconstitution studies are under way

- ◆ Mucin-secreting porosomes are 100 nm cup-shaped secretory portal at the plasma membrane; the porosome has been isolated from Calu-3 cells and its proteome determined
- ◆ Porosome structure has been determined using EM and AFM
- ◆ Distribution of lipids and various proteins especially CFTR within the porosome complex is under way

using the established lipid bilayer EPC9 system⁶. These experiments further determine if the entire porosome complex has been isolated prior to determination of its composition and molecular structure-function. Electron microscopy (EM) especially single particle cryo electron tomography, small angle X-ray solution scattering (SAXS), supra resolution microscopy (SRM), and AFM analyses are being used and complemented by techniques from structural mass spectrometry and proteomics to obtain molecular details of the mucin-secreting porosome structure. The mass spectrometry studies include subunit stoichiometry, interacting subunits, and site of contact between subunits. Changes to porosome subunit composition and subunit interactions during the secretory process are being studied using these approaches. CXL-MS and multiple quantitative mass spectrometry techniques are being utilized to determine details of the protein-protein interactions within the native mucin-secreting porosome complex, which is central to building a structural model of the complex for a molecular understanding of its structure-function.

New and recently developed crosslinkers⁷⁹ combined with tandem mass spectrometry are being carried out, which will provide identities of interacting subunits and provide the identities of specific residues crosslinked both between and within subunits in the porosome complex. Results from these studies will provide information on interaction domains and distance constraints on protein structures. Quantitative mass spectrometry using iTRAQ are also being carried out, which will provide additional information on changes in porosome subunits composition and dynamics, as a function of the secretion status of the organelle. Immuno-AFM⁵, immuno-EM, and SAXS⁸⁰ (Figure 5) studies on isolated Calu-3 porosomes as in porosomes of the exocrine pancreas and neurons, are being performed to determine the distribution of

some of the major proteins within the complex. Similar to studies using SRM on the nuclear pore complex⁸¹, SRM is being employed to obtain additional information on the structure of the mucin-secreting porosome complex. Finally, computational approaches are being employed, such as coarse-grain molecular docking studies⁸²⁻⁹⁷, homology modeled interactions⁹⁸⁻¹⁰⁰, and fitting of known atomic structures of protein-protein interactions and complexes¹⁰¹⁻¹⁰⁸. It is becoming increasingly clear that the ultrastructural and mass spectrometry methods show promise in providing complementary information and the high degree of cross-validation required to build an accurate structural model of the mucin-secreting porosome complex. Collectively, the outlined studies briefly discussed here will enable an understanding at the molecular level, the elegant mechanism of porosome-mediated secretion (Figure 6) in Calu-3 and other cells.

Conflict of Interest

The authors do not declare any conflict of interest.

Acknowledgements

The author thanks members of his laboratory for lively discussions on the subject. This work was supported by the NIH grant R01EB000303 (to BPJ).

References:

1. Cho SJ, Jeftinija K, Glavaski A, Jeftinija S, Jena BP, Anderson LL. Structure and dynamics of the fusion pores in live GH-secreting cells revealed using atomic force microscopy. *Endocrinology* 2002;143:1144-8.
2. Cho SJ, Quinn AS, Stromer MH, Dash S, Cho J, Taatjes DJ, et al. Structure and dynamics of the fusion pore in live cells. *Cell Biol Int* 2002;26:35-42.
3. Cho WJ, Jeremic A, Rognlien KT, Zhvania MG, Lazrshvili I, Tamar B, et al. Structure, isolation, composition and reconstitution of the neuronal fusion pore. *Cell Biol Int* 2004;28:699-708.
4. Cho WJ, Jeremic A, Jin H, Ren G, Jena BP. Neuronal fusion pore assembly requires membrane cholesterol.

- Cell Biol Int 2007;31:1301-8.
5. Jena BP, Cho SJ, Jeremic A, Stromer MH, Abu-Hamdah R. Structure and composition of the fusion pore. *Biophys J* 2003;84:1-7.
 6. Jeremic A, Kelly M, Cho SJ, Stromer MH, Jena BP. Reconstituted fusion pore. *Biophys J* 2003;85:2035-43.
 7. Schneider SW, Sriharan KC, Geibel JP, Oberleithner H, Jena BP. Surface dynamics in living acinar cells imaged by atomic force microscopy: Identification of plasma membrane structures involved in exocytosis. *Proc Natl Acad Sci U S A* 1997;94:316-21.
 8. Jena BP. Molecular machinery and mechanism of cell secretion. *Exp Biol Med* 2005;230:307-19.
 9. Jena BP. Secretion machinery at the cell plasma membrane. *Curr Opin Struct Biol* 2007;17:437-43.
 10. Cho WJ, Ren G., Jena BP. EM 3D contour maps provide protein assembly at the nanoscale within the neuronal porosome complex. *J Microscopy* 2008;232:106-11.
 11. Hou X, Lewis KT, Wu Q, Wang S, Chen X, Flack A, Mao G, Taatjes DJ, Su F, Jena BP. Proteome of the porosome complex in human airways epithelia: Interaction with the cystic fibrosis transmembrane conductance regulator (CFTR). *Journal of Proteomics* 96: 82-91, 2014.
 12. Lee J-S, Jeremic A, Shin L, Cho WJ, Chen X, Jena BP. Neuronal porosome proteome: molecular dynamics and architecture. *J Proteomics* 2012;75:3952-3962.
 13. Guggino WB, Stanton BA. New insights into cystic fibrosis: molecular switches that regulate CFTR. *Nature Rev* 2006;7:426-436.
 14. Schwiebert EM, Egan ME, Hwang T-H, Fulmer SB, Allen SS, Cutting GR, Guggino WB. CFTR regulates outwardly rectifying chloride channels through an autocrine mechanism involving ATP. *Cell* 1995;81:1063-1073.
 15. Naren AP, Nelson DJ, Xie W, Jovov B, Tousson A, Pevsner J, Bennett MK, Benos DJ, Quick MW, Kirk KL. Regulation of CFTR chloride channels by syntaxin and Munc 18 isoforms. *Nature* 1997;390:302-305.
 16. Naren AP, Cobb B, Li C, Roy K, Nelson D, Heda GD, Liao J, Kirk KL, Sorscher EJ, Hanrahan J, Clancy JP. A macromolecular complex of β 2AR, CFTR and EBP50 is regulated by protein kinase A. *Proc Natl Acad Sci USA* 2003;100:342-346.
 17. Stutts MJ, Rossier BC, Boucher RC. Cystic fibrosis transmembrane conductance regulator inverts protein kinase A-mediated regulation of epithelial sodium channel single channel kinetics. *J Biol Chem* 1997;272:14037-14040.
 18. Shumaker H, Amlal H, Frizzell R, Ulrich CD, Soleimani M. CFTR drives $\text{Na}^+-\text{NHCO}_3^-$ cotransport in pancreatic duct cells: a basis for defective HCO_3^- secretion in CF. *Am J Physiol Cell Physiol* 1999;276:C16-C25.
 19. Ji HL, Chalfant ML, Jovo B, Lockhart JP, Parker SB, Fuller CM, Stanton BA, Benos DJ. The cytosolic termini of the β - and γ -ENaC subunits are involved in the functional interactions between cystic fibrosis transmembrane conductance regulator and epithelial sodium channel. *J Biol Chem* 2000;275:27947-27956.
 20. Jiang QS, Li J, Dubroff R, Ahn YJ, Foskett JK, Engelhardt J, Kleyman TR. Epithelial sodium channels regulate cystic fibrosis transmembrane conductance regulator chloride channels in *Xenopus* oocytes. *J Biol Chem* 2000;275:13266-13274.
 21. Sun F, Hug MJ, Bradbury NA, Frizzell RA. Protein kinase A associates with cystic fibrosis transmembrane conductance regulator via an interaction with ezrin. *J. Biol. Chem.* 2000;275:14360-14366.
 22. Cheung KH, Leung CT, Leung GP, Wong PY. Synergistic effects of cystic fibrosis transmembrane conductance regulator and aquaporin-9 in the rat epididymis. *Biol Reprod* 2003;68:1505-1510.
 23. Ganeshan R, Di A, Nelson DJ, Quick MW, Kirk KL. The interaction between syntaxin 1A and cystic fibrosis transmembrane conductance regulator Cl⁻ channels is mechanistically distinct from syntaxin 1A-SNARE interactions. *J Biol Chem* 2003;278:2876-2885.
 24. Ko SBH, Zeng W, Dorwart MR, Lou X, Kim KH, Millen L, Goto H, Naruse S, Soyombo A, Thomas PJ, Muallem S. Gating of CFTR by the STAS domain of SLC26 transporters. *Nature Cell Biol* 2004;6:343-350.
 25. Li, C., Roy, K., Dandridge, K. & Naren, A. P. Molecular assembly of cystic fibrosis transmembrane conductance regulator in plasma membrane. *J. Biol. Chem.* 2004;279:24673-24684.
 26. Yoo D, Flagg TP, Olsen O, Raghuram V, Foskett JK, Welling PA. Assembly and trafficking of a multiprotein ROMK (Kir 1.1) channel complex by PDZ interactions. *J Biol Chem* 2004;279:6863-6873.
 27. Guggino WB. The cystic fibrosis transmembrane regulator forms macromolecular complexes with PDZ domain scaffold proteins. *Proc Am Thor Soc* 2004;1:28-32.
 28. Riordan JR. Assembly of functional CFTR chloride channels. *Annu Rev Physiol* 2005;67:701-718.
 29. Quinton PM. Chloride impermeability in cystic fibrosis. *Nature*. 1983;301:421-422.
 30. Riordan JR, Rommens JM, Kerem BS, Alon N, Rozmahel R, Grzelczak Z, Zielenski J, Lok S, Plavsic N, Chou JL, Drumm ML, Iannuzzi MC, Collins FC, Tsui L-C. Identification of the cystic fibrosis gene: cloning and characterization of complementary DNA. *Science* 1989;245:1066-1073.
 31. Gustafsson JK, Ermund A, Ambrot D, Johansson MEV, Nilsson HE, Thorell K, Hebert H, Sjövall H, Hansson GC. Bicarbonate and functional CFTR channel are required for proper mucin secretion and link cystic fibrosis with its mucus phenotype. *ScJ Exp Med* 2012;209:1263-1272.
 32. Knowles MR, Boucher RC. Mucus clearance as a primary innate defense mechanism for mammalian

- airways. *J Clin Invest* 2002;109:571–577.
33. Hovenberg HW, Davies JR, Carlstedt I. Different mucins are produced by the surface epithelium and the submucosa in human trachea: identification of MUC5AC as a major mucin from the goblet cells. *Biochem J* 1996;318:319–324.
 34. Wickstrom C, Davies JR, Eriksen GV, Veerman EC, Carlstedt I. MUC5B is a major gel-forming, oligomeric mucin from human salivary gland, respiratory tract and endocervix: identification of glycoforms and C-terminal cleavage. *Biochem J* 1998;334:685–93.
 35. Finkelstein A, Zimmerberg J, Cohen FS. Osmotic swelling of vesicles: its role in the fusion of vesicles with planar phospholipid bilayer membranes and its possible role in exocytosis. *Annu Rev Physiol* 1986;48:163-174.
 36. Holz RW. The role of osmotic forces in exocytosis from adrenal chromaffin cells. *Annu Rev Physiol* 1986;48:175-189.
 37. Fernandez JM, Villalon M, Verdugo P. Reversible condensation of mast cell secretory products in vitro. *Biophys J* 1991;59:1022-1027.
 38. Monck JR, Oberhauser AF, Alvarez de Toledo G, Fernandez JM. Is swelling of the secretory granule matrix the force that dilates the exocytotic fusion pore? *Biophys J* 1991;59:39-47.
 39. Jena BP, Schneider SW, Geibel JP, Webster P, Oberleithner H, Sritharan KC. Gi regulation of secretory vesicle swelling examined by atomic force microscopy. *Proc Natl Acad Sci USA* 1997;94:13317-13322.
 40. Cho S-J, Sattar AKM, Jeong E-H, Satchi M, Cho J, Dash S, Mayes MS, Stromer MH, Jena BP. Aquaporin 1 regulates GTP-induced rapid gating of water in secretory vesicles. *Proc Natl Acad Sci USA* 2002;99:4720-4724.
 41. Kelly M, Cho W-J, Jeremic A, Abu-Hamdah R, Jena BP. Vesicle swelling regulates content expulsion during secretion. *Cell Biol Int* 2004;28:709-716.
 42. Kelly M, Abu-Hamdah R, Cho S-J, Ilie AL, Jena BP. Patch clamped single pancreatic zymogen granules: direct measurement of ion channel activities at the granule membrane. *Pancreatol* 2005;5:443-449.
 43. Lee J-S, Cho W-J, Shin L, Jena BP. Involvement of cholesterol in synaptic vesicle swelling. *Exp Biol Med* 2010;235:470-477.
 44. Shin L, Basi N, Lee J-S, Cho W-J, Chen Z, Abu-Hamdah R, Oupicky D, Jena BP. Involvement of vH⁺-ATPase in synaptic vesicle swelling. *J Neurosci Res* 2010;88:95-101.
 45. Chen Z-H, Lee, J-S, Shin L, Cho W-J, Jena BP. Involvement of β -adrenergic receptor in synaptic vesicle swelling and implication in neurotransmitter release. *J Cell Mol Med* 2011;15:572-576.
 46. Cho S-J, Wakade A, Pappas GD, Jena BP. New structure involved in transient membrane fusion and exocytosis. *New York Acad Sci Annal* 2002;971:254-256.
 47. Trimble WS, Cowan DW, Scheller RH. VAMP-1: A synaptic vesicle-associated integral membrane protein. *Proc Natl Acad Sci USA* 85: 4538-4542, 1988.
 48. Bennett MK, Calakos N, Scheller RH. Syntaxin: A synaptic protein implicated in docking of synaptic vesicles at presynaptic active zones. *Science* 257: 255-259, 1992.
 49. Oyler GA, Higgins GA, Hart RA, Battenberg E, Billingsley M, Bloom FE, Wilson MC. The identification of a novel synaptosomal-associated protein, SNAP-25, differentially expressed by neuronal subpopulations. *J Cell Biol* 109: 3039-3052, 1989.
 50. Weber T, Zemelman BV, McNew JA, Westerman B, Gmachl M, Parlati F, Söllner TH, Rothman JE. SNAREpins: minimal machinery for membrane fusion. *Cell* 92: 759-772, 1998.
 51. Sutton RB, Fasshauer D, Jahn R, Brunger AT. Crystal structure of a SNARE complex involved in synaptic exocytosis at 2.4 Å resolution. *Nature* 395: 347–353, 1998.
 52. Cho SJ, Kelly M, Rognlien KT, Cho JA, Hörber JKH, Jena BP. SNAREs in opposing bilayers interact in a circular array to form conducting pores. *Biophys J* 83: 5: 2522-2527, 2002.
 53. Jeremic A, Kelly M, Cho JH, Cho SJ, Hörber JKH, Jena BP. Calcium drives fusion of SNARE-apposed bilayers. *Cell Biol Int* 28: 19-31, 2004.
 54. Jeremic A, Cho WJ, Jena BP. Membrane fusion: what may transpire at the atomic level. *J Biol Phys & Chem* 4: 139-142, 2004.
 55. Cho WJ, Jeremic A, Jena BP. Size of supramolecular SNARE complex: membrane-directed self-assembly. *J Am Chem Soc* 127: 10156-10157, 2005.
 56. Jeremic A, Quinn AS, Cho WJ, Taatjes DJ, Jena BP. Energy-dependent disassembly of self-assembled SNARE complex: observation at nanometer resolution using atomic force microscopy. *J Am Chem Soc* 128: 26-27, 2006.
 57. Cook JD, Cho WJ, Stemmler TL, Jena BP. Circular dichroism (CD) spectroscopy of the assembly and disassembly of SNAREs: the proteins involved in membrane fusion in cells. *Chem Phys Lett* 462: 6-9, 2008.
 58. Shin L, Cho WJ, Cook J, Stemmler T, Jena BP. Membrane lipids influence protein complex assembly-disassembly. *J Am Chem Soc* 132: 5596-5597, 2010.
 59. Potoff JJ, Issa Z, Manke CW Jr, Jena BP. Ca²⁺-Dimethylphosphate complex formation: providing insight into Ca²⁺ mediated local dehydration and membrane fusion in cells. *Cell Biol Int* 32: 361-366, 2008.
 60. Cho WJ, Lee JS, Ren G, Zhang L, Shin L, Manke CW, Potoff J, Kotaria N, Zhvania MG, Jena BP. Membrane-directed molecular assembly of the neuronal SNARE complex. *J Cell Mol Med* 15: 31-37, 2011.
 61. Misura KM, Scheller RH, Weis WI. Three-dimensional

- structure of the neuronal-Sec1–syntaxin 1a complex. *Nature* 404: 355–362, 2000.
62. Südhof TC. The synaptic vesicle cycle. *Annu Rev Neurosci* 27: 509–547, 2004.
 63. Portis A, Newton C, Pangborn W, Papahadjopoulos D. Studies on the mechanism of membrane fusion: evidence for an intermembrane Ca^{2+} phospholipid complex, synergism with Mg^{2+} , and inhibition by spectrin. *Biochemistry* 18: 780-790, 1979.
 64. Pobbati AV, Stein A, Fasshauer D. N- to C-terminal SNARE complex assembly promotes rapid membrane fusion. *Science* 313: 673–676, 2006.
 65. Jahn R, Scheller RH. SNAREs—engines for membrane fusion. *Nature Rev Mol Cell Biol* 7: 631–643, 2006.
 66. Shen J, Tareste DC, Paumet F, Rothman JE, Melia TJ. Selective activation of cognate SNAREpins by Sec1/Munc18 proteins. *Cell* 128: 183–195, 2007.
 67. Martens S, Kozlov MM, McMahon HT. How synaptotagmin promotes membrane fusion. *Science* 316: 1205–1208, 2007.
 68. Wickner W, Schekman R. Membrane fusion. *Nature Struct Mol Biol* 15: 658–664, 2008.
 69. Chapman ER. How does synaptotagmin trigger neurotransmitter release? *Annu Rev Biochem* 77: 615–641, 2008.
 70. Hui E, Johnson CP, Yao J, Dunning FM, Chapman ER. Synaptotagmin-mediated bending of the target membrane is a critical step in Ca^{2+} -regulated fusion. *Cell* 138: 709–721, 2009.
 71. Südhof TC, Rothman JE. Membrane fusion: grappling with SNARE and SM proteins. *Science* 323: 474–477, 2009.
 72. Stein A, Weber G, Wahl MC, Jahn R. Helical extension of the neuronal SNARE complex into the membrane. *Nature* 460: 525–528, 2009.
 73. Brunger AT, Weninger K, Bowen M, Chu S. Single-molecule studies of the neuronal SNARE fusion machinery. *Annu Rev Biochem* 78: 903–928, 2009.
 74. Südhof TC, Rizo J. Synaptic vesicle exocytosis. *Cold Spring Harb Perspect Biol* 3: a005637, 2011.
 75. Yao J, Gaffaney JD, Kwon SE, Chapman ER. Doc2 is a Ca^{2+} sensor required for asynchronous neurotransmitter release. *Cell* 147: 666–677, 2011.
 76. Diao J, Ishitsuka Y, Bae WR. Single-molecule FRET study of SNARE-mediated membrane fusion. *Biosci Rep* 31: 457–463, 2011.
 77. Szklarczyk D, Franceschini A, Kuhn M, Simonovic M, Roth A, Minguéz P, Doerks T, Stark M, Mueller J, Bork P, Jensen LJ, von Mering C. The STRING database in 2011: functional interaction networks of proteins, globally integrated and scored. *Nucleic Acid Res* 39: D561-D568, 2011.
 78. Lam AD, Tryoen-Toth P, Tsai B, Vatile N, Stuenkel EL. SNARE-catalyzed fusion events are regulated by syntaxin1A-lipid interactions. *Mol Biol Cell* 19: 485–497, 2008.
 79. Clifford-Nunn B, Showalter HD, Andrews PC. Quaternary diamines as mass spectrometry cleavable crosslinkers for protein interactions. *Journal of the American Society for Mass Spectrometry* 23:201-212, 2012.
 80. Kovari, L.C., Brunzelle, J.S., Lewis, K.T., Cho, W.J., Lee, J-S., Taatjes, D.J., Jena, B.P. (2014) X-ray solution structure of the native neuronal porosome-synaptic vesicle complex: Implication in neurotransmitter release. *Micron* 56:37-43.
 81. Szymborska A, de Marco A, Daigle N, Cordes VC, Briggs JAG, Ellenberg J. Nuclear pore scaffold structure analyzed by super-resolution microscopy and particle averaging. *Science* 341:655-658, 2013.
 82. Gray JJ, Moughon SE, Kortemme T, Schueler-Furman O, Misura KM, Morozov AV, Baker D. Protein-protein docking predictions for the CAPRI experiment. *Proteins* 52:118-122, 2003.
 83. Smith GR, Sternberg MJ. Prediction of protein-protein interactions by docking methods. *Curr Opin Struct Biol* 12:28-35, 2002.
 84. Janin J, Henrick K, Moult J, Eyck LT, Sternberg MJ, Vajda S, Vakser I, Wodak SJ. CAPRI: a Critical Assessment of PRedicted Interactions. *Proteins* 52:2-9, 2003.
 85. Schneidman-Duhovny D, Inbar Y, Polak V, Shatsky M, Halperin I, Benyamini H, Barzilai A, Dror O, Haspel N, Nussinov R, Wolfson HJ. Taking geometry to its edge: fast unbound rigid (and hinge-bent) docking. *Proteins* 52:107-112, 2003.
 86. Katchalski-Katzir E, Shariv I, Eisenstein M, Friesem AA, Aflalo C, Vakser IA. Molecular surface recognition: determination of geometric fit between proteins and their ligands by correlation techniques. *Proc Natl Acad Sci USA* 89:2195-2199, 1992.
 87. Gabb HA, Jackson RM, Sternberg MJ. Modelling protein docking using shape complementarity, electrostatics and biochemical information. *J Mol Biol* 272:106-120, 1997.
 88. Moont G, Sternberg MJ. *Modeling protein-protein and protein-DNA docking*, edited by Lengauer T. Weinheim: Wiley-VCH, 2001.
 89. Jackson RM, Gabb HA, Sternberg MJ. Rapid refinement of protein interfaces incorporating solvation: application to the docking problem. *J Mol Biol* 276:265-285, 1998.
 90. Vakser IA. Protein docking for low-resolution structures. *Protein Eng* 8:371-377, 1995.
 91. Mandell JG, Roberts VA, Pique ME, Kotlovyy V, Mitchell JC, Nelson E, Tsigelny I, Ten Eyck LF. Protein docking using continuum electrostatics and geometric fit. *Protein Eng* 14:105-113, 2001.
 92. Chen R, Li L, Weng Z. ZDOCK: an initial-stage protein-docking algorithm. *Proteins* 52:80-87, 2003.
 93. Ritchie DW, Kemp GJ. Protein docking using spherical polar Fourier correlations. *Proteins* 39:178-194, 2000.
 94. Fernandez-Recio J, Totrov M, Abagyan R. Soft

- protein- protein docking in internal coordinates. *Protein Sci* 11:280-291, 2002.
95. Gray JJ, Moughon S, Wang C, Schueler-Furman O, Kuhlman B, Rohl CA, Baker D. Protein-protein docking with simultaneous optimization of rigid-body displacement and side-chain conformations. *J Mol Biol* 331:281-299, 2003.
96. Gabdoulhine RR, Wade RC. Protein-protein association: investigation of factors influencing association rates by Brownian dynamics simulations. *J Mol Biol* 306:1139-1155, 2001.
97. Fitzjohn PW, Bates PA. Guided docking: first step to locate potential binding sites. *Proteins* 52:28-32, 2003.
98. Aloy P, Russell RB. Interrogating protein interaction networks through structural biology. *Proc Natl Acad Sci USA* 99:5896-5901, 2002.
99. Aloy P, Bottcher B, Ceulemans H, Leutwein C, Mellwig C, Fischer S, Gavin AC, Bork P, Superti-Furga G, Serrano L, Russell RB. Structure-based assembly of protein complexes in yeast. *Science* 303:2026-2029, 2004.
100. Pieper U, Eswar N, Braberg H, Madhusudhan MS, Davis FP, Stuart AC, Mirkovic N, Rossi A, Marti-Renom MA, Fiser A, Webb B, Greenblatt D, Huang CC, Ferrin TE, Sali A. MODBASE, a database of annotated comparative protein structure models, and associated resources. *Nucleic Acids Res* 32: D217-D222, 2004.
101. Marti-Renom MA, Stuart AC, Fiser A, Sanchez R, Melo F, Sali A. Comparative protein structure modeling of genes and genomes. *Annu Rev Biophys Biomol Struct* 29:291-325, 2000.
102. Volkman N, Hanein D, Ouyang G, Trybus KM, DeRosier DJ, Lowey S. Evidence for cleft closure in actomyosin upon ADP release. *Nat Struct Biol* 7:1147-1155, 2000.
103. Roseman AM. Docking structures of domains into maps from cryo-electron microscopy using local correlation. *Acta Crystallogr D Biol Crystallogr* 56:1332-1340, 2000.
104. Wriggers W, Birmanns S. Using situs for flexible and rigid-body fitting of multiresolution single-molecule data. *J Struct Biol* 133:193-202, 2001.
105. Ceulemans H, Russell RB. Fast fitting of atomic structures to lower resolution electron density maps by surface overlap maximization. *J Mol Biol* 338:783-793, 2004.
106. Volkman N, Hanein D. Quantitative fitting of atomic models into observed densities derived by electron microscopy. *J Struct Biol* 125:176-184, 1999.
107. Rossmann MG, Bernal R, Pletnev SV. Combining electron microscopic with x-ray crystallographic structures. *J Struct Biol* 136:190-200, 2001.
108. Chiu W, Baker ML, Jiang W, Zhou ZH. Deriving folds of macromolecular complexes through electron cryomicroscopy and bioinformatics approaches. *Curr Opin Struct Biol* 12:263-269, 2002.

DISCOVERIES is a peer-reviewed, open access, online, multidisciplinary and integrative journal, publishing high impact and innovative manuscripts from all areas related to MEDICINE, BIOLOGY and CHEMISTRY; © 2014, Applied Systems