

Supplementary Material

Testosterone Levels and Type 2 Diabetes– No Correlation with Age, Differential Predictive Value in Men and Women

Running Title: Testosterone levels and Type 2 Diabetes

Mahir Karakas^{1,2}, MD; Sarina Schäfer^{1,2}, MD; Sebastian Appelbaum¹, MSc; Francisco Ojeda¹, PhD; Kari Kuulasmaa³, PhD; Burkhard-Ekkehart Brueckmann¹; Filip Berisha¹, MD; Benedict Schulte-Steinberg¹, MD; Pekka Jousilahti³, MD; Stefan Blankenberg^{1,2}, MD; Tarja Palosaari³, MSc; Veikko Salomaa³, MD and Tanja Zeller^{1,2*}, PhD

¹Department of General and Interventional Cardiology, University Heart Center, 20246 Hamburg, Germany, m.karakas@uke.de, sar.schaefer@uke.de, sebastian.appelbaum@tu-dortmund.de, f.ojeda-echevarria@uke.de, Burkhard.Brueckmann@uk-erlangen.de, f.berisha@uke.de, b.schulte-steinberg@uke.de, s.blankenberg@uke.de, t.zeller@uke.de

²German Center for Cardiovascular Research (DZHK), Partner Site Hamburg, Lübeck, Kiel, Hamburg, 20246 Hamburg Germany, m.karakas@uke.de, sar.schaefer@uke.de, s.blankenberg@uke.de, t.zeller@uke.de

³National Institute for Health and Welfare, 00271Helsinki, Finland, kari.kuulasmaa@thl.fi, pekka.jousilahti@thl.fi, tarja.palosaari@thl.fi, veikko.salomaa@thl.fi

Table S1.

Distribution of baseline testosterone levels according to sex [nmol/L]

	Quartile 1	Quartile 2	Quartile 3	Quartile 4
Men, IQR	0; 12.91	12.91; 17.12	17.12; 22.02	22.02; 35.0
Women, IQR	0; 0.87	0.87; 1.15	1.15; 1.56	1.56; 35.0

Values are shown as normalized median values; IQR = interquartile range

Figure S1

Distribution of baseline testosterone levels according to sex. A and B are shown for absolute testosterone levels, C and D are shown for log transformed testosterone levels.

