

Introduction

Aromatic hydrocarbons such as benzene, toluene, and ethylbenzene emitted by solvent volatilization, vehicle exhaust, and other anthropogenic sources are usually organic pollutants in the atmosphere [1, 2]. They are inherently toxic and harmful to human health [3, 4]. Some aromatic compounds discharged into the atmosphere exist in the gas phase and are mainly oxidized by OH radicals to form semi-volatile and non-volatile organics. These organics then form secondary organic aerosol (SOA) by self-condensation or gas-particle partitioning [5, 6]. The other part of aromatic compounds dissolves in liquid aerosol, water droplets, clouds, and mist, then undergoes photooxidation to generate phenols, organic acids, and other low volatile products. These products remain in the particle phase after water evaporation, leading to the formation of SOA particles [7, 8]. SOA can scatter and absorb sunlight, reduce visibility, and have become a driving factor that interferes with regional climate [9, 10]. The contribution of SOA generated by aqueous reactions to atmospheric SOA is comparable to that of SOA formed from gaseous reactions and can explain the results of field observations that cannot be interpreted by gaseous formation methods [11, 12]. With the increasing impact of climate change, the study on the formation and optics of aqueous SOA has turned into a hot topic in atmospheric chemistry.

With the acceleration of urbanization, a large amount of heavy metal dust and fine particles have been emitted from human sources such as industry and transportation [13]. The heavy metal components in these fine particles are converted into soluble metal ions via atmospheric processes [14]. The field measurement results indicate that Cu, Fe, Ni, etc. are usual heavy metal ions in atmospheric aerosols [15]. Heavy metal ions have a catalytic effect, and OH, HO₂ radicals, and other reactive oxygen species (ROS) can be produced by Fenton-like chemical catalysis of H₂O₂ and O₃, affecting the oxidation potential of the atmosphere [16-18]. ROS participate in the oxidation of atmospheric VOCs and the aging process of aerosol particles, thereby impacting the components and optics of aqueous SOA.

Experiments conducted by Nguyen et al. [19] confirmed that when Fe²⁺ ions were present in the aqueous reaction system of glycolaldehyde and hydrogen peroxide, glycolic acid, oxalic acid, and other carboxylic acids significantly increased, and the oxygen-carbon ratio (O/C) for the formed aqueous SOA was as high as 0.90. Kameel et al. [20] found that Fe²⁺ ions can catalyze the aqueous reaction of isoprene with hydrogen peroxide to generate polyols, aldehydes, and carboxylic acids, thereby remarkably promoting the formation of aqueous SOA. Zhang et al. [21] demonstrated through field measurement experiments using an aerosol mass spectrometer that Fe³⁺ ions containing aerosol particles can promote the production of oxalate, glyoxylate, and other oxidized organic compounds. While Nimer et al. [22], Ling et al. [23], and Al-Abadleh et al. [24]

found that Fe³⁺ ions could promote methoxyphenol and catechol to form secondary brown carbon products via complexation, polymerization, and other aqueous reactions. However, the above experiments [19-24] only focus on the influences of heavy metal ions on the chemical components of aqueous SOA. The optical properties of aqueous SOA in presence of heavy metal ions need to be further studied.

Toluene is the most abundant aromatic compound in the atmosphere. The chemical composition and reaction mechanism for OH-initiated photooxidation of toluene to form SOA particles in atmospheric photochemical reactions have been studied detailedly using a smog chamber [6, 25-28]. The Henry constant of toluene in the aqueous phase at room temperature is 1.4×10⁻³ mol•m⁻³•Pa⁻¹ [29], which has certain water solubility and can be dissolved in rainwater, liquid aerosol, and other aqueous phases [30]. Although Heath et al. [31] have carried out a study of the effects of temperature, pH, and other factors on the aqueous reaction rate of benzene and OH radicals and the yield of phenol, the effects of heavy metal ions on the aqueous reaction products of aromatic compounds still need to be further studied. Copper is a common heavy metal element, mainly derived from garbage incineration and biomass combustion. Copper mainly exists in the form of Cu²⁺ ions in aqueous phases [13, 15], nevertheless, the influences of Cu²⁺ ions on the formation and composition of aqueous SOA are few reported.

Our group used TSI 9302 to atomize the mixed solution of aldehydes and inorganic ammonium salts and dried it via a silica gel diffusion tube to generate SOA particles. Components of aqueous SOA are on-line and off-line, detected using mass spectrometry and spectroscopy; 4-methyl-imidazole and other imidazoles are identified [32]. On this basis, on-line and off-line apparatus is used to measure constituents of SOA generated by aqueous photooxidation of toluene in presence of Cu²⁺ ions in the current study. The averaged mass absorption coefficient (<MAC>) of aqueous SOA in 200-600 nm is measured by solvent extraction-continuous spectroscopy, as proposed by Updyke et al. [33] and Powelson et al. [34], and the influences of Cu²⁺ ions on the composition and optics of aqueous SOA are investigated. These provide the basis for studying the components and optics of aqueous SOA in presence of heavy metal ions.

Experimental Procedures

Materials

Toluene (> 99%), copper chloride (> 99%), hydrogen peroxide (30%), and hydrochloric acid (36%~38%) are all bought from China National Pharmaceutical Group Chemical Reagent Co., Ltd.




Fig. 4. Averaged positive ion mass spectra for 200 SOA particles generated by aqueous photooxidation of toluene in presence of 4 $\mu\text{mol/L}$ copper ions.




Fig. 5. UV-Vis absorption spectra of collection solutions for SOA particles generated by aqueous photooxidation of toluene in presence of different concentration of copper ions.

In addition, oxygen molecules can also be added to the OH-toluene adduct to form peroxy radicals, which undergo isomerization to generate oxygen bridge radicals and produce 2-butendial and methylglyoxal through ring fracture reactions. Therefore, cresol, aromatic ether, and aldehyde compounds are the products of aqueous photooxidation of toluene initiated by OH radicals. These products have low volatility and remain in the particle phase to form SOA after water evaporation [7, 8].

The collection solution of aqueous SOA particles is not separated by a chromatographic column and directly enters a mass spectrometer to measure the mass spectra after electrospray ionization. Electrospray ionization is a soft ionization technology that will not break the molecular ion peak by using a low-voltage method. The negative ionization mode causes the measured organic molecule to dissociate and lose hydrogen ions, resulting in a deprotonated molecular ion peak ($[\text{M}-\text{H}]^{-}$), providing information on the molecular weight of




Fig. 6. Possible reaction mechanism for formation of phenolic and aldehyde compounds by aqueous photooxidation of toluene initiated by OH radical.




Fig. 7. Electrospray ionization negative ion mass spectra of SOA particles generated by aqueous photooxidation of toluene in absence of copper ions.

the measured component [35]. Electrospray negative ion mass spectra for the collection solution of aqueous SOA in absence of Cu^{2+} ions shown in Fig. 7 emerge $[\text{M-H}]^-$ peaks at $m/z = 71, 83, 91, 107,$ and 213 . Among them, the peak of $m/z = 91$ may refer to the unreacted deprotonated molecular ion peak of toluene ($\text{C}_6\text{H}_5\text{CH}_2^-$). In view of the possible reaction mechanism between toluene and OH radicals displayed in Fig. 6 and the molecular weight information of the products, the strongest mass peak at $m/z = 107$ is considered the deprotonated molecular ion peak of cresol ($\text{CH}_3\text{C}_6\text{H}_4\text{O}^-$). While peaks at $m/z = 71,$ $m/z = 83,$ and $m/z = 213$ may refer to deprotonated molecular ion peaks of methylglyoxal (CH_3COCO^-), 2-butenal (HCOCH=CHCO^-), and 2,3'-dimethyl-4'-hydroxy-diphenyl ether ($\text{C}_{14}\text{H}_{13}\text{O}_2^-$). These further verify the existence of cresol, aromatic ether, and aldehyde compounds in SOA particles generated by the aqueous photooxidation of toluene in absence of Cu^{2+} ions.

When there are Cu^{2+} ions of different concentrations in an aqueous system, as shown in Fig. 5, the absorption curve of each collection solution of aqueous SOA particles does not change significantly, but the absorbance of each collection solution at 277 nm is greater than that of a solution without Cu^{2+} ions. As can be seen from Fig. 5, when the concentration of Cu^{2+} is in the range of 0.25 - 4 $\mu\text{mol/L}$, the absorbance of collection solution at 277 nm gradually increases with an increment of Cu^{2+} ions. In absence of Cu^{2+} ions, the absorbance of collection solution for aqueous SOA particles at 277 nm is 0.254, while in presence of 4 $\mu\text{mol/L}$ Cu^{2+} ions, the absorbance of the collection solution at 277 nm is 1.489, about 6 times that of the absence of Cu^{2+} ions. These show that the content of phenolic compounds in aqueous SOA particles increases significantly when Cu^{2+} is present in the aqueous system. This is mainly because Cu^{2+} ions belong to fourth-period transition-state metal ions and have catalytic effects.




Fig. 8. Possible reaction mechanism for formation of polyhydroxyphenols and hydroxyphenyl ether polymers by aqueous photooxidation of toluene initiated by OH radical.

They form Fenton-like reagents with hydrogen peroxide, and the Cu element catalyzes hydrogen peroxide to produce OH radicals through the conversion of Cu^{2+} and Cu^+ valence states [44]. Therefore, when Cu^{2+} ions are present in an aqueous system, hydrogen peroxide can generate more OH radicals under ultraviolet irradiation and Cu^{2+} ion catalysis. As illustrated in Fig. 8, the OH radical undergoes an addition reaction with toluene, and the OH radical is added to the benzene ring to form cresol. The excessive OH radicals then continue to react with cresol to produce polyhydroxyphenol products such as dihydroxybenzene and trihydroxybenzene [45]. The resulting aqueous SOA contains more phenolic compounds and the absorbance of the collection solution at 277 nm increases significantly. Due to the constant concentration of hydrogen peroxide and toluene in each experiment, when the concentration of Cu^{2+} ions increases to a certain concentration (4 $\mu\text{mol/L}$), hydrogen peroxide is fully catalyzed to produce OH radicals, and the absorbance of the collection solution at 277 nm reaches its maximum value. Afterward, when the concentration of Cu^{2+} ions continues to increase, the amount of OH radicals formed almost stays constant, so the absorbance of the collection solution at 277 nm basically remains unchanged.

In order to characterize the chemical components of SOA particles generated by aqueous photooxidation of toluene in presence of 4 $\mu\text{mol/L}$ Cu^{2+} ions in detail, off-line electrospray ionization mass spectrometry is carried out, and the negative ion mass spectra measured are shown in Fig. 9. Apart from peaks of $m/z = 71$ (methylglyoxal deprotonation molecular ion (CH_3COCO^-) peak), $m/z = 83$ (2-butenedial deprotonation molecular ion ($\text{HCOCH}=\text{CHCO}^-$) peak), $m/z = 107$

(cresol deprotonation molecular ion ($\text{CH}_3\text{C}_6\text{H}_4\text{O}^-$) peak), and $m/z = 213$ (2,3'- dimethyl-4'-hydroxy diphenyl ether deprotonation molecular ion ($\text{C}_{14}\text{H}_{13}\text{O}_2^-$) peak), there are strong peaks of $m/z = 123$, 139, 320, 427, 534, and 641. According to the reaction mechanism shown in Fig. 9 and the molecular weight information of the products, $m/z = 123$ and 139 are considered deprotonated molecular ion peaks for methyl dihydroxybenzene ($\text{CH}_3\text{C}_6\text{H}_4\text{O}_2^-$) and methyl trihydroxybenzene ($\text{CH}_3\text{C}_6\text{H}_4\text{O}_3^-$). Also, the mass charge ratio of $m/z = 213$, 320, 427, 534, and 641 differs by 107, which may refer to deprotonated molecular ion peaks of hydroxy-phenyl ether polymers. On the basis of the experimental results of Sun et al. [43], as displayed in Fig. 8, OH radicals can react with cresol to generate cresol oxygen radicals, which nucleophilically attacks cresol to produce hydroxyphenyl ether dimer (2,3'-dimethyl-4'- hydroxy-diphenyl ether), which generates deprotonated molecular ions ($\text{C}_{14}\text{H}_{13}\text{O}_2^-$, $m/z = 213$) by electrospray ionization. Similarly, under the activation of OH radicals, hydroxyphenyl ether dimers continue to react with cresol to produce hydroxyphenyl ether trimers (producing deprotonated molecular ion $\text{C}_{21}\text{H}_{20}\text{O}_3^-$, $m/z = 320$), tetramers (producing deprotonated molecular ion $\text{C}_{28}\text{H}_{27}\text{O}_4^-$, $m/z = 427$), pentamers (producing deprotonated molecular ion $\text{C}_{35}\text{H}_{34}\text{O}_5^-$, $m/z = 534$), and hexamers (producing deprotonated molecular ion $\text{C}_{42}\text{H}_{41}\text{O}_6^-$, $m/z = 641$). These further confirm that Cu^{2+} ions catalyze the production of more OH radicals from hydrogen peroxide, leading to an increase in the generation of methyl dihydroxybenzene and methyl trihydroxybenzene and triggering the polymerization of cresol to form high molecular weight products of hydroxyphenyl ether polymers.




Fig. 9. Electrospray ionization negative ion mass spectra of SOA particles generated by aqueous photooxidation of toluene in presence of 4 $\mu\text{mol/L}$ copper ions.




Fig. 10. Averaged mass absorption coefficient (<MAC>) of SOA particles generated by aqueous photooxidation of toluene in presence of different concentration of copper ions.

Optical Characterization of Aqueous SOA Particles

Aerosol particles affect climate change via light absorption, and the averaged mass absorption coefficient (<MAC>) is often used to characterize the impact of particles on light absorption [33, 34]. The total organic carbon concentration of the collection solution for SOA generated by aqueous photooxidation of toluene in presence of different concentrations of Cu^{2+} ions is measured using a TOC-L analyzer. Combined with the

UV-Vis absorption spectra for the collection solution at 200-600 nm shown in Fig. 5, <MAC> is obtained according to equations (1) and (2). As shown in Fig. 10, when Cu^{2+} ions are not present, the <MAC> of SOA generated by aqueous photooxidation of toluene in 200-600 nm is $38 \text{ cm}^2/\text{g}$, which is a bit higher than that of 1,3,5-trimethylbenzene SOA formed in absence of NO_x , measured by Updyke et al. [33] ($20 \text{ cm}^2/\text{g}$). The <MAC> of aqueous SOA particles obviously increases when Cu^{2+} ions are present in the system and gradually

increases with an increment of Cu^{2+} ions. However, when the concentration of Cu^{2+} ions exceeds $4 \mu\text{mol/L}$, the optical properties of aqueous SOA particles tend to be stable.

Phenolic compounds have a $\text{C}=\text{C}$ bond and electron-removing functional group of $-\text{OH}$ in the benzene ring, which can absorb photons and have a certain degree of light absorption ability. The greater the number of $-\text{OH}$ functional groups contained in a phenolic compound molecule, the stronger its light absorption ability [46]. According to the above chemical component detection results, the major constituents of SOA generated by aqueous photooxidation of toluene in absence of Cu^{2+} ions are cresol, aromatic ether, and aldehyde compounds. These product molecules only have $\text{C}=\text{C}$ and $\text{C}=\text{O}$ bonds, without strong chromophores or color aids. They contain a small number of $-\text{OH}$ functional groups. Thus, aqueous SOA particles have weak absorption abilities [46]. When Cu^{2+} ions are present in an aqueous system, Cu^{2+} ions catalyze the production of OH radicals from hydrogen peroxide. As the concentration of Cu^{2+} ions increases, the catalytic ability increases, and the number of OH radicals formed in the system increases, resulting in an increase in the formation of methyl dihydroxybenzene, methyl trihydroxybenzene, and hydroxyphenyl ether polymers formed by the polymerization of cresol. These products contain a large number of $-\text{OH}$ functional groups, thus enhancing their ability to absorb light, resulting in an increment of $\langle \text{MAC} \rangle$ for the generated aqueous SOA particles. Due to the constant concentration of toluene and hydrogen peroxide in each experiment, when Cu^{2+} ions rise to a certain concentration ($4 \mu\text{mol/L}$), they catalyze the majority of hydrogen peroxide to produce OH radicals, and the $\langle \text{MAC} \rangle$ of formed aqueous SOA particles reaches a maximum value of $306 \text{ cm}^2/\text{g}$, which is a little higher than that of ammonia-aged aromatic SOA reported by Updyke et al. [33] ($\sim 300 \text{ cm}^2/\text{g}$). Since then, the concentration of Cu^{2+} ions has continued to rise, but OH radicals no longer increase. The formed polyhydroxyphenols and hydroxyphenyl ether polymers remained basically unchanged, and the optics of aqueous SOA particles tended to stabilize.

It is noteworthy that some researchers have recently carried out studies on the formation and aging of toluene and other aromatic SOA [47-50]. Mitra et al. [47] have investigated the effects of water vapors, active nitrogen, and acidity on the light-absorbing components and optics of toluene SOA and found that the mass absorption cross-section (MAC) of toluene SOA increased with $[\text{NO}_x/\Delta\text{HCl}]$. Also, under the condition of 80% RH, acidity facilitated the production of brown carbon components, while the presence of ammonia led to an increase in MAC for toluene SOA. Wang et al. [48] have studied the influences of ferric chloride fine particles on the components and optics of toluene SOA. They observed that methylcatechol and other phenolic compounds produced by the photooxidation of toluene will condense on surfaces for ferric chloride

fine particles and then combine with iron ions to form metallo-organic complexes with strong light absorption ability. As a result, the yield of toluene SOA increased, and the light absorption ability of SOA (MAC) was enhanced significantly. Campbell et al. [49] have coated the surface of CuSO_4 and FeSO_4 seed particles with β -pinene and naphthalene SOA particles and used two novel online instruments to quantify the OH radical and reactive oxygen species produced. The effects of metal-organic interaction and particle-phase chemistry on the oxidative potential (OP) of SOA were studied. They found a range of antagonistic and synergistic interactions for Fe(II), Cu(II) and SOA, highlighting metal-ascorbate and naphthoquinone-ascorbate reactions as vital drivers of OP. Malecha and Nizkorodov [50] have collected 135-trimethylbenzene, isoprene, and other anthropogenic and biogenic SOA particles in uncoated CaF_2 windows as substrates, annealed and aged by direct photolysis by UV-LED. Oxygenated volatile organic compounds such as acetaldehyde, acetone, formic acid, and acetic acid were detected during the photodegradation of SOA particles. They estimate that under summer conditions in Los Angeles, SOA would lose at least 1% of its mass during photolysis within a 24-hour period. The photodegradation process of SOA would affect the budget of gaseous organic compounds in the atmosphere.

However, the above experiments focus on the study of SOA produced by gaseous photooxidation of toluene and other volatile organic compounds. Compared with these studies [47-50], based on the solubility of toluene in liquid aerosol, rainwater, and other aqueous phases, the device shown in Fig. 1 is used to carry out the aqueous photooxidation reaction of toluene in presence of Cu^{2+} ions in this study. The reaction product solution was atomized by TSI 9302, and atmospheric aqueous SOA particles were simulated to form after water evaporation. The composition and optics of the formed aqueous SOA are on-line and off-line, characterized by mass spectrometry and spectroscopy. ALTOFMS on-line measurement combined with UV-Vis absorption spectroscopy and electrospray ionization mass spectrometry are used for off-line detection to verify that the main components of SOA particles generated by aqueous photooxidation of toluene in presence of Cu^{2+} ions are polyhydroxyphenols and hydroxyphenyl ether polymers. These experimental results are consistent with those of Chang et al. [45]. Their results showed that the infrared spectra for aqueous reaction products of phenol and other phenolic compounds initiated by hydroxyl radicals are similar to those of aerosol light-absorbing substances of humic-like substances (HULIS) measured in the field. The hydroxyphenyl ether polymer formed from the aqueous reaction of phenolic compounds is one of the major sources of HULIS in aerosol particles. Especially in industrial areas heavily polluted by automobile exhaust, there are high concentrations of aromatic compounds and heavy metal fine particles in the atmosphere. Under conditions of

- R.M., SAVARY V., SETYAN A., WENGER J.C. Scanning electron microscopy-energy dispersive X-ray spectrometry (SEM-EDX) and aerosol time-of-flight mass spectrometry (ATOFMS) single particle analysis of metallurgy plant emissions. *Environmental Pollution*. **210**, 9, **2016**.
43. SUN Y.L., ZHANG Q., ANASTASIO C., SUN J. Insights into secondary organic aerosol formed via aqueous-phase reactions of phenolic compounds based on high resolution mass spectrometry. *Atmospheric Chemistry and Physics*. **10** (2), 4809, **2010**.
 44. SINGH D.K., GUPTA T. Role of ammonium ion and transition metals in the formation of secondary organic aerosol and metallo-organic complex within fog processed ambient deliquescent submicron particles collected in central part of Indo-Gangetic Plain. *Chemosphere*. **181**, 725, **2017**.
 45. CHANG J.L., THOMPSON J.E. Characterization of colored products formed during irradiation of aqueous solutions containing H₂O₂ and phenolic compounds. *Atmospheric Environment*. **44** (4), 541, **2010**.
 46. LASKIN A., LASKIN J., NIZKORODOV S.A. Chemistry of atmospheric brown carbon. *Chemical Reviews*. **115** (10), 4335, **2015**.
 47. MITRA K., MISHRA H.R., PEI X.Y., PATHAK R.K. Secondary organic aerosol (SOA) from photo-oxidation of toluene: 1 influence of reactive nitrogen, acidity and water vapours on optical properties. *Atmosphere*. **13** (7), 1099, **2022**.
 48. WANG W.C., HUANG M.Q., HU H.M., ZHAO W.X., HU C.J., GU X.J., ZHANG W.J. Characterization of chemical components and optical properties of toluene secondary organic aerosol in presence of ferric chloride fine particles. *Atmosphere*. **14** (7), 1075, **2023**.
 49. CAMPBELL S.J., UTINGER B., BARTH A., PAULSON S.E., KALBERER M. Iron and copper alter the oxidative potential of secondary organic aerosol: insights from online measurements and model development. *Environmental Science & Technology*. **57** (36), 13546, **2023**.
 50. MALECHA K.T., NIZKORODOV S.A. Photodegradation of secondary organic aerosol particles as a source of small, oxygenated volatile organic compounds. *Environmental Science & Technology*. **50** (18), 9990, **2016**.