

## Supplementary Material

### **Concentrations of the endocannabinoid *N*-arachidonylethanolamine in the follicular fluid of women with endometriosis: the role of M1 polarised macrophages**

*Bruno M. Fonseca<sup>A,D</sup>, Beatriz Moreira-Pinto<sup>A</sup>, Lia Costa<sup>A,B</sup>, Eduarda Felgueira<sup>B</sup>, Pedro Oliveira<sup>C</sup> and Irene Rebelo<sup>A</sup>*

<sup>A</sup>UCIBIO, REQUIMTE, Laboratorio de Bioquímica, Departamento de Ciências Biológicas, Faculdade de Farmácia da Universidade do Porto, Rua de Jorge Viterbo Ferreira no. 228 4050-313 Porto, Portugal.

<sup>B</sup>Unidade de Medicina da Reprodução Dra. Ingeborg Chaves, Centro Hospitalar de Vila Nova de Gaia, Espinho, Portugal.

<sup>C</sup>Unidade de Investigação em Epidemiologia (EPIUnit), Departamento de Estudo de Populações, Instituto de Ciências Biomédicas Abel Salazar, Universidade do Porto, Porto, Portugal.

<sup>D</sup>Corresponding author. Email: [brunofonseca@ff.up.pt](mailto:brunofonseca@ff.up.pt)

**Table S1.** Primer sequences for FAAH, NAPE-PLD and GAPDH genes. Detailed description of genes tested, primer pairs' sequences and qRT-PCR conditions.

**Table S1.** Primer sequences for FAAH, NAPE-PLD and GAPDH genes. Detailed description of genes tested, primer pairs' sequences and qRT-PCR conditions.

Symbol	Accession Number	Description	Primers (5'-3')	Amplification conditions <sup>1)</sup>	T <sub>m</sub> (°C)	Reference
FAAH	NM_001441.2	Homo sapiens fatty acid amide hydrolase	S: GGCCGTCAGCTACACTATGC AS: ATCAGTCGCTCCACCTCCC	95°C, 03 sec 59°C, 30 sec 72°C, 90 sec	87	[14]
NAPE-PLD	NM_001122838.1	Homo sapiens <i>N</i> -acyl phosphatidylethanolamine phospholipase D	S: AGATGGCTGATAATGAGAA AS: TTCTCCTCCCACCAGTC	95°C, 03 sec 58°C, 30 sec 72°C, 90 sec	83.5	-
GAPDH	NC_000012.12	Homo sapiens glyceraldehyde-3-phosphate dehydrogenase	S: AGAACATCATCCCTGCCTC AS: GCCAAATTCGTTGTCATACC	95°C, 03 sec 55°C, 30 sec 72°C, 90 sec	78	[14]

<sup>1)</sup>Cycling conditions were as follows: an initial step at 95°C 3 min for enzyme activation in all cases, followed by up to 40 cycles of denaturation, a annealing and primer extension as described. \*The primers for Na pe-pld were designed in-house using Primer Premier (version 5.0 software; Premier Biosoft International, Palo Alto, CA; www.premierbiosoft.com).