



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

In situ measurements of the Majorite-Akimotoite-Perovskite phase transition boundaries in MgSiO₃

Citation for published version:

Hirose, K, Komabayashi, T, Murakami, M & Funakoshi, K-I 2001, 'In situ measurements of the Majorite-Akimotoite-Perovskite phase transition boundaries in MgSiO₃', *Geophysical Research Letters*, vol. 28, no. 23, pp. 4351-4354. <https://doi.org/10.1029/2001GL013549>

Digital Object Identifier (DOI):

[10.1029/2001GL013549](https://doi.org/10.1029/2001GL013549)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Geophysical Research Letters

Publisher Rights Statement:

Published in Geophysical Research Letters by the American Geophysical Union (2001)

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



In Situ Measurements of the Majorite-Akimotoite-Perovskite Phase Transition Boundaries in MgSiO_3

Kei Hirose, Tetsuya Komabayashi, and Motohiko Murakami

Department of Earth and Planetary Sciences, Tokyo Institute of Technology, Tokyo, Japan

Ken-ichi Funakoshi

Japan Synchrotron Radiation Research Institute, Mikazuki-cho, Hyogo, Japan

Abstract. Here we report the phase boundaries between majorite, akimotoite (ilmenite), and perovskite in MgSiO_3 determined by in situ measurements in a multi-anvil apparatus. We used both gold and platinum as internal pressure standards at high pressure and temperature. Our results obtained at 1400-2000°C, together with previous results obtained at 1000-1200°C by *Ono et al.* [2001], precisely locate the akimotoite-perovskite transition boundary at P (GPa) = $25.09 - 0.0027 \times T$ (°C), based on the P - V - T equation of state of gold [*Anderson et al.*, 1989]. Our new experimental data show the position of triple point to be 20.0 GPa and 1920°C. The present measurements reconfirm our earlier study [*Hirose et al.*, 2001] that Anderson's gold pressure scale gives slightly higher pressures than the platinum pressure scale proposed either by *Jamieson et al.* [1982] or *Holmes et al.* [1989].

1. Introduction

MgSiO_3 -rich perovskite is the predominant mineral in the Earth's lower mantle. Attempts have been made repeatedly to determine its stability at high pressure and temperature, including recent in situ synchrotron X-ray diffraction measurements in a multianvil apparatus [*Kato et al.*, 1995; *Kuroda et al.*, 2000; *Ono et al.*, 2001]. The phase transition between akimotoite and perovskite was detected by observing the change in diffraction pattern during either a temperature or a pressure cycle in these previous studies. This method works well if the transition occurs relatively fast. However, a precise determination of the phase boundary was hindered in these studies by (1) the loss of diffraction peaks of akimotoite due to rapid grain growth at high temperature and (2) the slow phase transition between the high-pressure phases by a kinetic effect. In the present in situ measurement, therefore, the reaction kinetics is a concern.

We determined the majorite - akimotoite - perovskite phase transition boundaries in MgSiO_3 in a wide temperature range from 1400 up to 2000°C. To avoid kinetic effects, we collected only one data point per run at a P - T condition of interest, like a conventional quench experiment. Both gold and platinum were used as internal pressure standards,

and their consistency was examined by a simultaneous measurement. Our new experimental data also determine the position of the triple point.

2. Experimental

High-pressure experiments with in situ X-ray diffraction measurements were conducted at SPring-8 using the multi-anvil apparatus (SPEED-1500) [*Utsumi et al.*, 1998]. The same experimental techniques were applied as *Hirose et al.* [2001]. We used a high-pressure 8/3 assembly with a cylindrical Re-heater (25- μm thickness), similar to that described in *Bertka and Fei* [1997]. Temperatures were monitored by the W5%Re-W26%Re thermocouple. The anhydrous glass starting material with MgSiO_3 composition was loaded near the thermocouple. X-ray diffractions were collected from a $50 \times 100 \mu\text{m}$ area of the pressure marker (mixture of Au or Pt and MgO) at high pressure and high temperature. Its position was precisely chosen using the CCD image looking through the anvil gap (Figure 1). The temperature difference over the entire region of the pressure marker was estimated to be less than 30°C at 2000°C assuming no pressure gradient. This temperature gradient could cause the uncertainty in pressure calculation less than 0.2 GPa using P - V - T equation of state. The primary pressure scales used here are the P - V - T equation of state of gold proposed by *Anderson et al.* [1989] and of platinum by *Jamieson et al.* [1982] (Table 1). The diffraction lines of Au (111, 200, 220,




Figure 1. X-ray CCD image of the assembly at 21 GPa. The white square represents the area of $50 \times 100\text{-}\mu\text{m}$, from which X-ray diffraction was collected. Both gold and platinum were closely loaded but separated by a thin layer of MgO to avoid metal alloying.

Copyright 2001 by the American Geophysical Union.

Paper number 2001GL013549.
0094-8276/01/2001GL013549\$05.00

Table 1. Experimental conditions and results

Run no.	Temperature (°C)	duration (min)	Pressure (GPa)				Run product
			Au*		Pt**		
			initial	final	initial	final	
1	1400	60	20.701	20.226	20.147	19.688	Ak
2	1400	60	21.350	20.994	20.646	20.290	Ak+Pv(trace)
3	1850	30	20.155	19.895			Ak+Pv
4	1850	30	20.659	20.254			Pv
5	1900	15	19.979	19.794			Ak
6	2000	4	20.02***		19.174	n.d.	Mj
7	2000	10	20.34***		19.248	19.057	Pv

*Au scale by *Anderson et al.* [1989].


**Pt scale by *Jamieson et al.* [1982].

***Pressure was estimated by load-pressure relationship (Figure 4).

Ak, akimotoite; Pv, perovskite; Mj, majorite.

311) or Pt (111, 200, 220, 311, 222) were used to calculate the unit cell volumes. The precision of the pressure determination was normally within 0.1-0.2 GPa. The sharpening of diffraction peaks and the consistency of unit-cell parameter calculated from each diffraction line of gold showed that the deviatoric stress approached zero at high temperature above 800-1000°C [*e.g.*, *Weidner et al.*, 1992 and 1998].


To avoid kinetic problems, we took the glass sample to a *P-T* condition of interest, and kept that condition (constant load force of the apparatus and temperature) for 4-60 mins while repeatedly taking diffraction patterns of the pressure maker. Pressure dropped by 0.18-0.48 GPa during heating (Table 1). Then, we quenched the sample to room temperature immediately by turning off the power, so that we could examine the quenched sample. The experimental procedure adopted here allowed us to obtain only one data point for each high *P-T* run, unlike other in situ measurements that can obtain multiple data points in a single run. Phase identification of the quenched sample was made by micro-Raman spectroscopy measurements (Figure 2). The micro-probe analysis showed that the bulk of the sample showed

**Figure 2.** Examples of Raman spectra of the recovered samples.

no evidence of chemical reaction with the surrounding MgO or Al₂O₃.

3. Phase boundaries in MgSiO₃

Seven runs were conducted to determine the phase transition boundaries. The results are shown in Figure 3 based on Anderson's gold pressure scale. In the two 2000°C-runs, only platinum was loaded as a pressure standard, because we had lost the diffraction peaks of gold during the temper-

**Figure 3.** Phase transition boundaries in MgSiO₃ based on Anderson's gold scale [*Anderson et al.*, 1989]. Open symbols, perovskite (Pv); solid symbols, akimotoite (Ak); open symbol with cross, majorite (Mj). The half-filled squares show the coexistence of akimotoite and perovskite. The squares represent the data points obtained in this study. Bars represent initial pressure after reaching a temperature of interest and final pressure before quenching (Table 1). The circles are from [*Ono et al.*, 2001]. Pressures in the runs at 2000°C (gray symbols) were estimated from the calibration curve (Figure 4). Errors in these two runs are from the reproducibility of pressure at a given load force of the apparatus.

ature increase to 2000°C due to grain coarsening near the melting temperature. For these two runs conducted at 650 and 680-ton, pressures in Anderson's gold scale were best estimated from the pressure calibration curve at each temperature (pressure as a function of load force and temperature) (Figure 4). They were established by the total of eight runs we have made at 1750-2000°C using the same high-pressure assembly with the multi-anvil apparatus at SPring-8 [Hirose *et al.*, 2001].

The precise phase boundary was not defined in the previous in situ X-ray observations by Kato *et al.* [1985] and Kuroda *et al.* [2000]. And the Clapeyron slope was not well determined by Ono *et al.* [2001], because the phase transition was observed only in the limited temperature range at 1000-1200°C. Our new experimental data obtained at 1400-2000°C, and together with those by Ono *et al.* [2001], tightly constrain the akimotoite-perovskite transition boundary. The boundary shown in Figure 3 is represented by a linear equation P (GPa) = 25.09 - 0.0027 × T (°C) based on Anderson's gold pressure scale.

The Clapeyron slope of the majorite (tetragonal garnet) - perovskite transition boundary was not precisely determined in the present experiments. A thermodynamical calculation using the physical parameters shown by Akaogi and Ito [1999] gives the slope of 0.0013 GPa/°C for this boundary. If we take this value in Figure 3, the position of the triple point should be at 20.0 GPa and 1920°C.

The phase transition pressures were determined also by the platinum pressure standard (Figure 5). The simultaneous measurement of both the metal pressure standards in a single run at 1400°C showed that Jamieson's platinum scale gives 0.55-0.70 GPa lower pressures than Anderson's gold scale (Table 1) [Hirose *et al.*, 2001].

The choice of equation of state, as well as the choice of pressure standard, could lead to a significant difference in the determination of phase transition pressures by the in situ X-ray diffraction measurements (Figure 5). The phase




Figure 4. Pressure calibration curve for the SPEED-1500 at SPring-8 at high temperatures, based on Anderson's gold pressure scale. Only initial pressures were used for calibration.




Figure 5. Phase transition boundaries in MgSiO₃ based on the different pressure standards and the equations of state. The circles represent the data points obtained in this study, using platinum as a pressure standard. The symbols as in Figure 3. The boundary determined by quench experiments [Ito and Takahashi, 1989] is also shown.

transition pressures in MgSiO₃ were recalculated using a different equation of state of gold [Jamieson *et al.*, 1982] and platinum [Holmes *et al.*, 1989], respectively, from the same unit cell volume data. The akimotoite-perovskite transition pressure based on Jamieson's gold equation of state is 2 GPa higher than that from Anderson's at 1600°C, and is consistent with that determined by conventional quench experiments [Ito and Takahashi, 1989]. It was, however, demonstrated that Jamieson's gold pressure scale deviates widely from other pressure scales such as NaCl, Pt, W, and Mo [Hirose *et al.*, 2001]. On the other hand, pressures calculated from both the platinum equations of state are relatively similar. Holmes's equation gives 0.6 GPa higher pressure at 1600°C. Shim *et al.* [2001] suggested from laser-heated diamond cell experiments that Holmes's platinum scale gives 2 GPa higher pressures than Anderson's gold scale in a similar pressure range, which may explain their different results for the post-spinel phase transition pressure from those by Irfune *et al.* [1998]. The present simultaneous measurements of both the pressure standards by the multi-anvil apparatus, however, showed that the platinum scale either by Holmes *et al.* [1989] or Jamieson *et al.* [1982] predicts even lower pressures than Anderson's gold scale.

Acknowledgments. We thank S. Ono, Y. Nishihara, G. Helffrich and J. Ando for helpful discussions. The comments by anonymous reviewers greatly improved the manuscript. The in situ X-ray experiments were carried out at SPring-8 (2000A0238-ND-np and 2001A0290-CD-np).

References

- Akaogi, M., and E. Ito, Calorimetric study on majorite-perovskite transition in the system Mg₄Si₄O₁₂ - Mg₃Al₂Si₃O₁₂; transition boundaries with positive pressure-temperature slopes, *Phys. Earth Planet. Inter.*, 114, 129-140, 1999.

- Anderson, O.L., D. G. Isaak, and S. Yamamoto, Anharmonicity and the equation of state for gold, *J. Appl. Phys.*, *65*, 1534-1543, 1989.
- Bertka, C.M., and Y. Fei, Mineralogy of Martian interior up to core-mantle boundary pressures, *J. Geophys. Res.*, *102*, 5251-5264, 1997.
- Hirose, K., Y. Fei, S. Ono, T. Yagi, and K. Funakoshi, In situ measurements of the phase transition boundary in $Mg_3Al_2Si_3O_{12}$: implications for the nature of the seismic discontinuities in the Earth's mantle, *Earth Planet. Sci. Lett.*, *184*, 567-573, 2001.
- Holmes, N. C., J. A. Moriarty, G. R. Gathers, and W. J. Nellis, The equation of state of platinum to 660 GPa (6.6 MBar), *J. Appl. Phys.*, *66*, 2962-2967, 1989.
- Irifune, T., N. Nishiyama, K. Kuroda, et al., The postspinel phase boundary in Mg_2SiO_4 determined by in situ X-ray diffraction, *Science*, *279*, 1698-1700, 1998.
- Ito, E., and E. Takahashi, Postspinel transformations in the system Mg_2SiO_4 - Fe_2SiO_4 and some geophysical implications, *J. Geophys. Res.*, *94*, 10637-10646, 1989.
- Jamieson, J. C., J. N. Fritz, and M. H. Manghnani, Pressure measurement at high temperature in x-ray diffraction studies: gold as a primary standard, in *High-Pressure Research in Geophysics*, edited by S. Akimoto and M.H. Manghnani, pp. 27-48, CAPJ, Tokyo, 1982.
- Kato, T, E. Ohtani, H. Morishima, et al., In situ X-ray observation of high-pressure phase transitions of $MgSiO_3$ and thermal expansion of $MgSiO_3$ perovskite at 25 GPa by double-stage multianvil system, *J. Geophys. Res.*, *100*, 20475-20481, 1995.
- Kuroda, K., T. Irifune, T. Inoue, et al., Determination of the phase boundary between ilmenite and perovskite in $MgSiO_3$ by in situ X-ray diffraction and quench experiments, *Phys. Chem. Mine.*, *27*, 523-532, 2000.
- Ono, S., T. Katsura, E. Ito, et al., In situ observation of ilmenite-perovskite phase transition in $MgSiO_3$ using synchrotron radiation, *Geophys. Res. Lett.*, *28*, 835-838, 2001.
- Shim, S-H, T. S. Duffy, and G. Shen, Evidence that the post-spinel transformation in Mg_2SiO_4 is responsible for the 660-km seismic discontinuity, *Nature*, *411*, 571-574, 2001.
- Utsumi, W., K. Funakoshi, S. Urakawa, et al., SPring-8 beamlines for high pressure science with multi-anvil apparatus, *Rev. High Press. Sci. Technol.*, *7*, 1484-1486, 1998.
- Weidner, D. J., M. T. Vaughan, J. Jo and Y. Wang, Characterization of stress, pressure, and temperature in SAM85, A diamond type high pressure apparatus, in *High-Pressure Research: Application to Earth and Planetary Sciences*, edited by Y. Shono and M.H. Manghnani, pp. 13-17, Terrapub/AGU, 1992.
- Weidner, D. J., Y. Wang, G. Chen, Rheology measurements at high pressure and temperature, in *Properties of Earth and Planetary Materials at High Pressure and Temperature*, edited by M.H. Manghnani and T. Yagi, pp. 473-482, Geophysical Monograph, 101, AGU, 1998.

K. Funakoshi, Japan Synchrotron Radiation Research Institute, Mikazuki-cho, Hyogo 679-5198, Japan.

K. Hirose, T. Komabayashi, and M. Murakami, Department of Earth and Planetary Sciences, Tokyo Institute of Technology, 2-12-1 Ookayama, Meguro, Tokyo 152-8551, Japan (e-mail: kei@geo.titech.ac.jp).

(Received May 30, 2001; revised September 6, 2001; accepted September 12, 2001.)