

Supporting Information to:

Antioxidant, Antimalarial and Antimicrobial Activities of Tannin-Rich Fractions, Ellagitannins and Phenolic Acids from *Punica granatum*

Muntha K. Reddy¹

Sashi K. Gupta¹

Melissa R. Jacob²

Shabana I. Khan²

Daneel Ferreira^{1,2}

Affiliation

¹ Department of Pharmacognosy, School of Pharmacy, University, The University of Mississippi, MS, USA

² National Center for Natural Products Research, Research Institute of Pharmaceutical Sciences, School of Pharmacy, University, The University of Mississippi, MS, USA

Correspondence

Dr. D. Ferreira

Department of Pharmacognosy and National Center for Natural Products Research

Research Institute of Pharmaceutical Sciences

School of Pharmacy

The University of Mississippi

University

MS 38677

USA

Phone: +1-662-915-7026

Fax: +1-662-915-6975

E-mail: dferreir@olemiss.edu




Fig. 1S Effect of TPT on ROS inhibitory activity.




Fig. 2S Effects of XAD-EtOAc, XAD-BuOH, XAD-H₂O, XAD-PJ, ellagic acid (1), gallagic acid (2), punicalins (3), and punicalagins (4) on ROS inhibitory activity. All sample solutions were made as µg/mL concentrations. The antioxidant dose dependent effect was measured by plotting graphs (concentration of test solutions vs. production of DCF).