

Supplementary Material S1 Brazilian versions of outcome measures used in stroke rehabilitation and respective levels of assessment

Outcome measures	Level of assessment
National Institutes of Health Stroke Scale ¹	Stroke deficit
Modified Rankin Scale ¹	Global disability
Barthel index ¹	Activities of daily living
Hospital Anxiety and Depression Scale ²	Depression
Montreal-Toulouse Language Assessment Battery ³	Language
Cancelation and Bisection Tests ⁴	Unilateral spatial neglect
Catherine Bergego Scale ⁵	Unilateral spatial neglect
Mini-mental State Examination ⁶	Cognition
Montreal Cognitive Assessment ⁷	Cognition
Fugl-Meyer scale ⁸	Motor-somatosensory
Nottingham Sensory Assessment ⁹	Somatosensory
Jebsen-Taylor test ¹⁰	Activities of the upper limbs
Motor Activity Log-Brazil (MAL) ¹¹	Activities of the upper limbs
Wolf-motor Function Test ¹²	Activities of the upper limbs
Test d'Évaluation des Membres Supérieurs de Personnes Âgées (TEMPA) ¹³	Activities of the upper limbs
Berg Balance Scale ¹⁴	Balance
Balance Evaluation Systems Test (BEST) ¹⁵	Balance
Activities-specific balance confidence scale ¹⁶	Balance
Dynamic Gait Index ¹⁷	Gait and balance
Functional Ambulation Category ¹⁸	Gait
Trunk Impairment Scale ¹⁹	Trunk control
Postural Assessment Scale for Stroke Patients ²⁰	Trunk control
Self-efficacy Questionnaire ²¹	Activities of daily living
SATIS-Stroke questionnaire ²²	Quality of life
Stroke impact scale 3.0 ²³	Healthy status/Quality of life

Conflict of Interests

The authors have no conflict of interests to declare.

References

- Cincura C, Pontes-Neto OM, Neville IS, et al. Validation of the National Institutes of Health Stroke Scale, modified Rankin Scale and Barthel Index in Brazil: the role of cultural adaptation and structured interviewing. *Cerebrovasc Dis* 2009;27(02):119–122
- Pais-Ribeiro J, Silva I, Ferreira T, Martins A, Meneses R, Baltar M. Validation study of a Portuguese version of the Hospital Anxiety and Depression Scale. *Psychol Health Med* 2007;12(02):225–235, quiz 235–237. Doi: 10.1080/13548500500524088
- Pagliarin KC, Ortiz KZ, Barreto SdosS, et al. Montreal-Toulouse Language Assessment Battery: evidence of criterion validity from patients with aphasia. *J Neurol Sci* 2015;357(1-2):246–251
- Luvizutto GJ, Fogaroli MO, Theotonio RM, Moura Neto E, Nunes HRC, Bazan R. Norm scores of cancelation and bisection tests for unilateral spatial neglect: data from a Brazilian population. *Clinics (São Paulo)* 2020;75:e1468
- Marques CLS, de Souza JT, Gonçalves MG, et al. Validation of the Catherine Bergego Scale in patients with unilateral spatial neglect after stroke. *Dement Neuropsychol* 2019;13(01):82–88
- Brucki SM, Nitrini R, Caramelli P, Bertolucci PH, Okamoto IH. Sugestões para o uso do mini-exame do estado mental no Brasil. [Suggestions for utilization of the mini-mental state examination in Brazil]. *Arq Neuropsiquiatr*. 2003 Sep;61(3B):777–81. Portuguese. doi: Doi: 10.1590/s0004-282 × 2003000500014
- Memória CM, Yassuda MS, Nakano EY, Forlenza OV. Brief screening for mild cognitive impairment: validation of the Brazilian version of the Montreal cognitive assessment. *Int J Geriatr Psychiatry* 2013;28(01):34–40. Doi: 10.1002/gps.3787
- Michaelsen SM, Rocha AS, Knabben RJ, Rodrigues LP, Fernandes CG. Translation, adaptation and inter-rater reliability of the administration manual for the Fugl-Meyer assessment. *Rev Bras Fisioter* 2011;15(01):80–88
- Lima DH, Queiroz AP, De Salvo G, Yoneyama SM, Oberg TD, Lima NM. Brazilian version of the Nottingham Sensory Assessment: validity, agreement and reliability. *Rev Bras Fisioter* 2010;14(02):166–174

- 10 Ferreiro KN, Santos RL, Conforto AB. Psychometric properties of the portuguese version of the Jebsen-Taylor test for adults with mild hemiparesis. *Rev Bras Fisioter*. 2010 Sep-Oct;14(05):377–82.. Doi: 10.1590/s1413-35552010005000018
- 11 Pereira ND, Ovando AC, Michaelsen SM, et al. Motor Activity Log-Brazil: reliability and relationships with motor impairments in individuals with chronic stroke. *Arq Neuropsiquiatr* 2012;70(03):196–201. Doi: 10.1590/s0004-282 × 2012000300008
- 12 Pereira ND, Michaelsen SM, Menezes IS, Ovando AC, Lima RC, Teixeira-Salmela LF. Reliability of the Brazilian version of the Wolf Motor Function Test in adults with hemiparesis. *Rev Bras Fisioter* 2011;15(03):257–265
- 13 Michaelsen SM, Natalio MA, Silva AG, Pagnussat AS. Confiabilidade da tradução e adaptação do test d'évaluation des membres supérieurs de personnes âgées (TEMPA) para o português e validação para adultos com hemiparesia. *Rev Bras Fisioter*. 2008;12(06):511–519
- 14 Miyamoto ST, Lombardi Junior I, Berg KO, Ramos LR, Natour J. Brazilian version of the Berg balance scale. *Braz J Med Biol Res* 2004;37(09):1411–1421
- 15 Rodrigues LC, Marques AP, Barros PB, Michaelsen SM. Reliability of the Balance Evaluation Systems Test (BESTest) and BESTest sections for adults with hemiparesis. *Braz J Phys Ther* 2014;18(03):276–281. Doi: 10.1590/bjpt-rbf.2014.0033
- 16 Marques AP, Mendes YC, Taddei U, Pereira CA, Assumpção A. Brazilian-Portuguese translation and cross cultural adaptation of the activities-specific balance confidence (ABC) scale. *Braz J Phys Ther* 2013;17(02):170–178
- 17 De Castro SM, Perracini MR, Ganança FF. Dynamic Gait Index–Brazilian version. *Rev Bras Otorrinolaringol (Engl Ed)* 2006;72(06):817–825. Doi: 10.1016/s1808-8694(15)31050-8
- 18 Elord C, Corrêa FI, Pereira GS, Silva SM, Corrêa JC. Translation into Brazilian Portuguese, cross-cultural adaptation, reliability and validation of the Functional Ambulation Classification for the categorization of ambulation following a stroke in a clinical setting. *Rev Neurol* 2020;70(10):365–371. Doi: 10.33588/rn.7010.2019400
- 19 Lima NMFV, Rodrigues SY, Fillipo TM, de Oliveira R, Oberg TD, Cacho EWA. Brazilian version of the Trunk Impairment Scale: a reliability study in post-stroke subjects. *Fisioter Pesqui* 2008;15(03):248–253
- 20 Yoneyama SM, Roiz RM, Oliveira TM, Oberg TD, Lima NMFV. [Validação da versão brasileira da Escala de Avaliação Postural para Pacientes após Acidente Vascular Encefálico]. *Acta Fisiatr* 2008;15(02):96–100
- 21 Makhoul MP, Pinto EB, Mazzini NA, Winstein C, Torriani-Pasin C. Translation and validation of the stroke self-efficacy questionnaire to a Portuguese version in stroke survivors. *Top Stroke Rehabil* 2020;27(06):462–472
- 22 Pereira GS, Corrêa FI, Júlio CE, et al. Clinimetric properties of the SATIS-Stroke questionnaire in the Brazilian population: A satisfaction assessment measure addressing activities and participation after a stroke. *Braz J Phys Ther* 2021;25(06):719–726
- 23 Carod-Artal FJ, Coral LF, Trizotto DS, Moreira CM. The stroke impact scale 3.0: evaluation of acceptability, reliability, and validity of the Brazilian version. *Stroke* 2008;39(09):2477–2484